



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07H 21/02, 21/04, C12N 5/00, 5/04, 5/06, 5/10, 5/16, 15/00, 15/09, 15/10, 15/11, 15/12, C12P 21/04, 21/06</p>	A1	<p>(11) International Publication Number: WO 98/56804</p> <p>(43) International Publication Date: 17 December 1998 (17.12.98)</p>																																																																																						
<table style="width: 100%; border: none;"> <tr> <td style="width: 45%; vertical-align: top; padding: 5px;"> <p>(21) International Application Number: PCT/US98/12125</p> <p>(22) International Filing Date: 11 June 1998 (11.06.98)</p> <p>(30) Priority Data:</p> <table style="width: 100%; border: none;"> <tr><td style="width: 30%;">60/049,547</td><td style="width: 40%;">13 June 1997 (13.06.97)</td><td style="width: 30%;">US</td></tr> <tr><td>60/049,548</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,549</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,550</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/050,566</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,606</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,607</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,608</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,609</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,610</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,611</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/050,901</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/052,989</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/051,919</td><td>8 July 1997 (08.07.97)</td><td>US</td></tr> <tr><td>60/055,984</td><td>18 August 1997 (18.08.97)</td><td>US</td></tr> <tr><td>60/058,665</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/058,668</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/058,669</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/058,750</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/058,971</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/058,972</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/058,975</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/060,834</td><td>2 October 1997 (02.10.97)</td><td>US</td></tr> <tr><td>60/060,841</td><td>2 October 1997 (02.10.97)</td><td>US</td></tr> <tr><td>60/060,844</td><td>2 October 1997 (02.10.97)</td><td>US</td></tr> <tr><td>60/060,865</td><td>2 October 1997 (02.10.97)</td><td>US</td></tr> <tr><td>60/061,059</td><td>2 October 1997 (02.10.97)</td><td>US</td></tr> <tr><td>60/061,060</td><td>2 October 1997 (02.10.97)</td><td>US</td></tr> </table> </td> <td style="width: 55%; vertical-align: top; padding: 5px;"> <p>(71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): MOORE, Paul, A. [GB/US]; Apartment 104, 1908 Holly Ridge Drive, McLean, VA 22102 (US). SHI, Yanggu [CN/US]; 437 West Side Drive, Gaithersburg, MD 20878 (US). ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US). RUBEN, Steven, M. [US/US]; 18528 Heritage Hills Drive, Olney, MD 20832 (US). LAFLEUR, David, W. [US/US]; 1615 Q Street, N.W. #807, Washington, DC 20009 (US). OLSEN, Henrik, S. [DK/US]; 182 Kendrick Place #24, Gaithersburg, MD 20878 (US). EBNER, Reinhard [DE/US]; 9906 Shelburne Terrace #316, Gaithersburg, MD 20878 (US). BREWER, Laurie, A. [US/US]; 14920 Mt. Nebo Road, Poolesville, MD 20837 (US). YOUNG, Paul [US/US]; 122 Beckwith Street, Gaithersburg, MD 20878 (US). GREENE, John, M. [US/US]; 872 Diamond Drive, Gaithersburg, MD 20878 (US). FERRIE, Ann, M. [US/US]; 13202 L Astoria Hill Court, Germantown, MD 20874 (US). YU, Guo-Liang [CN/US]; 13524 Straw Bale Lane, Darnestown, MD 20878 (US). NI, Jian [CN/US]; 5502 Manorfield Road, Rockville, MD 20853 (US). FENG, Ping [CN/US]; 4 Relda Court, Gaithersburg, MD 20878 (US).</p> <p>(74) Agents: BROOKES, A., Anders et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).</p> <p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p style="text-align: center;">Published With international search report.</p> </td> </tr> </table>			<p>(21) International Application Number: PCT/US98/12125</p> <p>(22) International Filing Date: 11 June 1998 (11.06.98)</p> <p>(30) Priority Data:</p> <table style="width: 100%; border: none;"> <tr><td style="width: 30%;">60/049,547</td><td style="width: 40%;">13 June 1997 (13.06.97)</td><td style="width: 30%;">US</td></tr> <tr><td>60/049,548</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,549</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,550</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/050,566</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,606</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,607</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,608</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,609</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,610</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,611</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/050,901</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/052,989</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/051,919</td><td>8 July 1997 (08.07.97)</td><td>US</td></tr> <tr><td>60/055,984</td><td>18 August 1997 (18.08.97)</td><td>US</td></tr> <tr><td>60/058,665</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/058,668</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/058,669</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/058,750</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/058,971</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/058,972</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/058,975</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/060,834</td><td>2 October 1997 (02.10.97)</td><td>US</td></tr> <tr><td>60/060,841</td><td>2 October 1997 (02.10.97)</td><td>US</td></tr> <tr><td>60/060,844</td><td>2 October 1997 (02.10.97)</td><td>US</td></tr> <tr><td>60/060,865</td><td>2 October 1997 (02.10.97)</td><td>US</td></tr> <tr><td>60/061,059</td><td>2 October 1997 (02.10.97)</td><td>US</td></tr> <tr><td>60/061,060</td><td>2 October 1997 (02.10.97)</td><td>US</td></tr> </table>	60/049,547	13 June 1997 (13.06.97)	US	60/049,548	13 June 1997 (13.06.97)	US	60/049,549	13 June 1997 (13.06.97)	US	60/049,550	13 June 1997 (13.06.97)	US	60/050,566	13 June 1997 (13.06.97)	US	60/049,606	13 June 1997 (13.06.97)	US	60/049,607	13 June 1997 (13.06.97)	US	60/049,608	13 June 1997 (13.06.97)	US	60/049,609	13 June 1997 (13.06.97)	US	60/049,610	13 June 1997 (13.06.97)	US	60/049,611	13 June 1997 (13.06.97)	US	60/050,901	13 June 1997 (13.06.97)	US	60/052,989	13 June 1997 (13.06.97)	US	60/051,919	8 July 1997 (08.07.97)	US	60/055,984	18 August 1997 (18.08.97)	US	60/058,665	12 September 1997 (12.09.97)	US	60/058,668	12 September 1997 (12.09.97)	US	60/058,669	12 September 1997 (12.09.97)	US	60/058,750	12 September 1997 (12.09.97)	US	60/058,971	12 September 1997 (12.09.97)	US	60/058,972	12 September 1997 (12.09.97)	US	60/058,975	12 September 1997 (12.09.97)	US	60/060,834	2 October 1997 (02.10.97)	US	60/060,841	2 October 1997 (02.10.97)	US	60/060,844	2 October 1997 (02.10.97)	US	60/060,865	2 October 1997 (02.10.97)	US	60/061,059	2 October 1997 (02.10.97)	US	60/061,060	2 October 1997 (02.10.97)	US	<p>(71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): MOORE, Paul, A. [GB/US]; Apartment 104, 1908 Holly Ridge Drive, McLean, VA 22102 (US). SHI, Yanggu [CN/US]; 437 West Side Drive, Gaithersburg, MD 20878 (US). ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US). RUBEN, Steven, M. [US/US]; 18528 Heritage Hills Drive, Olney, MD 20832 (US). LAFLEUR, David, W. [US/US]; 1615 Q Street, N.W. #807, Washington, DC 20009 (US). OLSEN, Henrik, S. [DK/US]; 182 Kendrick Place #24, Gaithersburg, MD 20878 (US). EBNER, Reinhard [DE/US]; 9906 Shelburne Terrace #316, Gaithersburg, MD 20878 (US). BREWER, Laurie, A. [US/US]; 14920 Mt. Nebo Road, Poolesville, MD 20837 (US). YOUNG, Paul [US/US]; 122 Beckwith Street, Gaithersburg, MD 20878 (US). GREENE, John, M. [US/US]; 872 Diamond Drive, Gaithersburg, MD 20878 (US). FERRIE, Ann, M. [US/US]; 13202 L Astoria Hill Court, Germantown, MD 20874 (US). YU, Guo-Liang [CN/US]; 13524 Straw Bale Lane, Darnestown, MD 20878 (US). NI, Jian [CN/US]; 5502 Manorfield Road, Rockville, MD 20853 (US). FENG, Ping [CN/US]; 4 Relda Court, Gaithersburg, MD 20878 (US).</p> <p>(74) Agents: BROOKES, A., Anders et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).</p> <p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p style="text-align: center;">Published With international search report.</p>
<p>(21) International Application Number: PCT/US98/12125</p> <p>(22) International Filing Date: 11 June 1998 (11.06.98)</p> <p>(30) Priority Data:</p> <table style="width: 100%; border: none;"> <tr><td style="width: 30%;">60/049,547</td><td style="width: 40%;">13 June 1997 (13.06.97)</td><td style="width: 30%;">US</td></tr> <tr><td>60/049,548</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,549</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,550</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/050,566</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,606</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,607</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,608</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,609</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,610</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/049,611</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/050,901</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/052,989</td><td>13 June 1997 (13.06.97)</td><td>US</td></tr> <tr><td>60/051,919</td><td>8 July 1997 (08.07.97)</td><td>US</td></tr> <tr><td>60/055,984</td><td>18 August 1997 (18.08.97)</td><td>US</td></tr> <tr><td>60/058,665</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/058,668</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/058,669</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/058,750</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/058,971</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/058,972</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/058,975</td><td>12 September 1997 (12.09.97)</td><td>US</td></tr> <tr><td>60/060,834</td><td>2 October 1997 (02.10.97)</td><td>US</td></tr> <tr><td>60/060,841</td><td>2 October 1997 (02.10.97)</td><td>US</td></tr> <tr><td>60/060,844</td><td>2 October 1997 (02.10.97)</td><td>US</td></tr> <tr><td>60/060,865</td><td>2 October 1997 (02.10.97)</td><td>US</td></tr> <tr><td>60/061,059</td><td>2 October 1997 (02.10.97)</td><td>US</td></tr> <tr><td>60/061,060</td><td>2 October 1997 (02.10.97)</td><td>US</td></tr> </table>	60/049,547	13 June 1997 (13.06.97)	US	60/049,548	13 June 1997 (13.06.97)	US	60/049,549	13 June 1997 (13.06.97)	US	60/049,550	13 June 1997 (13.06.97)	US	60/050,566	13 June 1997 (13.06.97)	US	60/049,606	13 June 1997 (13.06.97)	US	60/049,607	13 June 1997 (13.06.97)	US	60/049,608	13 June 1997 (13.06.97)	US	60/049,609	13 June 1997 (13.06.97)	US	60/049,610	13 June 1997 (13.06.97)	US	60/049,611	13 June 1997 (13.06.97)	US	60/050,901	13 June 1997 (13.06.97)	US	60/052,989	13 June 1997 (13.06.97)	US	60/051,919	8 July 1997 (08.07.97)	US	60/055,984	18 August 1997 (18.08.97)	US	60/058,665	12 September 1997 (12.09.97)	US	60/058,668	12 September 1997 (12.09.97)	US	60/058,669	12 September 1997 (12.09.97)	US	60/058,750	12 September 1997 (12.09.97)	US	60/058,971	12 September 1997 (12.09.97)	US	60/058,972	12 September 1997 (12.09.97)	US	60/058,975	12 September 1997 (12.09.97)	US	60/060,834	2 October 1997 (02.10.97)	US	60/060,841	2 October 1997 (02.10.97)	US	60/060,844	2 October 1997 (02.10.97)	US	60/060,865	2 October 1997 (02.10.97)	US	60/061,059	2 October 1997 (02.10.97)	US	60/061,060	2 October 1997 (02.10.97)	US	<p>(71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): MOORE, Paul, A. [GB/US]; Apartment 104, 1908 Holly Ridge Drive, McLean, VA 22102 (US). SHI, Yanggu [CN/US]; 437 West Side Drive, Gaithersburg, MD 20878 (US). ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US). RUBEN, Steven, M. [US/US]; 18528 Heritage Hills Drive, Olney, MD 20832 (US). LAFLEUR, David, W. [US/US]; 1615 Q Street, N.W. #807, Washington, DC 20009 (US). OLSEN, Henrik, S. [DK/US]; 182 Kendrick Place #24, Gaithersburg, MD 20878 (US). EBNER, Reinhard [DE/US]; 9906 Shelburne Terrace #316, Gaithersburg, MD 20878 (US). BREWER, Laurie, A. [US/US]; 14920 Mt. Nebo Road, Poolesville, MD 20837 (US). YOUNG, Paul [US/US]; 122 Beckwith Street, Gaithersburg, MD 20878 (US). GREENE, John, M. [US/US]; 872 Diamond Drive, Gaithersburg, MD 20878 (US). FERRIE, Ann, M. [US/US]; 13202 L Astoria Hill Court, Germantown, MD 20874 (US). YU, Guo-Liang [CN/US]; 13524 Straw Bale Lane, Darnestown, MD 20878 (US). NI, Jian [CN/US]; 5502 Manorfield Road, Rockville, MD 20853 (US). FENG, Ping [CN/US]; 4 Relda Court, Gaithersburg, MD 20878 (US).</p> <p>(74) Agents: BROOKES, A., Anders et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).</p> <p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p style="text-align: center;">Published With international search report.</p>			
60/049,547	13 June 1997 (13.06.97)	US																																																																																						
60/049,548	13 June 1997 (13.06.97)	US																																																																																						
60/049,549	13 June 1997 (13.06.97)	US																																																																																						
60/049,550	13 June 1997 (13.06.97)	US																																																																																						
60/050,566	13 June 1997 (13.06.97)	US																																																																																						
60/049,606	13 June 1997 (13.06.97)	US																																																																																						
60/049,607	13 June 1997 (13.06.97)	US																																																																																						
60/049,608	13 June 1997 (13.06.97)	US																																																																																						
60/049,609	13 June 1997 (13.06.97)	US																																																																																						
60/049,610	13 June 1997 (13.06.97)	US																																																																																						
60/049,611	13 June 1997 (13.06.97)	US																																																																																						
60/050,901	13 June 1997 (13.06.97)	US																																																																																						
60/052,989	13 June 1997 (13.06.97)	US																																																																																						
60/051,919	8 July 1997 (08.07.97)	US																																																																																						
60/055,984	18 August 1997 (18.08.97)	US																																																																																						
60/058,665	12 September 1997 (12.09.97)	US																																																																																						
60/058,668	12 September 1997 (12.09.97)	US																																																																																						
60/058,669	12 September 1997 (12.09.97)	US																																																																																						
60/058,750	12 September 1997 (12.09.97)	US																																																																																						
60/058,971	12 September 1997 (12.09.97)	US																																																																																						
60/058,972	12 September 1997 (12.09.97)	US																																																																																						
60/058,975	12 September 1997 (12.09.97)	US																																																																																						
60/060,834	2 October 1997 (02.10.97)	US																																																																																						
60/060,841	2 October 1997 (02.10.97)	US																																																																																						
60/060,844	2 October 1997 (02.10.97)	US																																																																																						
60/060,865	2 October 1997 (02.10.97)	US																																																																																						
60/061,059	2 October 1997 (02.10.97)	US																																																																																						
60/061,060	2 October 1997 (02.10.97)	US																																																																																						
<p>(54) Title: 86 HUMAN SECRETED PROTEINS</p> <p>(57) Abstract</p> <p>The present invention relates to 86 novel human secreted proteins and isolated nucleic acids containing the coding regions of the genes encoding such proteins. Also provided are vectors, host cells, antibodies, and recombinant methods for producing human secreted proteins. The invention further relates to diagnostic and therapeutic methods useful for diagnosing and treating disorders related to these novel human secreted proteins.</p>																																																																																								

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

86 Human Secreted Proteins

Field of the Invention

This invention relates to newly identified polynucleotides and the polypeptides encoded by these polynucleotides, uses of such polynucleotides and polypeptides, and
5 their production.

Background of the Invention

Unlike bacterium, which exist as a single compartment surrounded by a membrane, human cells and other eucaryotes are subdivided by membranes into many functionally distinct compartments. Each membrane-bounded compartment, or
10 organelle, contains different proteins essential for the function of the organelle. The cell uses "sorting signals," which are amino acid motifs located within the protein, to target proteins to particular cellular organelles.

One type of sorting signal, called a signal sequence, a signal peptide, or a leader sequence, directs a class of proteins to an organelle called the endoplasmic reticulum
15 (ER). The ER separates the membrane-bounded proteins from all other types of proteins. Once localized to the ER, both groups of proteins can be further directed to another organelle called the Golgi apparatus. Here, the Golgi distributes the proteins to vesicles, including secretory vesicles, the cell membrane, lysosomes, and the other organelles.

20 Proteins targeted to the ER by a signal sequence can be released into the extracellular space as a secreted protein. For example, vesicles containing secreted proteins can fuse with the cell membrane and release their contents into the extracellular space - a process called exocytosis. Exocytosis can occur constitutively or after receipt of a triggering signal. In the latter case, the proteins are stored in secretory vesicles (or
25 secretory granules) until exocytosis is triggered. Similarly, proteins residing on the cell membrane can also be secreted into the extracellular space by proteolytic cleavage of a "linker" holding the protein to the membrane.

Despite the great progress made in recent years, only a small number of genes encoding human secreted proteins have been identified. These secreted proteins include
30 the commercially valuable human insulin, interferon, Factor VIII, human growth hormone, tissue plasminogen activator, and erythropoietin. Thus, in light of the pervasive role of secreted proteins in human physiology, a need exists for identifying and characterizing novel human secreted proteins and the genes that encode them. This knowledge will allow one to detect, to treat, and to prevent medical disorders by using
35 secreted proteins or the genes that encode them.

Summary of the Invention

The present invention relates to novel polynucleotides and the encoded polypeptides. Moreover, the present invention relates to vectors, host cells, antibodies, and recombinant methods for producing the polypeptides and polynucleotides. Also provided are diagnostic methods for detecting disorders related to the polypeptides, and therapeutic methods for treating such disorders. The invention further relates to screening methods for identifying binding partners of the polypeptides.

Detailed Description

Definitions

The following definitions are provided to facilitate understanding of certain terms used throughout this specification.

In the present invention, "isolated" refers to material removed from its original environment (e.g., the natural environment if it is naturally occurring), and thus is altered "by the hand of man" from its natural state. For example, an isolated polynucleotide could be part of a vector or a composition of matter, or could be contained within a cell, and still be "isolated" because that vector, composition of matter, or particular cell is not the original environment of the polynucleotide.

In the present invention, a "secreted" protein refers to those proteins capable of being directed to the ER, secretory vesicles, or the extracellular space as a result of a signal sequence, as well as those proteins released into the extracellular space without necessarily containing a signal sequence. If the secreted protein is released into the extracellular space, the secreted protein can undergo extracellular processing to produce a "mature" protein. Release into the extracellular space can occur by many mechanisms, including exocytosis and proteolytic cleavage.

As used herein, a "polynucleotide" refers to a molecule having a nucleic acid sequence contained in SEQ ID NO:X or the cDNA contained within the clone deposited with the ATCC. For example, the polynucleotide can contain the nucleotide sequence of the full length cDNA sequence, including the 5' and 3' untranslated sequences, the coding region, with or without the signal sequence, the secreted protein coding region, as well as fragments, epitopes, domains, and variants of the nucleic acid sequence. Moreover, as used herein, a "polypeptide" refers to a molecule having the translated amino acid sequence generated from the polynucleotide as broadly defined.

In the present invention, the full length sequence identified as SEQ ID NO:X was often generated by overlapping sequences contained in multiple clones (contig

analysis). A representative clone containing all or most of the sequence for SEQ ID NO:X was deposited with the American Type Culture Collection ("ATCC"). As shown in Table 1, each clone is identified by a cDNA Clone ID (Identifier) and the ATCC Deposit Number. The ATCC is located at 10801 University Boulevard,
5 Manassas, Virginia 20110-2209, USA. The ATCC deposit was made pursuant to the terms of the Budapest Treaty on the international recognition of the deposit of microorganisms for purposes of patent procedure.

A "polynucleotide" of the present invention also includes those polynucleotides capable of hybridizing, under stringent hybridization conditions, to sequences contained
10 in SEQ ID NO:X, the complement thereof, or the cDNA within the clone deposited with the ATCC. "Stringent hybridization conditions" refers to an overnight incubation at 42°C in a solution comprising 50% formamide, 5x SSC (750 mM NaCl, 75 mM sodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the
15 filters in 0.1x SSC at about 65°C.

Also contemplated are nucleic acid molecules that hybridize to the polynucleotides of the present invention at lower stringency hybridization conditions. Changes in the stringency of hybridization and signal detection are primarily accomplished through the manipulation of formamide concentration (lower percentages
20 of formamide result in lowered stringency); salt conditions, or temperature. For example, lower stringency conditions include an overnight incubation at 37°C in a solution comprising 6X SSPE (20X SSPE = 3M NaCl; 0.2M NaH₂PO₄; 0.02M EDTA, pH 7.4), 0.5% SDS, 30% formamide, 100 µg/ml salmon sperm blocking DNA; followed by washes at 50°C with 1XSSPE, 0.1% SDS. In addition, to achieve even
25 lower stringency, washes performed following stringent hybridization can be done at higher salt concentrations (e.g. 5X SSC).

Note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to suppress background in hybridization experiments. Typical blocking reagents include
30 Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above, due to problems with compatibility.

Of course, a polynucleotide which hybridizes only to polyA+ sequences (such
35 as any 3' terminal polyA+ tract of a cDNA shown in the sequence listing), or to a

complementary stretch of T (or U) residues, would not be included in the definition of "polynucleotide," since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone).

5 The polynucleotide of the present invention can be composed of any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. For example, polynucleotides can be composed of single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be
10 single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, the polynucleotide can be composed of triple-stranded regions comprising RNA or DNA or both RNA and DNA. A polynucleotide may also contain one or more modified bases or DNA or RNA backbones modified for stability
15 or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications can be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically, or metabolically modified forms.

 The polypeptide of the present invention can be composed of amino acids joined
20 to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres, and may contain amino acids other than the 20 gene-encoded amino acids. The polypeptides may be modified by either natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs,
25 as well as in a voluminous research literature. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be
30 branched, for example, as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched, and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a
35 nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine,

formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993); POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, pgs. 1-12 (1983); Seifter et al., Meth Enzymol 182:626-646 (1990); Rattan et al., Ann NY Acad Sci 663:48-62 (1992).)

"SEQ ID NO:X" refers to a polynucleotide sequence while "SEQ ID NO:Y" refers to a polypeptide sequence, both sequences identified by an integer specified in Table 1.

"A polypeptide having biological activity" refers to polypeptides exhibiting activity similar, but not necessarily identical to, an activity of a polypeptide of the present invention, including mature forms, as measured in a particular biological assay, with or without dose dependency. In the case where dose dependency does exist, it need not be identical to that of the polypeptide, but rather substantially similar to the dose-dependence in a given activity as compared to the polypeptide of the present invention (i.e., the candidate polypeptide will exhibit greater activity or not more than about 25-fold less and, preferably, not more than about tenfold less activity, and most preferably, not more than about three-fold less activity relative to the polypeptide of the present invention.)

Polynucleotides and Polypeptides of the Invention

FEATURES OF PROTEIN ENCODED BY GENE NO: 1

The translation product of this gene shares sequence homology with LIM-homeobox domain proteins, such as T-cell translocation protein, which are thought to be important in development and leukemogenesis. In addition, translation product of this gene shares homology with the human breast tumor autoantigen (See Accession No. gil1914877). In one embodiment the polypeptides of the invention comprise the sequence:

MNGSHKDPLLPPASARTPSLPPAPPAQAPLPWKPSGFARISPPPPLAILQYRG
KADHGESGQQLAAAPGDGRLPLLEAVRRLRGQDCGPLSALCHGQLLAQVPVQ
VLLPGAXGDIGTSCYTKSGMILCRNDYIRLFGNSGACSACGQSIPASELVMRA
QGNVYHLKCFTCTCRNRLVPGDRFHYINGSLFCEHDRPTALINGHLNSLQSN

PLLPDQKVCKVRVMQNACLHLRFVHHRWIPCXFSRQVTFVASTSASSMPLHLL
 (SEQ ID NO:211); MARTRTPSSPFLLLRELPPSLQLRQPRRPFGSRAASLAFHRR
 RLSQYCNIGEKQTMVNP GSSSQPPV TAGSLSWKRCAGCGGKIADRFLLYA
 (SEQ ID NO:212); LFGNSGACSACGQSIPASELVMRA (SEQ ID NO:213);
 5 HDRPTALINGHLNSLQSNP (SEQ ID NO:214); and/or LVPGDRFH YING (SEQ ID
 NO:215). Polynucleotide fragments encoding these polypeptide fragments are also
 encompassed by the invention.

This gene is expressed primarily in fetal brain, osteosarcoma, IL-1/TNF treated
 synovial, and estradiol treated endometrial stromal cells, and to a lesser extent in
 10 chondrosarcoma, smooth muscle and number of other tissues.

Therefore, polynucleotides and polypeptides of the invention are useful as
 reagents for differential identification of the tissue(s) or cell type(s) present in a
 biological sample and for diagnosis of diseases and conditions which include, but are
 not limited to, developmental defects or leukemia. Similarly, polypeptides and
 15 antibodies directed to these polypeptides are useful in providing immunological probes
 for differential identification of the tissue(s) or cell type(s). For a number of disorders
 of the above tissues or cells, particularly of the hematopoietic system and immune
 system, expression of this gene at significantly higher or lower levels may be routinely
 detected in certain tissues and cell types (e.g., brain and other tissue of the nervous
 20 system, bone cells, synovial tissue, endometrial tissue and other reproductive tissue,
 cartilage cells, smooth muscle, and blood cells and cells and tissue of the immune
 system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma,
 urine, synovial fluid or spinal fluid) or another tissue or cell sample or another tissue or
 cell sample or another tissue or cell sample taken from an individual having such a
 25 disorder, relative to the standard gene expression level, i.e., the expression level in
 healthy tissue or bodily fluid or bodily fluid or bodily fluid from an individual not
 having the disorder. Preferred epitopes include those comprising a sequence shown in
 SEQ ID NO. 111 as residues: Met-1 to Cys-9.

The tissue distribution and homology to the LIM-homeodomain containing
 30 proteins, such as T-cell translocation factor, indicates that polynucleotides and
 polypeptides corresponding to this gene are useful for diagnosis and intervention of
 leukemia and other developmental defects. Because of the importance of the LIM-
 homeodomain proteins in development and their correlation to number of leukemic
 diseases, the molecule can be either used as a diagnostic or prognostic indicator for
 35 leukemia progression or a therapeutic target. In addition, polynucleotides and
 polypeptides corresponding to this gene are useful for the detection/treatment of
 neurodegenerative disease states and behavioral disorders such as Alzheimer's Disease,

Parkinson's Disease, Huntington's Disease, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, and autism. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system. Furthermore, homology to the breast auto-antigen may suggest this gene is useful in the detection, prevention, and or treatment of breast cancer and/or other proliferative disorders.

FEATURES OF PROTEIN ENCODED BY GENE NO: 2

Translation product of gene has homology to a highly conserved member of the human calpain family of proteases, Calpain large subunit 1 gene (See Accession No.T32454). Calpains are thought to play a defining role in protein regulation, particularly during development. One embodiment for this gene is the polypeptide fragments comprising the following amino acid sequence:

MKYMGGCAKVMCKYYVILYQGLEYP LLXSGDPETSPPWILRADCIVLSSRNFH
SNXGRLTINKIYVIGGGKYRGEVTNGAK (SEQ ID NO:216);
MGQSELYSSILRNLGVFLVYTRGGFLLSPLLHGTLTCAHS (SEQ ID NO:217);
MVL LLLTVASYTVFWMIGDVLDILFLWNFEYTTLY (SEQ ID NO:218);
MELYNLCPICYFSTVLTTTYIYFVYSQSSXIRMKVP (SEQ ID NO:219);
MQIVIVLYCVRNKDKKKVCTCSVQTQFFPIFPILGCLNGCRTQE (SEQ ID
NO:220); MKYMGGCAKVMCKYYVILYQGLEYP LLX (SEQ ID NO:221);
LEYP LLXSGDPET SPPWILRADCIVLSSRNFH SNX (SEQ ID NO:222); and/or
RNFH SNXGRLTINKIY VIGGGKYRGEVTNGAK (SEQ ID NO:223). An
additional embodiment is the polynucleotide fragments encoding these polypeptide
fragments.

This gene is expressed primarily in caudate nucleus, dermatofibrosarcoma protuberance and apoptotic T-cells, and to a lesser extent in eosinophils, brain and smooth muscle.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurodegenerative diseases or immune disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the nervous system or immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., skin, T-cells and other blood

cells and cells and tissue of the immune system, brain and other tissue of the nervous system, and smooth muscle, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in caudate nucleus and apoptic T-cells indicates that polynucleotides and polypeptides corresponding to this gene are useful for detection or intervention of neurodegenerative diseases and behavioral disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's disease, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder or immune disorders, because the elevated level of the molecule in cells undergoing cell death may be the cause or consequence of these degenerative conditions. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, or disorders of the cardiovascular system.

FEATURES OF PROTEIN ENCODED BY GENE NO: 3

This gene maps to chromosome 15, and therefore, may be used as a marker in linkage analysis for chromosome 15. One embodiment for this gene is the polypeptide fragments comprising the following amino acid sequence: VTNEMSQGRGKYDFY IGLGLAMSSSIFIGGSFILKKKGLLRLARKGSMRAGQGGHAYLKEWLWWAGL LSMGAGEVANFAAYAFAPATLVTPLGALSVLVSAILSSYFLNERLNLHGKIGCL LSILG STVMVIHAPKEEEEIETLNE (SEQ ID NO:224);

VTNEMSQGRGKYDFYIGLGLAMSSSIFIGGSFILKKKGLLRLARKGSMRAGQG GHAYLKEWLWWAGLLSMGAGEVANF (SEQ ID NO:225); NFAAYAFAPATLVTPLGALSVLVSAILSSY (SEQ ID NO:226); and/or ERLNLHGKIGCLLSILGSTVMVIHAPKEEEEIETLNE (SEQ ID NO:227). An additional embodiment is the polynucleotide fragments encoding these polypeptide fragments

This gene is expressed primarily in colon carcinoma cell line, and to a lesser extent in aorta endothelial cells, T-cells, human erythroleukemia cells (HEL), and stromal cells (TF274).

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, colon carcinoma. Similarly, polypeptides and antibodies directed to

these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of colon carcinoma tissues, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., colon, aorta and other vascular tissue, T-cells and other cells and tissue of the immune system, and stromal cells, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 113 as residues: Asn-191 to Ser-196, Asn-208 to Gly-214.

The tissue distribution in colon carcinoma indicates that polynucleotides and polypeptides corresponding to this gene are useful for detection and intervention of colon carcinoma and/or other tumors. Additionally the significant presence in T-cell populations may indicate the involvement of the function of the gene product in cancer immunosurveillance. Furthermore, the tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of cancer and other proliferative disorders, in general. The expression in hematopoietic cells and tissues indicates that this protein may play a role in the proliferation, differentiation, and/or survival of hematopoietic cell lineages. Thus, this gene may be useful in the treatment of lymphoproliferative disorders, and in the maintenance and differentiation of various hematopoietic lineages from early hematopoietic stem and committed progenitor cells.

FEATURES OF PROTEIN ENCODED BY GENE NO: 4

This gene is expressed primarily in ovary.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, reproductive or endocrine disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the reproductive or endocrine systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., ovary and other reproductive tissue, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial

fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 114 as residues:

5 Pro-20 to Ser-25.

The tissue distribution in ovary indicates that polynucleotides and polypeptides corresponding to this gene are useful for assessing reproductive dysfunction or endocrine disorders, because factors secreted by ovary may be involved in reproductive processes, and in cases have global hormonal effects.

10

FEATURES OF PROTEIN ENCODED BY GENE NO: 5

This gene is expressed primarily in tissues in the central nervous system, including pineal gland, frontal cortex, and dura mater, and to a lesser extent in bladder, lung, T-cells and liver.

15

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurodegenerative diseases, endocrine disorders, and immune disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the nervous and endocrine systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., tissue of the nervous system, bladder, lung, liver, and T-cells and other cells and tissues of the immune system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID

20

25

30 NO. 115 as residues: Glu-14 to Arg-20.

The primary tissue distribution in the central nerve system indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection and intervention of neurodegenerative diseases or endocrinedisorders, because extracellular proteins in these tissues may function as a neurotrophic factor, a matrix protein for tissue integrity, a neuroguidance factor or as a hormone.

35

FEATURES OF PROTEIN ENCODED BY GENE NO: 6

This gene is expressed primarily in spleen, resting T-cells, colorectal tumor and pancreatic carcinoma, and to a lesser extent in number of tissues including prostate, synovial hypoxia, osteosarcoma, ulcerative colitis, myeloid progenitor cells, lung and placenta.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, inflammation, immunosurveillance of cancers, and immune and gastrointestinal disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly in carcinogenesis or the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., prostate, synovial tissue, bone cells, colon, myeloid progenitor cells, lung, cells and tissue of the immune system, cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 116 as residues: Arg-29 to Pro-37, Gln-46 to Val-56.

The primary tissue distribution in lymphatic tissues such as T-cells and spleen, as well as tumors and ulcerative tissues indicates that the protein product of this gene may be involved in the immuno response to or immunosurveillance of carcinogenesis and/or inflammatory conditions.

FEATURES OF PROTEIN ENCODED BY GENE NO: 7

The translation product of this gene shares very weak sequence homology with voltage dependent sodium channel protein and Bowman-Birk proteinase inhibitor which is thought to be important in membrane signaling or extracellular signaling cascades. One embodiment for this gene is the polypeptide fragments comprising the following amino acid sequence: RFKTLMTNKSEQDGDSSKTIEISDMKYHIFQ (SEQ ID NO:228); and/or LVEGKLFYAHKVLLVTXSNR (SEQ ID NO:229) (See Accession No. gnllPID1020763 (AB000216)). An additional embodiment is the polynucleotide fragments encoding these polypeptide fragments.

This gene is expressed primarily in prostate cancer.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, prostate cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of prostate cancer tissue, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., prostate and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 117 as residues: Glu-30 to Ser-35.

The tissue distribution in the prostate cancer and homology to sodium channel or proteinase inhibitor suggest that polynucleotides and polypeptides corresponding to this gene are useful for the intervention of cancer progression, because the gene product may be involved in multidrug resistance by altering the drug kinetics by serving the function as a channel transporter. Alternatively, the proteinase inhibitor like function may facilitate tumor metastasis. By targeting these functions, either through vaccine or small molecules, therapeutics may be rationally designed to slow the cancer progression.

FEATURES OF PROTEIN ENCODED BY GENE NO: 8

This gene is expressed primarily in ovary and to a lesser extent in the adrenal gland.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, female infertility and endocrine disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the female reproductive system and the endocrine system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., ovary and other reproductive tissue, and adrenal gland, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample

taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- 5 The tissue distribution of this gene in ovary and adrenal gland indicates that polynucleotides and polypeptides corresponding to this gene are useful for treatment/diagnosis of female infertility, endocrine disorders, ovarian function, amenorrhea, ovarian cancer and metabolic disorders.

FEATURES OF PROTEIN ENCODED BY GENE NO: 9

- 10 This gene is expressed only in prostate cancer.

- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, prostate disorders including cancer. Similarly, polypeptides and
15 antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the endocrine and male reproductive system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., prostate and cancerous and wounded
20 tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- The tissue distribution of this gene only in prostate cancerous tissue, indicates
25 that polynucleotides and polypeptides corresponding to this gene are useful for the treatment/diagnosis of male infertility, metabolic disorders, and prostate disorders including benign prostate hyperplasia and prostate cancer.

FEATURES OF PROTEIN ENCODED BY GENE NO: 10

- 30 This gene is expressed primarily in placenta and to a lesser extent in ovary.

- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, female infertility, pregnancy disorders, and ovarian cancer. Similarly,
35 polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the reproductive

system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., placenta, and ovary and other reproductive tissue, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 120 as residues: Gln-39 to Gly-73.

The tissue distribution of this gene in placenta and ovary indicates that polynucleotides and polypeptides corresponding to this gene are useful for treatment/diagnosis of female infertility, endocrine disorders, fetal deficiencies, ovarian failure, amenorrhea, and ovarian cancer.

FEATURES OF PROTEIN ENCODED BY GENE NO: 11

Gene shares homology with the gene for the Human 3' apolipoprotein B SAR element gene Rh32 (See Accession No. T31530).

This gene is expressed primarily in prostate and in the pancreas.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, prostate and pancreatic disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the endocrine system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., prostate and pancreas, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution of this gene in prostate and pancreas, indicates that polynucleotides and polypeptides corresponding to this gene are useful for treatment/diagnosis of male infertility, prostate disorders including benign prostate hyperplasia, prostate cancer, pancreatic cancer, type I and type II diabetes and hypoglycemia. Homology to a known human apolipoprotein may suggest this gene is useful for the detection, prevention, or treatment of various metabolic disorders,

particularly those secondary to lipoprotein disorders such as atherosclerosis, coronary heart disease, stroke, and hyperlipidemias.

FEATURES OF PROTEIN ENCODED BY GENE NO: 12

- 5 Gene has homology to conserved Beta-casein, an abundant milk protein (See Accession No.Q37894).

 This gene is expressed primarily in stomach.

- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the digestive tract and/or mammary glands. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the digestive system and breast, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., mammary tissue, and stomach and other gastrointestinal tissue, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.
- 10
15
20

- The tissue distribution of this gene indicates a role in the treatment/diagnosis of digestive disorders including stomach cancer and ulceration. Furthermore, the homology to conserved beta-casein may indicate this gene as having utility in the diagnosis and prevention of mammary gland disorders.
- 25

FEATURES OF PROTEIN ENCODED BY GENE NO: 13

 This gene is expressed in brain and lung.

- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurodegenerative disease states, behavioral abnormalities and pulmonary disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune, nervous, and pulmonary systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell
- 30
35

types (e.g., brain and other tissue of the nervous system, and lung, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioral disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder and panic disorder. In addition it could be used in the detection and treatment of pulmonary disease states such as lung lymphoma or sarcoma formation, pulmonary edema and embolism, bronchitis and cystic fibrosis.

FEATURES OF PROTEIN ENCODED BY GENE NO: 14

This gene is expressed exclusively in T-cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., T-cells and other cells and tissue of the immune system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for treatment/detection of immune disorders such as arthritis, asthma, immune deficiency diseases such as AIDS, and leukemia. Additionally, the expression in hematopoietic cells and tissues indicates that this protein may play a role in the proliferation, differentiation, and/or survival of hematopoietic cell lineages. Thus, this gene may be useful in the treatment of lymphoproliferative disorders, and in the maintenance and differentiation of various hematopoietic lineages from early hematopoietic stem and committed progenitor cells.

FEATURES OF PROTEIN ENCODED BY GENE NO: 15

This gene is expressed primarily in T-cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., T-cells and other cells and tissue of the immune system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 125 as residues: Ala-46 to Asp-51.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of immune disorders including: leukemias, lymphomas, auto-immunities, immunodeficiencies (e.g. AIDS), immuno-suppressive conditions (transplantation) and hematopoietic disorders.

FEATURES OF PROTEIN ENCODED BY GENE NO: 16

This gene is expressed primarily in endometrial tumors.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, cancer, particularly endometrial. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the female reproductive system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., endometrial cells and other reproductive cells or tissue, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having

such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of ovarian and other endometrial cancers, as well as reproductive dysfunction, prenatal disorders or fetal deficiencies.

FEATURES OF PROTEIN ENCODED BY GENE NO: 17

This gene is expressed primarily in a variety of osteoclastic cells: osteoclastoma stromal cells, osteosarcoma, chondrosarcoma and stromal cell culture. To a lesser extent, it is also seen in a variety of fetal and embryonic cell and tissue types.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, bone cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the skeletal and developmental systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., bone cells, cartilage, and stomal cells, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 127 as residues: Gln-34 to Gln-41, Asn-76 to Lys-82, Ser-85 to Lys-91.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for treatment and detection of a variety disorders and conditions affecting bone and the skeletal system, including: osteoperosis, fracture, osteosarcoma, osteoclastoma, chondrosarcoma, ossification and osteonecrosis, arthritis, tendonitis, chrondomalacia and inflammation.

FEATURES OF PROTEIN ENCODED BY GENE NO: 18

This gene is expressed primarily in smooth muscle.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are

not limited to, cardiovascular disorders including lymphatic system disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the cardiovascular and lymphatic systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., smooth muscles, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of conditions and pathologies of the cardiovascular system: heart disease, restenosis, atherosclerosis, stroke, angina, thrombosis, and wound healing.

FEATURES OF PROTEIN ENCODED BY GENE NO: 19

The translation product of this gene shares sequence homology with 5'-nucleotidase (See Accession No. 2668557) as well as the gene for alpha-1 collagen type X (See Accession No. gblX67348IMMCOL10A). One embodiment for this gene is the polypeptide fragments comprising the following amino acid sequence:

MAQHFSLAACDVVGFDLDHTLCRYNLPESAPLIYNSFAQFLVKEKGDKELLN
VTPEDWDFCCKGLALDLEDGNFLKLANNGTVLRASHGTKMMTPEVLAEAYG
KKEWKHFLSDTGMACRSKGYYFYDNYFDLPGALLCARVVDYLTCLNNGQKT
FDFWKDIVAAIQHNYKMSAFKENCIGIYFPEIKRDPGRYLHSCPESVKKWLRQL
KNAGKILLITSSHSDYCRLLCEYILGNDFTDLFDIVITNALKPGFFSHLPSQRPF
RTLENDEEQEALPSLDKPGWYSQGNVHLYELLKKMTGKPEPKVVYFGDSMH
SDIFPARHYSNWETVLILEELRGDEGTRSQRPEESEPLEKKKGKYEGPKAKPLNT
SSKKWGSFFIDSVLGLENTEDSLVYTWSCKRISTYSTIAIPSEIAIAELPLDYKFT
RFSSSNSKTAGYYPNPPLVLSSDETLISK (SEQ ID NO:233); and/or
TSSHSDYCRLLCEYILGNDFTDLFDIV (SEQ ID NO:234). An additional
embodiment is the polynucleotide fragments encoding these polypeptide fragments.
Additionally, another embodiment for this gene is the polynucleotide fragments
comprising the following sequence:

CCTTAAAAGCTGACATTTTATAATTGTGTTGTATAGCAGCAACTATATCCTTC
CAAAAATCAAATGTTTTTTGACCATTGTTTCAGTT (SEQ ID NO:230);
CCTTAAAAGCT GACATTTTATAATTGTGTTGTATAGCA (SEQ ID NO:231);

and/or CTTCCTCAAAAA TCAAATGTTTTTTTGACCATTGTTTCAGTT (SEQ ID NO:232). An additional embodiment is the polypeptide fragments encoded by these polynucleotide fragments. This gene maps to chromosome 6, and therefore, may be used as a marker in linkage analysis for chromosome 6.

5 This gene is expressed primarily in prostate and smooth muscle.

 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, prostate cancer and cardiovascular disorders. Similarly, polypeptides
10 and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the prostate and cardiovascular system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., prostate, and smooth muscle, and
15 cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

 The tissue distribution indicates that polynucleotides and polypeptides
20 corresponding to this gene are useful for the treatment and diagnosis of prostate cancer and other disorders. In addition the expression in smooth muscle would suggest a role for this gene product in the treatment and diagnosis of cardiovascular disorders such as hypertension, restenosis, atherosclerosis, stroke, angina, thrombosis, and other aspects of heart disease and respiration.

25

FEATURES OF PROTEIN ENCODED BY GENE NO: 20

 This gene is expressed primarily in endometrial tissue and to a lesser extent in synovium.

 Therefore, polynucleotides and polypeptides of the invention are useful as
30 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, endometrial cancer and arthritis. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of
35 the above tissues or cells, particularly of the reproductive and skeletal systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., endometrial tissue and other reproductive tissue,

and synovial tissue, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 130 as residues: Ser-19 to His-24, Pro-36 to Arg-43, Ala-61 to Gly-67, Pro-86 to Ala-95.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of endometrial cancers, as well as reproductive and developmental disorders (fetal deficiencies and other pre-natal conditions). In addition the expression of this gene product in synovium would suggest a role in the detection and treatment of disorders and conditions affecting the skeletal system, in particular the connective tissues (e.g. arthritis, trauma, tendonitis, chondromalacia and inflammation).

FEATURES OF PROTEIN ENCODED BY GENE NO: 21

This gene maps to chromosome 6, and therefore, may be used as a marker in linkage analysis for chromosome 6.

This gene is expressed primarily in keratinocytes, fetal tissue (especially fetal brain) and leukocytic cell types and tissues (e.g. B-cell, macrophages, Jurkat T-Cell, T cell helper cells, spleen, thymus and lymphoma).

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, integument and immune systems, as well as developmental disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the skin, immune and central nervous systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., keratinocytes, brain and other tissue of the nervous system, differentiating tissue, leukocytes and other cells and tissue of the immune system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of immune disorders including: leukemias, lymphomas, auto-immunities, immunodeficiencies (e.g. AIDS), immuno-suppressive conditions (transplantation) and hematopoietic disorders. Expression in keratinocytes would suggest a role for the gene product in the diagnosis treatment of skin disorders such as cancers (melanomas), eczema, psoriasis, wound healing and grafts. In addition the expression in fetal brain might implicate this gene product in the detection and treatment of developmental and neurodegenerative diseases of the brain and nervous system: behavioral or nervous system disorders, such as depression, schizophrenia, Alzheimer's disease, Parkinson's disease, Huntington's disease, mania, dementia, paranoia, addictive behavior and sleep disorders.

FEATURES OF PROTEIN ENCODED BY GENE NO: 22

Translation product of this gene shares significant homology with the conserved YME1 PROTEIN from *Saccharomyces cerevisiae*, which is a putative ATP-dependent protease thought to regulate the assembly of key respiratory chains within the mitochondria (See Accession No. P32795). Preferred polypeptide fragments comprise the following amino acid sequence:

MKTKNIPEAHQDAFKTGFAEGFLKAQALTQKTNDSLRRLILFVLLLFGIYGL
LKNPFLSVRFRTTGLDSA VDPVQMKNVTFEHVKGVEEAKQELQEVVEFLKNP
QKFITLGGKLPKGILLVGPPGTGKTLLARAVAGEADVPFYYASGSEFDEMFGV
VGASRIRNLFREAKANAPCVIFIDELDSVGGKRIESPMHPYSRQTINQLLAEMD
GFKPNEGVIIGATNFPEALDNALIRPGRFDMQVTVPDPVKGRTEILKWYLNK
IKFDXSVDPEIARGTVGFSGAELENLVNQAALKA AVDGKEMVTMKELGVFQR
QNSNGA (SEQ ID NO:235); MKTKNIPEAHQDAFKTGFAEG (SEQ ID NO:236);
PVQMKNVTFEHVKGVEEAKQELQ (SEQ ID NO:237);
SRQTINQLLAEMDGFKPN EGVII (SEQ ID NO:238); and/or
FSGAELENLVNQAALKA AVDGKEM (SEQ ID NO:239). Also preferred are
polynucleotide fragments encoding these polypeptide fragments.

This gene is expressed primarily in T-cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune and hematopoietic disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and hematopoietic systems,

expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., T-cells and other cells and tissue of the immune system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of immune disorders including: leukemias, lymphomas, auto-immunities, immunodeficiencies (e.g. AIDS), immuno-suppressive conditions (transplantation) and hematopoietic disorders. Furthermore, the homology of this gene indicates that it may play an important role in disorders affecting metabolism.

15 **FEATURES OF PROTEIN ENCODED BY GENE NO: 23**

This gene is expressed primarily in human chronic synovitis.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, synovial and other inflammatory disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the synovial tissue and immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., synovial tissue, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that the protein product of this gene are useful for study, diagnosis and treatment of inflammatory disorders such as chronic synovitis.

FEATURES OF PROTEIN ENCODED BY GENE NO: 24

This gene is expressed primarily in pituitary, breast cancer, and bone marrow; and to a lesser extent in breast, prostate, uterine cancer and cerebellum.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a

- biological sample and for diagnosis of diseases and conditions which include, but are not limited to, endocrine, reproductive disorders and cancers. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the reproductive, metabolic and endocrine systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., pituitary, mammary tissue, bone marrow, prostate, reproductive tissue, uterus, and brain and other tissue of the nervous system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 134 as residues: Asp-32 to Gln-38, Lys-88 to Ile-97.
- The tissue distribution indicates that the protein products of this gene are useful for the study, treatment and diagnosis of various endocrine disorders, reproductive diseases and disorders and cancers.

FEATURES OF PROTEIN ENCODED BY GENE NO: 25

- The translation product of this gene shares sequence homology with androgen withdrawal apoptosis protein in rat which is thought to be important in programmed cell death. Preferred polypeptides encoded by this gene comprise the following amino acid sequence:
- LPMWQVTAFLDHNIVTAQTTWKGLWMSCVVQSTGHMQCKVYDSVLALSTEV
 QAARALTVSAVLLAFVALFVTLAGAQCCTTCVAPGPAKARVALTGGVLYLFCGL
 LALVPLCWFANIVVREFYDPSVPVSQKYELGAXLYIGWAATALLMVGGCLLC
 GAWVCTGRPDLSFPVKYSAPRRPTATGDYDKKNYV (SEQ ID NO:240). This polypeptide is expected to contain multiple transmembrane domains. The extracellular portion of the polypeptide is expected to comprise residues 1-51 of the foregoing amino acid sequence. Therefore, particularly preferred polypeptides encoded by this gene comprise residues 1-51 of the foregoing amino acid sequence. Polynucleotides encoding the foregoing polypeptides are also provided.
- This gene is expressed primarily in human adult pulmonary and brain (striatum) tissue and to a lesser extent in thymus, synovium and testis.
- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are

not limited to, reproductive, metabolic, and neurodegenerative disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the reproductive, nervous, respiratory and metabolic systems expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., thymus, synovial tissue, testis and other reproductive tissue, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution and homology to androgen withdrawal apoptosis rat gene protein indicates that polynucleotides and polypeptides corresponding to this gene are useful for study, diagnosis and treatment of disorders in which the mechanism controlling programmed cell death is instrumental. This could include reproductive, neurodegenerative, and various metabolic disorders and diseases such as cancer.

FEATURES OF PROTEIN ENCODED BY GENE NO: 26

The translation product of this gene shares homology with both ubiquitin and a G-protein coupled receptor TM3 consensus polypeptide (see Genbank accession Nos. gnllPIDe331456 (AJ000657) and R50664, respectively). Preferred polypeptides encoded by this gene comprising the following amino acid sequence:

LHYFALSFVLILTEICLVSSGMGF (SEQ ID NO:241);
QLRNGIPPGRKALFCSGKPR LFTLGQGRTCA (SEQ ID NO:242); and/or
WSGLWVTTWNGSSGERTPSPWRRK RASQSAGRIASWMSF (SEQ ID NO:243).

An additional embodiment is polynucleotides encoding these polypeptides. This gene maps to chromosome 1, and therefore, may be used as a marker in linkage analysis for chromosome 1.

This gene is expressed primarily in activated T cells and to a lesser extent in CD34 depleted buffy coat.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune and hemopoietic disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the hemopoietic and immune system,

- expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., T-cells and other blood cells and other cells and tissue of the immune system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell
- 5 sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 136 as residues: Thr-15 to His-21, Gly-30 to Lys-39, Arg-113 to Met-118, Arg-178 to Ala-187.
- 10 The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex vivo culture, bone marrow
- 15 transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia. The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency etc. Furthermore, the homology to G-coupled proteins as well as to ubiquitin may implicate this gene as being important in regulation of gene expression and protein
- 20 sorting - both of which are vital to development and wound healing models. Therefore, the gene may provide utility in the diagnosis, prevention, and/or treatment of various developmental disorders.

FEATURES OF PROTEIN ENCODED BY GENE NO: 27

- 25 This gene is expressed primarily in activated T cells and to a lesser extent in fetal kidney.
- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are
- 30 not limited to, immune, developmental and metabolic diseases. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and metabolic systems, expression of this gene at significantly higher or lower levels may be routinely
- 35 detected in certain tissues and cell types (e.g., T-cells and other cells and tissue of the immune system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from

an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides
 5 corresponding to this gene are useful for the study and treatment of diseases and disorders of the immune, metabolic, and endocrine systems; such as renal diseases and T cell dysfunctions. Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved in immune functions. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immune deficiency
 10 diseases such as AIDS, and leukemia.

FEATURES OF PROTEIN ENCODED BY GENE NO: 28

The translation product of this gene shares sequence homology with Cystatin-related epididymal specific protein in mouse which is thought to be important in
 15 reproductive system function/regulation (See Genbank accession no.bbsl18813). Based on the structural similarity between these proteins, the translation product of this clone, hereinafter "Cystatin G", is expected to share biological activities with cystatin related proteins and other cysteine protease inhibitors. Such activities are known in the art and are described elsewhere herein. Preferred polypeptides encoded by this gene
 20 comprising the following amino acid sequence:
 MPRCRWLSLILLTIPLALVARKDPKKNETGVLRLKLPVNASNANVKQCLWFA
 MQEYNKESEDKYVFLVVKTLQAQLQVTNLLEYLIDVEIARSDCRKPLSTNEICAI
 QENSKLKRKLSCSFLVGALPWNGEFTVMEKKCEDA (SEQ ID NO:246);
 ARKDPKKNETGVLRLKLPVNASNANVKQCLWFAMQEYNKESEDKYVFLVVK
 25 TLQAQLQVTNLLEYLIDVEIARSDCRKPLSTNEICAIQENSKLKRKLSCSFLVGA
 LPWNGEFTVMEKKCEDA (SEQ ID NO:248);
 CLWFAMQEYNKESEDKYVFLVVKTLQAQLQVTNLLEYLIDVEIARSDCRKPLST
 NEICAIQENSKLKRKLSCSFLVGALPWNGEFTVMEKKC (SEQ ID NO:247);
 EYNKESEDKYVFLV (SEQ ID NO:244); and/or IDVEIARSDCRKPL (SEQ ID
 30 NO:245). An additional embodiment is the polynucleotide fragments encoding these polypeptide fragments. Preferred cystatin polypeptide fragments are shown to be active in the following assays: The methods used for active site titration of papain, titration of the molar enzyme inhibitory concentration in cystatin G preparations, and for determination of equilibrium constants for dissociation (K_i) of complexes between
 35 cystatin G and cysteine peptidases are described in detail in Hall et al., Biochem. J., 291:123-29 (1993) and Abrahamson, Methods Enzymol., 244:685-700 (1994), both of which are hereby incorporated herein by reference. The enzymes used for equilibrium

assays are papain (EC 3.4.22.2; from Sigma, St Louis, MO) and cathepsin B (EC 3.4.22.1; from Calbiochem, La Jolla, CA). The fluorogenic substrate used was Z-Phe-Arg-NHMec (10 mM; from Bachem Feinchemikalien, Bubendorf, Switzerland) and the assay buffer was 100 mM Na-phosphate buffer (pH 6.5 and 6.0 for papain and cathepsin B, respectively), containing 1 mM dithiothreitol and 2 mM EDTA. Steady state velocities are measured and K_i values were calculated according to Henderson, Biochem J., 127:321-333 (1972), incorporated herein by reference. Corrections for substrate competition are made using K_m values of 150 μ M for cathepsins B (Barrett and Kirschke, Methods Enzymol., 80:535-561 (1981) and 60 μ M for papain (Hall et al., Biochem. J., 291:123-29 (1992)), both of which are hereby incorporated herein by reference.

This gene is expressed primarily in human testes.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, reproductive disorders and cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the reproductive system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., testis and other reproductive tissue, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 138 as residues: Arg-21 to Thr-29.

The tissue distribution and homology to cystatin-related epididymal specific protein-mouse indicates that polynucleotides and polypeptides corresponding to this gene are useful for study, diagnosis and treatment of reproductive diseases and disorders. Cysteine proteinase inhibitors of the cystatin superfamily are ubiquitous in the body and are generally tight-binding inhibitors of papain-like cysteine proteinases, such as cathepsins B, H, L, S, and K (for review, see Ref. 1). They should therefore serve a protective function to regulate the activities of such endogenous proteinases, which otherwise may cause uncontrolled proteolysis and tissue damage. Cysteine proteinase activity can normally not be measured in body fluids, but can be detected extracellularly in conditions like endotoxin-induced sepsis (2), metastasizing cancer (3), and at local inflammatory processes in rheumatoid arthritis (4), purulent bronchiectasis

(5) and periodontitis (6), which indicates that a tight cystatin regulation is a necessity in the normal state. A deficiency state in which the levels of the intracellular cystatin, cystatin B, are lowered due to mutations has recently been shown to segregate with a form of progressive myoclonus epilepsy (7), which points to additional specialized functions of cystatins. Moreover, results showing that chicken cystatin inhibits polio virus replication (8), human cystatin C inhibits corona- and herpes simplex virus replication (9,10), and human cystatin A inhibits rhabdovirus-induced apoptosis (11) in cell cultures indicates that cystatins play additional roles in the human defense system. The cystatins constitute a superfamily of evolutionary related proteins, all composed of at least one 100-120 residue domain with conserved sequence motifs (12). The previously well characterized single-domain human members of superfamily could be grouped in two protein families. The Family 1 members, cystatins (or stefins) A and B, contain approximately 100 amino acid residues, lack disulfide bridges, and are not synthesized as preproteins with signal peptides. The Family 2 cystatins (cystatins C, D, S, SN, and SA) are secreted proteins of approx. 120 amino acid residues (Mr 13,000-14,000) and have two characteristic intrachain disulfide bonds. Recently, we identified an additional human cystatin superfamily member by EST1 sequencing in epithelial cell derived cDNA libraries which we named cystatin E (13). The same cystatin was independently discovered by differential display experiments as a mRNA species down-regulated in breast tumor tissue, but present in the surrounding epithelium and reported under the name cystatin M (14). Cystatin E/M is an atypical, secreted low-Mr cystatin in that it is a glycoprotein and just shows 30-35% sequence identity in alignments with the human Family 2 cystatins, which shows that additional cystatin families are yet to be identified (13). The cystatin E/M gene has been localized to chromosome 2 (15), whereas all human Family 2 cystatin genes are clustered on the short arm of chromosome 20 (16), which further stresses that cystatin E/M is just distantly related to the other secreted human low-Mr cystatins.

FEATURES OF PROTEIN ENCODED BY GENE NO: 29

The translation product of this gene shares sequence homology with the leukocyte-associated Ig-like receptor-1, putative inhibitory receptor which is thought to be important in regulation of various physiological functions (See Accession No. gil2352941 (AF013249). Preferred polypeptides encoded by this gene comprise the following amino acid sequence:

DSPDTEPGSSAGPTQRPSDNSHNEHAPASQGLKAEHL YILIGVS (SEQ ID NO:249); HRQNQIKQGPPRSKDEEQKPPQRPDLAVDVLERTADKATVNGL PEKDRETDTSALAAGSSQEVTYAQLDHWALTQRTARAVSPQSTKPMASITYAA

VARH (SEQ ID NO:250);

MSPHPTALLGLVLCLAQTIHTQEEDLPRPSISAEPGTVIPLGSHVTFVCRGPVGV
QTFRLERESRSTYNDTEDVSQASPSESEARFRIDSVSEGNAGPYRCIYYKPPKW

SEQSDY (SEQ ID NO:251); TALLGLVLCLAQTIHTQE (SEQ ID NO:252);

- 5 LPRPSISAEPGTVI (SEQ ID NO:253); CRGPVGVQTFRLERE (SEQ ID NO:254);
and/or VLERTADKATVNGLEKDRDTSALAAGSS (SEQ ID NO:255).

Additional embodiments of the invention include polynucleotides encoding these polypeptides.

- 10 This gene is expressed primarily in macrophages and T-cells and to a lesser extent in human fetal heart.

- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, developmental, inflammatory, and immune disorders. Similarly,
- 15 polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the growth and inflammatory systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., macrophages, T-cells
- 20 and other cells and tissue of the immune system, heart, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those
- 25 comprising a sequence shown in SEQ ID NO. 139 as residues: His-20 to Arg-28, Glu-61 to Val-74, Ser-78 to Ala-84, Lys-105 to Ser-117.

- The tissue distribution and homology to putative inhibitory receptor indicates that polynucleotides and polypeptides corresponding to this gene are useful for the study, diagnosis and treatment of functional disorders of the developing fetal heart;
- 30 including circulatory and vascular; and inflammatory disorders. In addition expression in macrophages and lymphocytes indicates a role in the treatment/detection of immune disorders including disorders such as arthritis, asthma, immune deficiency diseases such as AIDS, and leukemia.

35 **FEATURES OF PROTEIN ENCODED BY GENE NO: 30**

The translation product of this gene shares sequence homology with erythroid cell specific transcription factor- murine which is thought to be important in normal

physiological function of erythroid cells. In addition, the translation product of this gene also shares homology with the conserved 3-phosphoglycerate dehydrogenase gene which is essential component of metabolic biosynthetic pathways. Preferred polypeptides comprise the following amino acid sequence:

- 5 MNTPNGNSLSAAELTCGMIMCLARQIPQATASMKDGKWERKKFMGTELNGK
TLGILGLGRIGREVATRMQSFGMKTIGYDPIISPEVSASFGVQQLPLEEIWPLCDF
ITVHTPLLPSTTGLLNDNTFAQCKKGVRV VNCARGGIVDEGALLRALQSGQCA
GAALDVFTTEPPRDRAVDHENVISCPHLGASTKEAQSRCGEELAVQFVDMVK
GKSLTGVPVNAQALTSAFSPHTKPWIGLAEALGTLMRAWAGSPKGTIQVITQGT
10 SLKNAGNCLSPAVIVGLLKEASKQADVNLVNAKLLVKEAGLNVTTSHSPAAPG
EQGFGECLLAVALAGAPYQAVGLVQGTTPVLQGLNGAVFRPEVPLRRDLPLLL
FRTQTSDPAMLPTMIGLLAEAGVRLLSYQTSLSVSDGETWHVMGISSLLPSLEAW
KQHVTEAFQFHF (SEQ ID NO:256); MAFANLRKVLISDSLDPCCRKILQ (SEQ ID
NO:257); GGLQVVEKQNL SKEELIA (SEQ ID NO:258);
15 MCLARQIPQATASMKDGKWERKKFMGTEL (SEQ ID NO:259);
ALTSAFSPHTKPWIGLAEALGTLMRAWAG (SEQ ID NO:260); and/or
EVPLRRDLPLLLFRTQTSDPAMLPTMIGLLAEAGVR (SEQ ID NO:261). Also
preferred are polynucleotide fragments encoding these polypeptides. This gene maps to
chromosome 1, and therefore, may be used as a marker in linkage analysis for
20 chromosome 1.

This gene is expressed primarily in IL-1 induced smooth muscle and fetal kidney and to a lesser extent in myeloid progenitor cell line and bone marrow.

- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a
25 biological sample and for diagnosis of diseases and conditions which include, but are
not limited to, immune, hemopoietic, and cardiovascular disorders. Similarly,
polypeptides and antibodies directed to these polypeptides are useful in providing
immunological probes for differential identification of the tissue(s) or cell type(s). For a
number of disorders of the above tissues or cells, particularly of the hemopoietic and
30 immune system, expression of this gene at significantly higher or lower levels may be
routinely detected in certain tissues and cell types (e.g., smooth muscle, kidney,
myeloid progenitor cells, bone, and cancerous and wounded tissues) or bodily fluids
(e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell
sample taken from an individual having such a disorder, relative to the standard gene
35 expression level, i.e., the expression level in healthy tissue or bodily fluid from an
individual not having the disorder. Preferred epitopes include those comprising a
sequence shown in SEQ ID NO. 140 as residues: Met-1 to Asn-7, Met-33 to Lys-42,

Asn-123 to Cys-130, Glu-169 to Asp-174, Ser-192 to Gly-201, Thr-266 to Asn-273, Pro-318 to Phe-323.

The tissue distribution and homology to erythroid cell specific murine transcription factor indicates that polynucleotides and polypeptides corresponding to this gene are useful for study, diagnosis and treatment of disorders and diseases involving the hemopoietic and immune systems; the maturation of progenitor cells; and the development of various smooth muscle tissues (heart, etc.). In addition, homology to a key biosynthetic protein implicates this the protein product of this gene as being important in metabolism. Therefore, the protein may show utility in the diagnosis, prevention, and/or treatment of metabolic disorders and conditions.

FEATURES OF PROTEIN ENCODED BY GENE NO: 31

This gene is expressed primarily in human adult testes.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, reproductive disorders, particularly of the male genitalia. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the reproductive system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 141 as residues: Met-1 to Pro-8, Ser-45 to Thr-50.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the study, diagnosis, treatment, and possibly prevention of various male reproductive disorders and diseases including male impotence, failed libido and male secondary sex characteristics, infertility, and testicular cancer.

35 FEATURES OF PROTEIN ENCODED BY GENE NO: 32

This gene is expressed primarily in human adult testis.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, reproductive disorders and cancers of the male reproductive system.

5 Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the reproductive system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., testis and other reproductive
10 tissue, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

15 The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the study, diagnosis, treatment, and possibly prevention of various male reproductive disorders and diseases including male impotence, failed libido and male secondary sex characteristics, infertility, and testicular cancer.

20

FEATURES OF PROTEIN ENCODED BY GENE NO: 33

The translation product of this gene shares homology to the W09D10.1 protein of *Caenorhabditis elegans*. In addition, the gene also shares homology with the human protein hRIP, a protein known to be critical for HIV replication (See Accession
25 Nos. gnlIPIDe1186472 and W12713). Preferred polypeptides encoded by this gene comprise the following amino acid sequence:

MDLLGLDAPVACSIANSKTSNTLEKDLLASVPSPSSSGSRKVVGSMPTAGSA
GSVPENLNLFPPEGSKSEEIGKKQLSKDSILSLYGSQTXQMPTQAMFMAPAQM
AYPTAYPSFPGVTPPNSIMGSMMPPPVGMVAQPGASGMVAPMAMPAGYMGG
30 MQASMMGVPNGMMTTQQAGYMAAGMAAMPQTVYGVQPAQLQWNLTQMTQ
QMAGMNFYGANMMNYGQSMGGNGQAANQTLSPQMWKFGTRFLANLLE
EDNKFCADCQSKGPRWASWNIGVFICIRCAXIHRNLGVHISRVKSVNLDQWTQ
VQIQC (SEQ ID NO:267); MQXMGNGKANRLYEAYLPETFRRPQIDPAVEGFIR
DXYE (SEQ ID NO:268); EEDNKFCADCQSKGPRWASWN (SEQ ID NO:263);
35 GVFCIRCAXIHR NLGVHIS (SEQ ID NO:264); and/or SVNLDQWTQVQIQCMQX
MGNGKA (SEQ ID NO:265). Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in lymphoid tumors.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune and inflammatory disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune, hematopoietic and inflammatory, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., lymphoid tissue and other tissue and cells of the immune system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 143 as residues: Cys-21 to Trp-28.

The tissue distribution indicates that the protein products of this gene are useful for study, diagnosis and treatment of various immune disorders and diseases, including self-recognition and rejection functions of the immune system, hematopoietic disorders, and inflammatory disorders. Homology to the W09D10.1 of *C.elegans* and the hRIP implicates this gene as playing a role as an essential receptor for host-viral interactions including, but not limited to retroviral infections such as AIDS.

FEATURES OF PROTEIN ENCODED BY GENE NO: 34

The translation product of this gene shares homology to an *Arabidopsis thaliana* recombination and DNA-damage resistance/repair protein (See Accession No.gil166694). Preferred polypeptides encoded by this gene comprise the following amino acid sequence:

KYGKVGKCVIFEIPGAPDDEAVRIFLEFERVESAIKAVVDLNGRYFGGRVVKAC
FYNLDKFRVLDLA (SEQ ID NO:269); KAVDLGRYFGGR (SEQ ID NO:270);
and/or EAVRIFFRE (SEQ ID NO:271). Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in ovarian and other cancers.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, cancer, particularly of the female reproductive system. Similarly,

polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the reproductive system, expression of this gene at significantly higher or lower levels may be routinely
 5 detected in certain tissues and cell types (e.g., ovaries and other reproductive tissue, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the
 10 disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 144 as residues: Thr-11 to Trp-19, Ala-40 to Gln-47, Lys-58 to Arg-66, Asp-98 to Lys-110, Arg-114 to Glu-121.

The tissue distribution in tumors of ovarian origins combined with the homology to a known DNA damage repair enzyme indicates that polynucleotides and
 15 polypeptides corresponding to this gene are useful for diagnosis and intervention of tumors. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tumors and tissues.

FEATURES OF PROTEIN ENCODED BY GENE NO: 35

20 Translation product of this gene shares homology with human stomatin, intestinal surface antigens, as well as protein F30A10.5 of *Caenorhabditis elegans* (See Accession No.gnllPIDle276130). Preferred polypeptides encoded by this contig comprise the following amino acid sequence: RMGRFHRILEPGLNILIPVLDRIRYVQ
 SLKEIVINVPEQSAVTLDNVTLQIDGVLRLIMDPYKASYGVEDPEYAVTQLAQT
 25 TMRSELGKLSLDKVFREERESLNASIVDAINQAADCWGIRCLRYEIKDIHVPPRV
 KESMQMQVEAERRKRATVLESEGTRESAINVAEGKKQAQILASEAEKAEQINQA
 AGEASAVLAKAKAKAEAIRILAAALTQHNGDAAASLTVAEQYVSAFSKLAKDS
 NTILLPSNPGDVTSMAQAMGVYGALTKAPVPGTPDSLSSGSSRDVQGTDSL
 DEELDRVKMS (SEQ ID NO:272); ASYGVEDPEYAVTQLAQT MRSELGK (SEQ
 30 ID NO:273); MQMQVEAERRKRATVLESEGTRESAIN (SEQ ID NO:274);
 LTVAEQYVSAFSKLAKDSNTILLPSN (SEQ ID NO:275), and/or
 LLGATAPLVSLVPEVAAAVGNAGARGAXHWGPFAEGLSTGFWPR SARASSGL
 PRNTVVLVFPQQEAWVVE (SEQ ID NO:276). Polynucleotides encoding these
 polypeptides are also provided.

35 This gene is expressed primarily in activated T-cells and to a lesser extent in other cell types.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., T-cells and other cells and tissue of the immune system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 145 as residues: Arg-23 to Pro-33, Pro-184 to Ser-189, Ala-196 to Arg-201, Glu-208 to Ser-213, Glu-230 to Ile-237, Gly-326 to Leu-331, Gly-334 to Gln-340.

The tissue distribution indicates that the protein products of this gene are useful for the treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia. The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency etc. In addition, the homology to known intestinal antigens may suggest that the protein is important in the diagnosis, treatment, and/or prevention of gastrointestinal disorders.

FEATURES OF PROTEIN ENCODED BY GENE NO: 36

Translation product of this gene has homology to a human estrogen receptor variant from human breast cancer. Preferred polypeptides encoded by this gene comprise the following amino acid sequence: RMWRNGTHFWECKIVQPLWK TVWWFPRKLSIELPENLAILIGTYFK (SEQ ID NO:277); and/or LKRHFPEANK HVKRCSTSLDIREIQIKIMRY (SEQ ID NO:278). Polynucleotides encoding these polypeptides are also provided.

This gene is expressed primarily in ulcerative colitis.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a

biological sample and for diagnosis of diseases and conditions which include, but are not limited to, intestinal ulcers, inflammatory conditions and cancers, particular of the breast. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the gastrointestinal system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., colon and other gastrointestinal tissue, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution in colon and breast origins indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and intervention of tumors or other conditions within these tissues, in addition to other tumors where expression has been indicated. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tumors and tissues.

20 FEATURES OF PROTEIN ENCODED BY GENE NO: 37

This gene is expressed primarily in epithelial cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, cancers and skin disorders, particularly melanoma. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the skin and other epithelia, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 147 as residues: Met-1 to Tyr-6.

The tissue distribution in epithelial tissue indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and intervention of

tumors of this tissue. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

5 FEATURES OF PROTEIN ENCODED BY GENE NO: 38

This gene is expressed primarily in adult retina.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, diseases of the eye. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the eye, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., epithelial cells, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 148 as residues: Cys-14 to Lys-21.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for treatment and diagnosis of disorders of the eye.

25 FEATURES OF PROTEIN ENCODED BY GENE NO: 39

This gene is expressed primarily in bone marrow and fetal liver.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, hemopoietic disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the hemopoietic system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., bone marrow and liver, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard

gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for treatment and diagnosis of disorders of the hemopoietic system.

FEATURES OF PROTEIN ENCODED BY GENE NO: 40

This gene is expressed primarily in lymph node, fetal liver and brain.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, hemopoietic diseases and disorders of the CNS. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the hemopoietic and CNS, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., lymphoid tissue and other tissue of the immune system, liver, and brain and other tissue of the nervous system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that the protein products of this gene are useful for the diagnosis and treatment of cancer and other proliferative disorders. Expression in embryonic tissue and other cellular sources marked by proliferating cells indicates that this protein may play a role in the regulation or cellular division. Additionally, the expression in hematopoietic cells and tissues indicates that this protein may play a role in the proliferation, differentiation, and/or survival of hematopoietic cell lineages. Thus, this gene may be useful in the treatment of lymphoproliferative disorders, and in the maintenance and differentiation of various hematopoietic lineages from early hematopoietic stem and committed progenitor cells. In addition, polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioral disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, and autism. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental

disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system.

FEATURES OF PROTEIN ENCODED BY GENE NO: 41

- 5 The translation product of this gene shares sequence homology with fibropellin and epidermal growth factors which are thought to be important in growth and regeneration of epidermal cells (See Genbank Accession Nos. W11719 and gi1310660). Preferred polypeptides comprise the following amino acid sequence:
- 10 GTRPGESHANDLECSGKGKCTTKPSEATFSCTCEEQYVGTFCEEYDACQRKPC
 QNNASCIDANEKQDGSNFTCVCLPGYTGELCQSKIDYCLDPCRNGATCISSLS
 GFTCQCPEGYFGSACEEKVDPCASSPCQNNGTCYVDGVHFTCNCSPGFTGPTC
 AQLIDFCALSPCAHGTCRSVGTSYKCLCDPGYHGLYCEEEYNECLSAPCLNAA
 TCRDLVNGYECVCLAEYKGTCELKDPSCANVSCLNGATCDSGLNGTCICA
 PGFTGEECDIDINECDSPCHHGGSCLDQPNGYNCHCPHWVGANCEIHLQW
 15 KSGHMAESLTN (SEQ ID NO:279); GKCTTKPSEATFSCTCEEQYVGTF (SEQ
 ID NO:280); CAHG TCRSVGTSYKCLCDPGYH (SEQ ID NO:281); and/or
 CANVSCLNGATCDSGLNG TCICAPGFTGEECD (SEQ ID NO:282).
 Polynucleotides encoding these polypeptides are also provided.

- 20 This gene is expressed primarily in brain and kidney and to a lesser extent in several other tissues and organs.

- 25 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the neural and renal systems, particularly growth disorders such as cancer. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the neural and renal systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., brain and other
 30 tissue of the nervous system, and kidney, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- 35 The tissue distribution and homology to epidermal growth factor indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of growth disorders especially in the neural and renal systems. In

addition, polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioral disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, and autism.

- 5 In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system

FEATURES OF PROTEIN ENCODED BY GENE NO: 42

- 10 This gene is expressed primarily in brain, kidney and stromal cells.

- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, disorders of the CNS and hemopoietic system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the hemopoietic, renal and central nervous system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., brain and other tissue of the nervous system, kidney, and stromal cells, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 152 as residues: Lys-71 to Trp-76, Glu-99 to Gly-108, Arg-142 to Ser-149.
- 15
20
25

- The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioral disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, and autism. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, or disorders of the cardiovascular system. In addition, polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. The uses include
- 30
35

bone marrow cell ex vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia. The gene product is thought to be involved in lymphopoiesis, therefore, it can be used in immune disorders to modulate infection, inflammation, allergy, immunodeficiency, etc.

5

FEATURES OF PROTEIN ENCODED BY GENE NO: 43

The preferred polypeptide encoded by this gene comprise the following amino acid sequence: MAQNLKDLA₁GRLPAGPRGMGTALKLLLGAGAVAYGVRESVFT
VEGGHRAIFFNRIGGVQQDTILAEGLHFRIPWFQYPIIYDIRARPRKISSPTGSKD
10 LQMVNISLRVLSRPNAQELPSMYQRLGLDYEERVLPSIVNEVLKSVVAKFNASQ
LITQRAQVSLIRRELTERAKDFSLILDDVAITELSF₁SREYTA₁AVEAKQVAQQEAQ
RAQFLVEKAKQEQRQKIVQAEGEAEAAKMLGEALSKNPGYIKLRKIRAAQNIS
KTIATSQNRIYLTADNLVLNLQDESFTRGSDSLIKGKK (SEQ ID NO:283). The
gene product above share sequence similarity with prohibitin. Thus, these polypeptides
15 are expected to share biological activities with prohibitin. Such activities are known in
the art and discussed elsewhere herein.

This gene is expressed primarily in fetal brain.

Therefore, polynucleotides and polypeptides of the invention are useful as
reagents for differential identification of the tissue(s) or cell type(s) present in a
20 biological sample and for diagnosis of diseases and conditions which include, but are
not limited to, neural diseases. Similarly, polypeptides and antibodies directed to these
polypeptides are useful in providing immunological probes for differential identification
of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells,
particularly of the nervous system, expression of this gene at significantly higher or
25 lower levels may be routinely detected in certain tissues and cell types (e.g., brain and
other tissue of the nervous system, and cancerous and wounded tissues) or bodily
fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or
cell sample taken from an individual having such a disorder, relative to the standard
gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an
30 individual not having the disorder. Preferred epitopes include those comprising a
sequence shown in SEQ ID NO. 153 as residues: Ala-85 to Ser-91, Pro-93 to Asp-98,
Glu-167 to Lys-173, Gln-205 to Ala-210.

The tissue distribution and structural similarity to prohibitin indicates that the
protein products of this gene are useful for the detection/treatment of neurodegenerative
35 disease states and behavioral disorders such as Alzheimer's Disease, Parkinson's
Disease, Huntington's Disease, schizophrenia, mania, dementia, paranoia, obsessive
compulsive disorder, panic disorder, and autism. In addition, the gene or gene product

may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, sexually-linked disorders, and/or disorders of the cardiovascular system.

5 FEATURES OF PROTEIN ENCODED BY GENE NO: 44

The translation product of this gene shares sequence homology with the F44G4.1 gene of the *c. elegans* genome which has no known function (See Accession No.gnlIPIDle236516). The translation product of this gene also shares sequence homology with the human torsionA and torsionB gene products, a gene candidate for
10 the Torsion Dystonia disease locus (See Accession Nos gil2358279 (AF007871) and gil2358281 (AF007872)). One embodiment for this gene is the polypeptide fragments comprising the following amino acid sequence: KALALSFHGWSGTGKNFV (SEQ ID NO:284); NLIDYFIPFLPLEYRHRVRLCAR (SEQ ID NO:285); NLIDYFIPFLPLEYRHRVRLC (SEQ ID NO:286); CHQTLFIFDEAEKLHPGLLEVLGPHL (SEQ ID
15 NO:287); and/or PEKALALSFHGWSGTGKNFVA (SEQ ID NO:288). An additional embodiment is the polynucleotide fragments encoding these polypeptide fragments.

This gene is expressed primarily in tonsils.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a
20 biological sample and for diagnosis of diseases and conditions which include, but are not limited to, such as tonsilitis or adnoiditis. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at
25 significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., tonsils, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the
30 disorder.

The tissue distribution and homology to F44G4.1 gene of the *c. elegans* genome indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and detection of conditions affecting the tonsils. The tonsils have not been thoroughly studied and the actual function of this organ is not known,
35 but this gene could be used in determining what may trigger tonsillitis. Especially in children, where the tonsils seem to be most active. Furthermore, due to the homology

of this gene, it may display potential utility in the detection, diagnosis, and/or treatment for Torsion Dystonia disease.

FEATURES OF PROTEIN ENCODED BY GENE NO: 45

- 5 Has exact sequence homology on the nucleotide level as Human HepG2 3' region cDNA, but the function of this gene is not known.

 This gene is expressed primarily in osteoclastoma stromal cells and to a lesser extent in T-cells.

- Therefore, polynucleotides and polypeptides of the invention are useful as
10 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, leukemia and bone disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of
15 the above tissues or cells, particularly of the haemolymphoid system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., bone tissue, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene
20 expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

 The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of diseases such as leukemia.

25

FEATURES OF PROTEIN ENCODED BY GENE NO: 46

 This gene is expressed primarily in activated monocytes.

- Therefore, polynucleotides and polypeptides of the invention are useful as
30 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, immune disorders, including leukemia and allergies. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the lymphoid system,
35 expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., hemopoietic cells, bone marrow, and spleen, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial

fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 156 as residues:

5 Met-1 to Gly-7.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment in tissue repair and modeling since monocytes engage the synthesis and secretion of many cytokines which are soluble proteins that regulate highly diverse aspects of cellular biology. Monocytes are
 10 also important in the fact that their expression of Major Histocompatibility Factor II (MHCII) enable them to select and stimulate the appropriate lymphocytes to combat specific antigens in the blood. Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved in immune functions. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immune
 15 deficiency diseases such as AIDS, and leukemia.

FEATURES OF PROTEIN ENCODED BY GENE NO: 47

Translation product of this gene has homology to the Na⁺/H⁺-exchanging protein: Na⁺/H⁺ antiporter in *Methanobacterium thermoautotrophicum* as well as the
 20 Na⁺/H⁺ antiporter *cdu2'* in *Clostridium difficile* (See Accession Nos. gil2621849 (AE000854) and pirlJC5343/JC5343, respectively). Thus, it is likely that this gene has similar Na⁺/H⁺ antiporter activity. One embodiment for this gene are polypeptide fragments comprising the following amino acid sequence:
 NLKEKIFISFAWLPKATVQAAIG (SEQ ID NO:289) and/or
 25 WLPKATVQAAIGSVALD (SEQ ID NO:290). An additional embodiment is the polynucleotide fragments encoding these polypeptide fragments.

This gene is expressed primarily in osteoclastoma cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a
 30 biological sample and for diagnosis of diseases and conditions which include, but are not limited to, osteoporosis, leukemia. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the lymphoid and skeletal systems, expression of this
 35 gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., bone cells, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell

sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 157 as residues: His-35 to Gln-43.

- 5 The tissue distribution predominantly in osteoclastoma cells (the site of hematopoiesis) indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of bone related diseases including osteoporosis, osteopetrosis and leukemia. Furthermore, its homology to known transporter proteins may suggest the protein is useful in the diagnosis, treatment, and
- 10 prevention of various developmental and metabolic disorders, particularly those based upon ion and proton transport.

FEATURES OF PROTEIN ENCODED BY GENE NO: 48

- 15 This gene is expressed primarily in amygdala and to a lesser extent in amniotic cells.

- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, depression and other emotional behavioral problems. Similarly,
- 20 polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the nervous system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., brain and tissues of the nervous system, and
- 25 tissues of the reproductive system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid or amniotic fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- 30 The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of mental problems associated with emotional behavior and neurodegenerative states such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder and panic disorders, and
- 35 depression. The amygdala processes sensory information and relays this to other areas of the brain including the endocrine and autonomic domains of the hypothalamus and the brain stem. In addition, expression of this protein in amniotic cells suggests that

this protein would be useful in the diagnosis, prevention, and/or treatment of various developmental and/or reproductive system disorders.

FEATURES OF PROTEIN ENCODED BY GENE NO: 49

5 This gene is expressed primarily in stromal cells.

 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, leukemia and other cancers and disorders deriving from hematopoietic
10 cells. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the lymphoid system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., haematopoietic tissues, and
15 cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid, or lymph fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

20 The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex vivo culture, bone marrow
25 transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia. The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency etc.

30 FEATURES OF PROTEIN ENCODED BY GENE NO: 50

 This gene maps to chromosome 9, and therefore, may be used as a marker in linkage analysis for chromosome 9.

 This gene is expressed primarily in tumors, particularly skin and adrenal gland tumors, and to a lesser extent in bone marrow stromal cells and activated T cells.

35 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are

not limited to, cancer; hematopoietic and immune disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the skin, adrenal gland, and immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., endocrine glands, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 160 as residues: Glu-13 to Arg-22, Ser-58 to Trp-63.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection and treatment of cancer. Elevated levels of expression of this gene in a variety of tumors suggest that it may play a role in cell proliferation, the induction of angiogenesis, destruction of the basal lamina, or a variety of other physiological processes that support the growth and development of tumors and cancer. Alternatively, its expression in the hematopoietic compartment, particularly in the bone marrow stroma and by activated T cells suggest that it may represent a soluble factor capable of influencing a variety of hematopoietic lineages. Therefore, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of blood cells.

25 FEATURES OF PROTEIN ENCODED BY GENE NO: 51

This gene is expressed primarily in benign human breast tissue.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, breast cancer and other female reproductive disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the breast and reproductive tissues, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., breast tissue, secretory/ductile organs, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid or milk) or another tissue or cell

sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- 5 The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and/or diagnosis of breast cancer. Alternately, this protein may play an important role in lactation or represent a critical component secreted into the milk, which may have an important function in the immunoprotection, health, and/or nourishment of the infant upon breastfeeding. Protein, as well as, antibodies directed against the protein may show utility as a tumor
10 marker and/or immunotherapy targets for the above listed tumors and tissues

FEATURES OF PROTEIN ENCODED BY GENE NO: 52

- Translation product of this gene has homology with the conserved human ring finger proteins (See Accession No.gnllPIDle351238 (AJ001019)) which are thought to
15 be important in facilitating and regulating signal transduction pathways in eukaryotic cells. One embodiment for this gene is the polypeptide fragments comprising the following amino acid sequence: HDRTMQDIVYKLVPGGLQE (SEQ ID NO:291) and/or FASHDRTM QDIVYKLVPGGLQEGE (SEQ ID NO:292). An additional embodiment is the polynucleotide fragments encoding these polypeptide fragments.

- 20 This gene is expressed primarily in adult whole brain.

- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, neurodegenerative disorders; Schizophrenia; Alzheimer's; tumors of a
25 brain or neuronal cell origin. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the CNS and/or peripheral nervous system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell
30 types (e.g., brain and other tissue of the nervous system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those
35 comprising a sequence shown in SEQ ID NO. 162 as residues: Phe-39 to Gly-44.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative

disease states and behavioral disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder and panic disorder. In addition, considering the homology to the conserved ring finger proteins may suggest that the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo.

FEATURES OF PROTEIN ENCODED BY GENE NO: 53

Translation product of this gene shares homology with the human conserved Lst-1 gene product, a member of the TNF family of proteins (See Accession No.gil1127546). One embodiment for this gene is the polypeptide fragments comprising the following amino acid sequence: LVLSLGAWGWPSTCLWW (SEQ ID NO:293). An additional embodiment is the polynucleotide fragments encoding these polypeptide fragments.

This gene is expressed primarily in human 6-week old embryo.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, abnormal cell proliferation; defects in terminal tissue differentiation. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the embryo, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., proliferating and differentiating tissues, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid or amniotic fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and/or diagnosis of fetal disorders. Alternately, expression within embryonic tissues may reflect a role for this protein in proliferating cells. In such an event, this gene product may be useful in the treatment or diagnosis of abnormal cell proliferation, such as that involved in cancer. Similarly, embryonic development also involves decisions involving cell differentiation and/or apoptosis involved in pattern formation. Thus, this protein may also be involved in apoptosis or tissue differentiation, and could again be useful in cancer therapy.

FEATURES OF PROTEIN ENCODED BY GENE NO: 54

This gene is expressed primarily in human epithelioid sarcoma.

Therefore, polynucleotides and polypeptides of the invention are useful as
5 reagents for differential identification of the tissue(s) or cell type(s) present in a
biological sample and for diagnosis of diseases and conditions which include, but are
not limited to, epithelial sarcoma; tumors of an epithelial cell origin including the
underlying integument. Similarly, polypeptides and antibodies directed to these
polypeptides are useful in providing immunological probes for differential identification
10 of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells,
particularly of the skin and epithelial tissue layers, expression of this gene at
significantly higher or lower levels may be routinely detected in certain tissues and cell
types (e.g., epithelial cells and tissue, and cancerous and wounded tissues) or bodily
fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or
15 cell sample taken from an individual having such a disorder, relative to the standard
gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an
individual not having the disorder. Preferred epitopes include those comprising a
sequence shown in SEQ ID NO. 164 as residues: Met-1 to Tyr-6, Thr-24 to Cys-36.

The tissue distribution indicates that polynucleotides and polypeptides
20 corresponding to this gene are useful for the treatment and/or diagnosis of epithelial
cancer. This gene product displays enhanced expression in epithelial cell sarcoma, and
thus may be involved in cell proliferation, apoptosis, or in the control of angiogenesis.

FEATURES OF PROTEIN ENCODED BY GENE NO: 55

25 This gene is expressed primarily in endometrial tumors.

Therefore, polynucleotides and polypeptides of the invention are useful as
reagents for differential identification of the tissue(s) or cell type(s) present in a
biological sample and for diagnosis of diseases and conditions which include, but are
not limited to, endometrial cancer including other cancers of the female reproductive
30 system. Similarly, polypeptides and antibodies directed to these polypeptides are useful
in providing immunological probes for differential identification of the tissue(s) or cell
type(s). For a number of disorders of the above tissues or cells, particularly of the
endometrium and reproductive system, expression of this gene at significantly higher or
lower levels may be routinely detected in certain tissues and cell types (e.g.,
35 endometrial tissue as well as other tissues of the female reproductive system, and
cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial
fluid or spinal fluid) or another tissue or cell sample taken from an individual having

such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of cancers, particularly those of the endometrium and other reproductive organs. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tumors and tissues

FEATURES OF PROTEIN ENCODED BY GENE NO: 56

This gene is expressed primarily in metastatic melanoma and to a lesser extent in fetal lung.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, cancer of the integument system, particularly melanoma, as well as within the developing pulmonary system. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the skin, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., cells capable of forming melanin, epithelia, and lung, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid, or pulmonary surfactant) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 166 as residues: Asp-20 to Lys-25.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of cancer, particularly melanoma and more particularly, metastasizing melanomas. In addition, the tissue distribution also indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of cancer and other proliferative disorders. Expression in embryonic tissue and other cellular sources marked by proliferating cells indicates that this protein may play a role in the regulation or cellular division.

FEATURES OF PROTEIN ENCODED BY GENE NO: 57

This gene is expressed primarily in T-cell lymphoma.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, lymphomas and other immune derived cancers. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., T-cells and other cells and tissue of the immune system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 167 as residues: Met-1 to Asn-7.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of lymphomas, particularly T cell lymphomas, and other cancers. In addition, the tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of cancer and other proliferative disorders. Additionally, the expression in hematopoietic cells and tissues indicates that this protein may play a role in the proliferation, differentiation, and/or survival of hematopoietic cell lineages. Thus, this gene may be useful in the treatment of lymphoproliferative disorders, and in the maintenance and differentiation of various hematopoietic lineages from early hematopoietic stem and committed progenitor cells.

FEATURES OF PROTEIN ENCODED BY GENE NO: 58

This gene maps to chromosome 7, and therefore is useful in linkage analysis as a marker for chromosome 7.

This gene is expressed primarily in brain and to a lesser extent in spinal cord.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, CNS and PNS diseases and disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes

for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the nervous system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., brain, spinal cord and other tissue of the nervous system, and
5 cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 168 as residues:
10 Tyr-14 to Ala-30.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and behavioral disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, schizophrenia, mania, dementia, paranoia, obsessive
15 compulsive disorder, panic disorder, and autism.

FEATURES OF PROTEIN ENCODED BY GENE NO: 59

Translation product of this gene shares homology to the conserved *C. elegans* protein FER-1 (See Accession No.gil1373333). One embodiment for this gene is the
20 polypeptide fragments comprising the following amino acid sequence:
QGKLQMWVDVFPKSL (SEQ ID NO:294); PPFNITPRKAKKYYLR (SEQ ID NO:295); KTDVHYRSLDGEGNFNWRP (SEQ ID NO:296); and/or
PRLIIQIWDNDKFSLEDDY LGFLELDL (SEQ ID NO:297). An additional embodiment is the polynucleotide fragments encoding these polypeptide fragments.

25 This gene is expressed primarily in synovial fibroblasts and to a lesser extent in synovial hypoxia.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are
30 not limited to, synovial inflammation and other diseases of the joints. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the synovium, expression of this gene at significantly higher or lower levels may be routinely detected
35 in certain tissues and cell types (e.g., synovial tissue, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to

the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

5 The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of diseases affecting the synovium of the joints, such as rheumatoid arthritis, osteoarthritis, other inflammatory conditions affecting the joints, as well as in the detection and treatment of disorders and conditions affecting the skeletal system, in particular the connective tissues (e.g. trauma, tendonitis, chondromalacia and inflammation). Furthermore, the homology to a conserved *C.elegans* protein may suggest protein is important in human
10 development and thus is beneficial in the diagnosis, prevention, and treatment of developmental disorders.

FEATURES OF PROTEIN ENCODED BY GENE NO: 60

15 This gene is expressed primarily in endothelial cells and to a lesser extent in brain.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, inflammation and other disorders of the integument, in addition to
20 neurodegenerative and nervous system disorder, such as stroke. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the endothelial, circulatory, and nervous systems, expression of this gene at significantly higher or
25 lower levels may be routinely detected in certain tissues and cell types (e.g., endothelial cells, and brain and other tissue of the nervous system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily
30 fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 170 as residues: Ser-4 to Gly-13.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of inflammatory diseases primarily mediated through endothelial cells, such as sepsis, inflammatory
35 bowel disease, psoriasis, and Crohn's disease, as well as for stroke. Alternatively, the tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the detection/treatment of neurodegenerative disease states and

behavioral disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder and panic disorder. In addition, the gene or gene product may also play a role in the treatment and/or detection of developmental disorders associated with the developing embryo, or disorders of the cardiovascular system.

FEATURES OF PROTEIN ENCODED BY GENE NO: 61

This gene is expressed primarily in fetal brain.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, CNS and PNS disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the nervous system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., developing and differentiating tissues, brain and other tissue of the nervous system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid, or amniotic fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of neural disorders such as Alzheimer's disease, depression, paranoia, schizophrenia, autism, and particularly developmental brain disorders..

FEATURES OF PROTEIN ENCODED BY GENE NO: 62

Translation product of this gene shares homology with a conserved 4-nitrophenylphosphatase from *Schizosaccharomyces pombe* (See Accession No. gill1938421). One embodiment for this gene is the polypeptide fragments comprising the following amino acid sequence: AVMIGDDCRDDVGGA (SEQ ID NO:298), and/or ILVKTGKYRASDEEKIN (SEQ ID NO:299). An additional embodiment is the polynucleotide fragments encoding these polypeptide fragments. This gene maps to chromosome 18, and therefore, may be used as a marker in linkage analysis for chromosome 18.

This gene is expressed primarily in endometrial tumors and to a lesser extent in leukemia and lymphoma.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, cancer, particularly of the immune and hematopoietic systems. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the endometrium and white blood cells, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., endometrial and/or proliferating tissues, and cells and tissue of the immune system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid, or lymph) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 172 as residues: Val-19 to Cys-24.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for detection, diagnosis, and treatment of cancers, particularly those cancers affecting endometrial tissues and the lymphatic system. In addition, the tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of hematopoietic related disorders such as anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia. The gene product may also be involved in lymphopoiesis, therefore, it can be used in immune disorders such as infection, inflammation, allergy, immunodeficiency etc. Furthermore, homology to a conserved *S.pombe* protein may suggest protein is important in development. Therefore, protein may be beneficial in the diagnosis, prevention, and treatment of developmental disorders.

FEATURES OF PROTEIN ENCODED BY GENE NO: 63

The translation product of this gene shares sequence homology with ribosomal releasing factor which is thought to be important in protein synthesis.

This gene is expressed primarily in pancreatic tumors, placenta, testis, ovarian cancer, adipocytes, spleen, and fetal liver and heart.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for diagnosis of a number of diseases and conditions such as immune-
5 diseases, cardiovascular and endocrine diseases and others. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, cardiovascular system, digestive system and reproductive system. expression of this
10 gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., pancreas, testis and ovary and other reproductive tissue, adipocytes, spleen, liver, and heart, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid, or lymph) or another tissue or cell sample taken from an individual having such a disorder, relative to the
15 standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 173 as residues: Glu-36 to His-41, Thr-57 to Thr-70, Glu-87 to Met-92, Lys-100 to Lys-105, Ala-197 to Ser-227.

The tissue distribution and homology to ribosomal releasing factor indicates that
20 polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of many diseases, especially cancers and immuno-related diseases.

FEATURES OF PROTEIN ENCODED BY GENE NO: 64

The translation product of this gene shares sequence homology with
25 metalloprotease and also with thrombospondin, which is thought to be important in the activation of proteins and the processes of thrombopoiesis and metabolism.

This gene is expressed in many tissues, but especially in bladder, kidney, and ovary.

Therefore, polynucleotides and polypeptides of the invention are useful as
30 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of thrombopenia, hypertension, and other blood disfunctions. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of
35 the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., urogenital, and reproductive tissues, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma,

urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID

5 NO. 174 as residues: Gly-8 to Leu-14, Met-18 to Phe-30.

The tissue distribution and homology to thrombospondin indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of a variety of blood-related diseases.

10 **FEATURES OF PROTEIN ENCODED BY GENE NO: 65**

This gene is expressed primarily in tonsil, placenta, and fetal tissues.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of many diseases of the immune system. Similarly, 15 polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., immune and developmental tissues, and cancerous and wounded tissues) or bodily 20 fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid, or amniotic fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides 25 corresponding to this gene are useful for diagnosis and treatment of diseases of the immune system including many cancers such as lymphomas, leukemias, lymphocytomas, and the like.

FEATURES OF PROTEIN ENCODED BY GENE NO: 66

30 Polypeptides encoded by this gene share reasonable homology to steroid/thyroid hormone orphan nuclear receptor and to several additional orphan nuclear receptors isolated from several different tissues.

This gene is expressed primarily in testis.

Therefore, polynucleotides and polypeptides of the invention are useful as 35 reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of testicular tumors, impotence, and other reproductive disorders. Similarly, polypeptides and antibodies directed to these

polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the reproductive system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., male reproductive tissue, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid, or seminal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- 5 The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of diseases in the male reproductive system such as tumors of the testis and other reproductive disorders.

FEATURES OF PROTEIN ENCODED BY GENE NO: 67

- 15 Polypeptides encoded by polynucleotides comprising this gene have a high degree of sequence identity with CTGF-4.

In one embodiment, the polypeptides of the invention comprise the sequence: MDSMPEPASRCLLLLPLLLLLLLLLPAPELGPSQAGAEENDWVRLPSK
CEVCKYVAVELKVKPLRKRQDTEVIGTVYGILDQKASGVKYTKSDLRLIEVTET
20 ICKRLLDYSLHKERTGSXRFAKGMSETFETLHXLVHKGVKVVM DIPYELWNE
TSAEVADLKKQCDVLVEEFEEVIEDWYRNHQEEDLTEFLCANHVLK GKDTSCLEQWSGKKGDTAALGGKSKKK SIRAKAAGGRSSSSKQRKELGGLEGDPSP
EEDEGIQKASPLTHSPPEL (SEQ ID NO:300). Polynucleotides encoding these polypeptide sequences are also encompassed by the invention.

- 25 This gene is expressed in many tissues especially including cells in the immune system.

- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for the diagnosis of cancers, immunological disorders, and neural diseases (such as spinocerebellar ataxia, bipolar affective disorder, schizophrenia, and autism), and other diseases featuring anticipation, neurodegeneration, or abnormalities of neurodevelopment. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the nerve system, immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., immune cells and/or tissue, and cancerous and wounded tissues) or bodily

fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid, or lymph) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 177 as residues: Ser-3 to Ser-9, Gly-36 to Val-43, Leu-45 to Gly-51.

FEATURES OF PROTEIN ENCODED BY GENE NO: 68

Polypeptides encoded by polynucleotides comprising this gene contain a zinc finger homology domain. Such motifs are believed to be important for protein interactions, particularly with regard to gene regulation.

This gene is expressed primarily in T cells and the colon and, to a lesser extent, in the testes and placenta.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of many immune and digestive disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and digestive systems, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., immune, gastrointestinal, and reproductive system tissues, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid, or seminal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 178 as residues: Pro-12 to Lys-33, Asn-41 to His-46, Pro-48 to Ser-58, Gly-71 to Asp-78, Ala-94 to Gly-102, Ser-133 to Ser-140, Arg-197 to Lys-202.

The expression of this gene in T-cells indicates a potential role in the treatment and detection of immune disorders such as arthritis, asthma, immune deficiency diseases (such as AIDS), and leukemia. Expression of this gene in the colon indicates a potential role in the treatment and detection of colon disorders such as ulcers and colon cancer in addition to digestive disorders in general.

FEATURES OF PROTEIN ENCODED BY GENE NO: 69

The translation product of this gene shares sequence homology with neuroendocrine protein which is thought to be important in neuronal development and differentiation. A preferred embodiment of this gene comprises the following amino acid sequence: MDGQKKNWKDKVVDLLYWRDIKKTGVVFGASLFLLSLTVF
 5 SIVSVTAYIALALLSVTISFRIYKGVIAIQKSDEGHPFRAYLESEVAISEELVQKY
 SNSALGHVNCTIKELRRLFLVDDLVDLKFVAVLMWVFTYVGALFNGLTLLILAL
 ISLFSVPVIYERHQAQIDHYLGLANKNVKDAMAKIQAKIPGLKRKAE (SEQ ID
 NO:301). Particularly preferred are polynucleotides comprising polynucleotides
 10 encoding this polypeptide sequence.

This gene is expressed in many different tissues, but primarily in brain, and, to a lesser extent, in fetal tissue, placenta, bone marrow, and stromal cells.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for diagnosis of neurodegenerative diseases and developmental disorders.
 15 Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the nervous system and during development, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., neural, developmental, and hemopoietic cells and tissue, and cancerous and wounded tissues)
 20 or bodily fluids (e.g., serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those
 25 comprising a sequence shown in SEQ ID NO. 179 as residues: Gln-47 to Gly-52, Leu-169 to Glu-174.

The predominant tissue distribution in brain and homology to neuroendocrine protein indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of neurodegenerative diseases and behavioral
 30 disorders such as Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, schizophrenia, mania, dementia, paranoia, obsessive-compulsive disorder and panic disorder.

FEATURES OF PROTEIN ENCODED BY GENE NO: 70

35 Polypeptides encoded by polynucleotides comprising this gene share sequence identity with human hepatoma-derived growth factor (WPI 95-069304/10). As such, polynucleotides comprising this gene can be used for the recombinant production of the

protein, which can be used to encourage the growth of various animal cells, and for the purification of receptors. Additional embodiments of the invention comprise the following polypeptide sequences: MAVTSLLLGGRVCA (SEQ ID NO:302); PSLAVGSRPGGW RAQALLAGSRTPIPTGSRNGSCRRWRAP (SEQ ID NO:303); and/or MAVTSLLLGGRVCAPSLAVGSRPGGWRAQALLAGSRTPIPTGSRNGSCRRWRAP (SEQ ID NO:304). Also contemplated are polynucleotides comprising polynucleotides encoding the aforementioned polypeptide sequences.

This gene is expressed primarily in brain and to a lesser extent in endothelium, T- cell, and tumors.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of many neurodegenerative diseases (for example, Alzheimer's Disease, ALS, and the like) and cancers (including, but not limited to neuroblastoma, glioblastoma, Schwannoma, astrocytoma, and the like). Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the nervous system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., neural, and haematopoietic cells and tissue, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid or lymph) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 180 as residues: Pro-4 to Thr-10, Glu-25 to Trp-30, Leu-58 to Leu-69, Arg-82 to Thr-87, Ala-108 to His-115, Ser-124 to Glu-146, Pro-159 to Gly-176, Ser-182 to Glu-187, Leu-189 to Ser-198, Phe-208 to Asn-214.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment and diagnosis of many neurodegenerative diseases and cancers.

FEATURES OF PROTEIN ENCODED BY GENE NO: 71

The translation product of this gene shares sequence homology with acrosin, trypsin, as well as trypsinogen precursor which are thought to be important in cell-cell recognition and proteinase activity for protein cleavage and degradation. Preferred polynucleotide fragments comprise the following sequence:

GATGTTACACAGCTCTTTAATAATAGTGGCCATAGCTGTAATAACAATGACA

ACAGTAGGTAACGGTAGTCATACCAACAGTAGGGCAGTGCATTTTATATTAC
AACTGGTTTTCTTGCTCTAGTAGGCTTGGGGATGGGTGAAGACGGACAGGGC
TGGCGCAGACCCTTTCCTTCTCCTCTCCAGCCCACAGTGATCTGGGCTTTTA
CAGACAGCCTGCTTCCATTCACTAGTGTGGGAAAGTTCCTTCTTGGCTTAGC
5 AATACCCCTGAGACCTTGTTTCAGTGGGCTGTGTCTCTCCCTGGGATGCTGG
GAGCACCAAGTGTGGCCGAGCTAGGGCTGCTGACTTCCTCTGGGCGCCTCT
GGGCTGCGAGGGTCTCTTATAGGAATTGAGGCCCTTTGCTGCTCCAAGAAA
TGCGAGGCTGTGGGCARAGGGKTGTACCCAAGGGGACTCTTGCTCTGTGT
CTGACTTTGGGGRATCC (SEQ ID NO:305); CACAGCTCTTTAATAATAGTGGC
10 CATAGCTGTAATAACAATGACA ACAGTAGGTAACG (SEQ ID NO:306);
TGTGTCTCTCCCTGGGATGCTGGGAGCACCAAGTGTGGCCGAGCTAGGGCT
GCTGACTT (SEQ ID NO:307); GCGAGGGTCTCTTATAGGAATTGAGGCCCTT
TGCTGCTCCAAGAAATGCTGAGGCTGTGGGCARAGGGKTGTACCCAAGGG
GACT (SEQ ID NO:308). Also preferred are polypeptide fragments encoded by these
15 polynucleotide fragments.

This gene is expressed primarily in cheek carcinoma and to a lesser extent in uterine and pancreatic cancers.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a
20 biological sample and for diagnosis of diseases and conditions which include, but are not limited to, cheek cancers or cancers of uterine and pancreatic origins. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the neoplastic tissues,
25 expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., epithelial, endocrine, and reproductive tissues, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid, and saliva) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e.,
30 the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution and homology to acrosin and trypsin indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and intervention of cancers. The homology to acrosin and trypsin may indicate the gene
35 function in tumor metastasis or migration since in both cases cell-cell interaction and extracellular matrix degradation may be involved. The gene product can also be used as a target for cancer immunotherapy or as a diagnostic marker.

FEATURES OF PROTEIN ENCODED BY GENE NO: 72

This gene is expressed primarily in T helper cells I, T-cells stimulated with PHA for 24 hours, and in a placenta Nb2HP cDNA library.

5 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of many immunodeficiencies and disorders (especially autoimmune diseases). Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential
10 identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., immune, and haematopoietic cells and tissue, and cancerous and wounded tissue) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid and
15 lymph) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of autoimmune
20 diseases, immunodeficiencies, and other immune system disorders.

FEATURES OF PROTEIN ENCODED BY GENE NO: 73

This gene is expressed primarily in 7 week old early stage human, human chronic synovitis, and infant brain.

25 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of chronic synovitis. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders
30 of the above tissues or cells, particularly of the synovium, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., developmental, differentiating, and neural tissues, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid, and amniotic fluid) or another tissue or cell sample taken from an individual
35 having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the

disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 183 as residues: Ser-44 to Pro-49.

- The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of chronic synovitis and other disorders of the synovium.

FEATURES OF PROTEIN ENCODED BY GENE NO: 74

- Polypeptides encoded by polynucleotides comprising this gene exhibit sequence homology to a number of mucin-like extracellular or cell surface proteins. In one embodiment polypeptides of the invention comprise the following sequence:
- MVGPVTLHKKIHITTTVLFIVQIHILLIQAITQAK (SEQ ID NO:309); LQMHLMLQMTGLSILALLGKSTTTIVEQKFHNGKNQKSGLKENRDKKKQTRWQSTASQKIGITEER (SEQ ID NO:310); and/or MVGPVTLHKKIHITTTVLFIVQIHILLIQAITQAKLQMHLMLQMTGLSILALLGKSTTTIVEQKFHNGKNQKSGLKENRDKKKQTRWQSTASQKIGITEER (SEQ ID NO:311). Polynucleotides encoding the

aforementioned polypeptides are also contemplated embodiments of the invention.

This gene is expressed primarily in ovarian cancer, endometrial tumor, B-cell lymphoma, brain-medulloblastoma, hepatocellular tumor, osteosarcoma, and T- and B-cells.

- Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, Ovarian cancer, endometrial tumor, B-cell lymphoma, brain medulloblastoma, hepatocellular tumor, and osteosarcoma. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., brain and other tissue of the nervous system, bone, T-cells and other cells of the immune system, and B cells and other blood cells, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid and lymph) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 184 as residues: Met-1 to Lys-12, Leu-14 to Asn-35, Arg-42 to Asn-58, Ser-65 to Trp-90, Ser-95 to Asn-129, Phe-136 to Arg-144, Met-159 to Ala-167, Thr-179 to Tyr-187, Pro-190 to

Val-201, Gln-226 to Phe-235, Pro-254 to His-272, Thr-288 to Thr-293, Thr-383 to Ser-391, Asp-398 to Tyr-405, Ile-410 to Asn-416, Ala-449 to Lys-458.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of ovarian cancer, endometrial tumors, B-cell lymphoma, brain medulloblastoma, hepatocellular tumor, and osteosarcoma.

FEATURES OF PROTEIN ENCODED BY GENE NO: 75

An additional preferred polypeptide sequence derived from the polynucleotide of this contig comprises the following amino acid sequence: MQTCPLVGTLNMDG YTCVVSTSTFSWIISAWXLWKGSPSTSMPTMPETPLRLCCTKMPSIFSSLMTD GRA (SEQ ID NO:312). Polynucleotides encoding these polypeptides are also provided. This polypeptide sequence has sequence homology with a *Drosophila melanogaster* male germ-line specific transcript which encodes a putative protamine molecule (see, gil608696).

This gene is expressed primarily in breast tissue and to a lesser extent in various other fetal and adult cells and tissues, especially those comprising endocrine organs.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, developmental and reproductive defects. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the female reproductive system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., breast and/or other ductile secretory tissues, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid, and milk) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for study and treatment of developmental, reproductive and growth and metabolic disorders.

35 FEATURES OF PROTEIN ENCODED BY GENE NO: 76

In one embodiment, the polypeptides of the invention comprise the sequence: MTLIQNCWYSWLFFGFFFHFLRKSISIFSIFLVCFRILALGPTCFLVWFWKAFFR

HILIFICLSREVFPRCFLVYFR (SEQ ID NO:313). This polypeptide sequence has sequence homology with the MURF4 protein of *Herpetomonas muscarum* (S43288). Such RNA-editing enzymes may be useful as molecular targets in the intervention of the life cycle of trypanosomes and other protozoa. Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in fetal liver and spleen, osteosarcoma and bone marrow.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of liver tumors, osteosarcoma, and other cancers. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., hepatic, developmental, and differentiating tissue, bone cells, liver and spleen, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid, and lymph) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis of cancers such as liver tumor and osteosarcoma.

25 FEATURES OF PROTEIN ENCODED BY GENE NO: 77

This gene is expressed primarily in T cell lymphoma and monocytes.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of T-cell lymphoma. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., immune and hematopoietic cells and tissues, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid, and lymph) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in

healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 187 as residues: Thr-1 to Ser-9.

- 5 The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis and treatment of T-cell lymphoma.

FEATURES OF PROTEIN ENCODED BY GENE NO: 78

This gene is expressed primarily in tonsils and a bone marrow cell line.

- 10 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of immunological disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system,
- 15 expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., haematopoietic and immune cells and tissues, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid, and lymph) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the
- 20 expression level in healthy tissue or bodily fluid from an individual not having the disorder.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of immunological disorders.

25

FEATURES OF PROTEIN ENCODED BY GENE NO: 79

- In one embodiment, the polypeptides of the invention comprise the sequence: MGTRAQVTPGRLPIPPAPGLPFSAXEPLQGQLRRVSSSRGGFPGLALQLLRSE TVKAYVNNEINILASFF (SEQ ID NO:314) and/or MLVRTRPSQPLPLPGVGLGGP
- 30 RSGDPPESTELRKGPGLA (SEQ ID NO:315). Polynucleotides encoding these polypeptides are also encompassed by the invention.

This gene is expressed primarily in brain, placenta, bone marrow, keratinocyte, fetal liver, and spleen.

- 35 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of brain and skin related diseases. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing

immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune and skin system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., neural, reproductive, and hepatic tissues, 5 keratinocytes, and spleen, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a 10 sequence shown in SEQ ID NO. 189 as residues: Phe-13 to Leu-18.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the diagnosis and treatment of many brain and skin related diseases.

15 **FEATURES OF PROTEIN ENCODED BY GENE NO: 80**

The translation product of this gene shares sequence homology with mouse RNA Polymerase I which is thought to be important in gene transcription process.

This gene is expressed primarily in HEL cell line and aorta endothelial cells and to a lesser extent in Jurkat T-cells.

20 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis and treatment of cancer and autoimmune diseases. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell 25 type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., endothelial, haematopoietic tissues, cardiovascular tissue, and T-cells and other cells of the immune system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial 30 fluid, spinal fluid, and lymph) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 190 as residues: Lys-25 to Arg-32.

35 The tissue distribution and homology to mouse RNA polymerase I indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment of immune diseases and cardiovascular diseases.

FEATURES OF PROTEIN ENCODED BY GENE NO: 81

In one embodiment, the polypeptides of the invention comprise the sequence:
MCPVCGRALSSPGSLGRHLLIHSEDQRSNCAVCGARFTSHATFNSEKLPEVLN
5 MESLPTVHNEGPPSSAEGKDIAFSPVYPAGILLVCNNCAA YRKXLEAQTSPVX
KWALRRQNEPLEVRLQRLERERTAKKSRRDNETPEEREVRRMRDREAKRLQR
MQETDEQRARRLQRDREAMRLKRANETPEKRQARLIREREAKRLKRRLEKMD
MMLRAQFGQDPSAMAALAAEMNFFQLPVSGVELDXQLLGKMAFEEQNSSXLH
(SEQ ID NO:316). This polypeptide shares sequence homology with human trichohylin
10 which is thought to be important in gene regulation. Polynucleotides encoding this
polypeptide are also encompassed by the invention.

This gene is expressed primarily in brain tissue and to a lesser extent in
apoptotic T-cell and B-cell lymphoma.

Therefore, polynucleotides and polypeptides of the invention are useful as
15 reagents for differential identification of the tissue(s) or cell type(s) present in a
biological sample and for diagnosis and treatment of growth disorders,
neurodegenerative diseases, and endocrine disorders. Similarly, polypeptides and
antibodies directed to these polypeptides are useful in providing immunological probes
for differential identification of the tissue(s) or cell type(s). For a number of disorders
20 of the above tissues or cells, particularly of the neural and immune systems, expression
of this gene at significantly higher or lower levels may be routinely detected in certain
tissues and cell types (e.g., neural tissues, T-cells, B-cells and other cells and tissue of
the immune system, and cancerous and wounded tissues) or bodily fluids (e.g., serum,
plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from
25 an individual having such a disorder, relative to the standard gene expression level, i.e.,
the expression level in healthy tissue or bodily fluid from an individual not having the
disorder.

The tissue distribution and homology to DNA binding protein indicates that
polynucleotides and polypeptides corresponding to this gene are useful for the
30 diagnosis and treatment of immune and neurological diseases.

FEATURES OF PROTEIN ENCODED BY GENE NO: 82

In one embodiment, the polypeptides of the invention comprise the sequence:
MDHSHHMGMSYMDSNSTMQPSHHPTTSASHSHGGDSSMMMMPMTFYFG
35 FKNVELLFSGLVINTAGEMAGAFVAVFLLAMFYEGLKIARESLLRKSQVSIRYN
SMPVPGPNGTILMETHKTVGQQMLSFPHLLQTVLHIIQVVISYFLMLIFMTYNG
YLCIAXAAGAGTGYFLFSWKKA VVVDITEHCH (SEQ ID NO:317). This

polypeptide is thought to function in mediating the uptake of copper and other metal ions by cells. Polynucleotides encoding this polypeptide are also encompassed by the invention.

5 This gene is expressed primarily in osteosarcoma and to a lesser extent in T-cell and bone marrow stromal cell.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for treatment and diagnosis of osteosarcoma and copper and other metal uptake disorders. Similarly, polypeptides and antibodies directed to these
10 polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., hematopoietic tissue and cancerous and wounded tissues) or bodily fluids (e.g.,
15 serum, plasma, urine, synovial fluid, spinal fluid, and lymph) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 192 as residues: Ser-24 to Ser-29.

20 The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the prevention or treatment of osteosarcoma and copper or other metal uptake disorders.

FEATURES OF PROTEIN ENCODED BY GENE NO: 83

25 This gene is expressed primarily in skin tumor and to a lesser extent in apoptic T-cell.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are
30 not limited to, skin tumor. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the skin, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., epithelial and
35 hematopoietic tissues, and T-cells and other tissue of the immune system, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, and spinal fluid) or another tissue or cell sample taken from an individual having

such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 193 as residues: Leu-51 to Gly-77, Ile-117 to Pro-125.

- 5 The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for diagnosis the treatment of skin tumor.

FEATURES OF PROTEIN ENCODED BY GENE NO: 84

This gene is expressed primarily in testis.

- 10 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, infertility and endocrine disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes
- 15 for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the reproductive system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., reproductive tissue, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid, and seminal fluid) or
- 20 another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder.

- The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment of reproductive disease and
- 25 endocrine disorders.

FEATURES OF PROTEIN ENCODED BY GENE NO: 85

- In one embodiment, the polypeptides of the invention comprise the sequence:
- 30 MVQPCGACAKTXWKACSSCCSSPCCLQERWPXPXAXCPEXGPSSHPGIQALC
AVAVVYLSPSSRLDWSLAPLFVPSLAAGETPLTQPAWALTNTLGHGQPAQDR
LPALGHCAPISVLGLGSS (SEQ ID NO:318). Polynucleotides encoding this polypeptide sequence are also encompassed by the invention.

 This gene is expressed primarily in kidney cortex, frontal cortex, spinal cord and hippocampus.

- 35 Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are

not limited to, kidney fibrosis, schizophrenia and neurological disorders. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the neural system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., endothelial, neural and endocrine tissue, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder. Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 195 as residues: Cys-27 to Tyr-33, Thr-38 to Gly-43, Leu-125 to Gly-130.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment of neurological disorders and kidney diseases..

FEATURES OF PROTEIN ENCODED BY GENE NO: 86

This gene is expressed primarily in resting T-cell.

Therefore, polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include, but are not limited to, T-cell related diseases. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the immune system, expression of this gene at significantly higher or lower levels may be routinely detected in certain tissues and cell types (e.g., hematopoietic and immune cells and tissues, and cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid, spinal fluid, and lymph) or another tissue or cell sample taken from an individual having such a disorder, relative to the standard gene expression level, (i.e., the expression level in healthy tissue or bodily fluid from an individual not having the disorder). Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 196 as residues: Thr-54 to Ile-59.

The tissue distribution indicates that polynucleotides and polypeptides corresponding to this gene are useful for the treatment of immune diseases.

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
1	HOAAE80	209012 04/28/97 209089 06/05/97	Uni-ZAP XR	11	1220	264	1220	288	288	111	1	26	27	31
2	HODDN92	209012 04/28/97 209089 06/05/97	Uni-ZAP XR	12	1939	294	1939	434	434	112	1	26	27	35
3	HOSBI96	209012 04/28/97 209089 06/05/97	Uni-ZAP XR	13	2602	672	1811	690	690	113	1	30	31	219
4	HOVAI58	209012 04/28/97 209089 06/05/97	pSport1	14	808	1	808	28	28	114	1	26	27	31
5	HPBDD36	209012 04/28/97 209089 06/05/97	pBluescript SK-	15	864	87	831	147	147	115	1	18	19	26
6	HPDDC77	209012 04/28/97 209089 06/05/97	pBluescript SK-	16	2361	455	1442	510	510	116	1	29	30	131
7	HPBED85	209012	Uni-ZAP XR	17	803	1	803	81	81	117	1	20	21	64

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
		04/28/97 209089 06/05/97												
8	HPFCX38	209012 04/28/97 209089 06/05/97	Uni-ZAP XR	18	1794	1051	1757		578	118	1			8
9	HPFCY51	209012 04/28/97 209089 06/05/97	Uni-ZAP XR	19	1037	1	1037	467	467	119	1	30	31	50
9	HPFCY51	209012 04/28/97 209089 06/05/97	Uni-ZAP XR	97	1052	1	1052	30	30	197	1			13
10	HPMGQ80	209012 04/28/97 209089 06/05/97	Uni-ZAP XR	20	1309	157	1309	360	360	120	1	19	20	76
11	HPRTG55	209012 04/28/97 209089 06/05/97	pBluescript	21	1081	55	1014	237	237	121	1	24	25	26
12	HROAN56	209012 04/28/97 209089 06/05/97	Uni-ZAP XR	22	807	1	807	26	26	122	1	19	20	23

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
13	HSAB142	209012 04/28/97 209089 06/05/97	pBluescript SK-	23	632	1	596	190	190	123	1	15	16	21
14	HSAUW44	209012 04/28/97 209089 06/05/97	Uni-ZAP XR	24	1358	1	1358	372	372	124	1	30	31	54
15	HSDES04	209012 04/28/97 209089 06/05/97	Uni-ZAP XR	25	1376	686	1376	146	146	125	1	33	34	318
15	HSDES04	209012 04/28/97 209089 06/05/97	Uni-ZAP XR	98	929	57	929	291	291	198	1	28	29	61
16	HSHBQ68	209012 04/28/97 209089 06/05/97	Uni-ZAP XR	26	2923	195	2642	211	211	126	1	23	24	58
17	HSKBO20	209012 04/28/97 209089 06/05/97	Uni-ZAP XR	27	775	1	501		308	127	1	28	29	98
18	HSKNM85	209012 04/28/97 209089	pBluescript	28	534	1	534	122	122	128	1	19	20	28

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
19	HSKXJ37	209012 04/28/97 209089 06/05/97	pBluescript	29	1827	67	1634	311	129	1	21	22	21
20	HSKZE52	209012 04/28/97 209089 06/05/97	Uni-ZAP XR	30	1479	418	1453	555	130	1	18	19	111
21	HWTAZ75	209012 04/28/97 209089 06/05/97	Uni-ZAP XR	31	987	448	963	133	131	1	1	2	114
22	HSRBA90	209012 04/28/97 209089 06/05/97	Uni-ZAP XR	32	2933	1437	2933	1670	132	1	19	20	21
23	HSVAG05	209090 06/05/97	Uni-ZAP XR	33	1366	1	1366	66	133	1	31	32	51
24	HSVBF78	209090 06/05/97	Uni-ZAP XR	34	667	141	621	64	134	1	28	29	99
25	HSXBO51	209090 06/05/97	Uni-ZAP XR	35	1710	388	1683	462	135	1	26	27	175
26	HT3BE24	209090 06/05/97	Uni-ZAP XR	36	1096	756	1091	422	136	1	15	16	187
26	HT3BE24	209090 06/05/97	Uni-ZAP XR	99	359	1	359	41	199	1	42	43	71

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
27	HT4AI54	209090 06/05/97	Uni-ZAP XR	37	2279	1387	2279	29	137	1	24	25	288
27	HT4AI54	209090 06/05/97	Uni-ZAP XR	100	952	1	952	199	200	1			10
28	HTEHU93	209090 06/05/97	Uni-ZAP XR	38	745	1	745	187	138	1	24	25	113
29	HTGCQ82	209090 06/05/97	Uni-ZAP XR	39	1718	70	1718	114	139	1	23	24	119
30	HTLAB25	209090 06/05/97	Uni-ZAP XR	40	1966	321	1966	449	140	1	1	2	438
31	HTLAV68	209090 06/05/97	Uni-ZAP XR	41	972	1	972	78	141	1	35	36	162
32	HTLDQ11	209090 06/05/97	Uni-ZAP XR	42	1536	1	1536	213	142	1	36	37	72
33	HTOBX52	209090 06/05/97	Uni-ZAP XR	43	2541	1743	2541	3	143	1	4	5	123
34	HTTCN24	209090 06/05/97	Uni-ZAP XR	44	2418	918	2290	188	144	1	30	31	138
34	HTTCN24	209090 06/05/97	Uni-ZAP XR	101	1545	123	1545	345	201	1	39	40	50
35	HTXCS21	209090 06/05/97	Uni-ZAP XR	45	1337	657	1309	76	145	1	24	25	356
35	HTXCS21	209090 06/05/97	Uni-ZAP XR	102	1322	641	1293		202	1			13
36	HUFAC49	209090 06/05/97	pSport1	46	1276	1	1276	105	146	1	17	18	39

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
37	HAIDK60	209090 06/05/97	Uni-ZAP XR	47	1282	1	1282	528	528	147	1	30	31	71
37	HAIDK60	209090 06/05/97	Uni-ZAP XR	103	276	1	276	14	14	203	1	25	26	38
38	HARAG28	209090 06/05/97	pBluescript SK-	48	645	1	645	150	150	148	1	16	17	33
38	HARAG28	209090 06/05/97	pBluescript SK-	104	381	1	381	154	154	204	1	18	19	34
39	HBMBB80	209090 06/05/97	pBluescript	49	1495	2	1495	23	23	149	1	30	31	78
39	HBMBB80	209090 06/05/97	pBluescript	105	638	1	638	196	196	205	1	16	17	26
40	HCEGR33	209090 06/05/97	Uni-ZAP XR	50	1630	1	1630	243	243	150	1	22	23	31
41	HSXBP68	209090 06/05/97	Uni-ZAP XR	51	2420	1009	2252	79	79	151	1	41	42	464
41	HSXBP68	209090 06/05/97	Uni-ZAP XR	106	2246	835	2079	985	985	206	1	32	33	105
42	HFFAT33	209090 06/05/97	Lambda ZAP II	52	1172	166	802	209	209	152	1	29	30	151
43	HFGAG96	209090 06/05/97	Uni-ZAP XR	53	1589	885	1446	189	189	153	1	33	34	299
43	HFGAG96	209090 06/05/97	Uni-ZAP XR	107	1105	1	1105		247	207	1	17	18	64
44	HETFJ05	209076 05/22/97	Uni-ZAP XR	54	2074	1	2065	75	75	154	1	24	25	397

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
45	HLTEY63	209076 05/22/97	Uni-ZAP XR	55	1483	1	1280	86	86	155	1	18	19	82
46	HMSJU68	209076 05/22/97	Uni-ZAP XR	56	1123	4	1123	272	272	156	1	31	32	49
47	HOSCZ41	209076 05/22/97	Uni-ZAP XR	57	1239	117	1222	178	178	157	1	20	21	50
48	HSHAV28	209076 05/22/97	Uni-ZAP XR	58	803	105	719		378	158	1			16
49	HSQEA85	209076 05/22/97	Uni-ZAP XR	59	995	1	995	98	98	159	1	23	24	52
50	HSTAG52	209076 05/22/97	Uni-ZAP XR	60	966	114	966	191	191	160	1	45	46	63
51	HBNAJ22	209076 05/22/97	Uni-ZAP XR	61	262	1	262	28	28	161	1	23	24	32
52	HBXGP76	209076 05/22/97	ZAP Express	62	753	1	753	34	34	162	1	34	35	94
53	HE6GL64	209076 05/22/97	Uni-ZAP XR	63	739	1	739	132	132	163	1	32	33	57
54	HESAL35	209076 05/22/97	Uni-ZAP XR	64	476	1	476	20	20	164	1	27	28	43
55	HETBB70	209076 05/22/97	Uni-ZAP XR	65	754	14	754		263	165	1	17	18	17
56	HLHAY19	209076 05/22/97	Uni-ZAP XR	66	1890	8	1890	18	18	166	1	22	23	28
57	HLTER45	209076 05/22/97	Uni-ZAP XR	67	1614	557	1614	578	578	167	1	25	26	36

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
58	HNHAL34	209076 05/22/97	Uni-ZAP XR	68	596	1	596	90	90	168	1	18	19	39
59	HOSFF78	209076 05/22/97	Uni-ZAP XR	69	1524	791	1524	846	846	169	1	34	35	46
60	HSKDV92	209076 05/22/97	Uni-ZAP XR	70	819	53	819		158	170	1	32	33	33
61	HFCCU63	209076 05/22/97	Uni-ZAP XR	71	1442	1	1442	12	12	171	1			4
62	HLTCS34	209076 05/22/97	Uni-ZAP XR	72	1223	1	1223	227	227	172	1	17	18	24
63	HPMCC16	209086 05/29/97	Uni-ZAP XR	73	1814	1024	1814	85	85	173	1	19	20	262
64	HOUQC17	209086 05/29/97	Uni-ZAP XR	74	4712	1	4693	508	508	174	1	51	52	967
65	HTDAG66	209086 05/29/97	pSport1	75	1885	262	1885	369	369	175	1			18
66	HTLBC79	209086 05/29/97	Uni-ZAP XR	76	890	1	890	17	17	176	1	1	2	205
67	HTOFC34	209086 05/29/97	Uni-ZAP XR	77	1657	356	1645	434	434	177	1	31	32	54
68	H2CBJ08	209086 05/29/97	pBluescript SK-	78	2015	13	2015	70	70	178	1	17	18	435
69	HAGFT48	209086 05/29/97	Uni-ZAP XR	79	1213	242	1213		290	179	1	23	24	174
70	HCE5M29	209086 05/29/97	Uni-ZAP XR	80	1391	23	1353	251	251	180	1	1	2	219

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
71	HTPBQ83	209076 05/22/97	Uni-ZAP XR	81	1008	146	1008		431	181	1			5
72	HCFNN01	209086 05/29/97	pSport1	82	1261	154	1261	254	254	182	1	27	28	43
73	HE7TF86	209086 05/29/97	Uni-ZAP XR	83	1045	241	986	426	426	183	1	23	24	58
74	HGBAC11	209086 05/29/97	Uni-ZAP XR	84	2877	1	2272	85	85	184	1	1	2	588
75	HHGAU81	209086 05/29/97	Lambda ZAP II	85	1367	747	1367	323	323	185	1	24	25	166
76	HLCAA05	209086 05/29/97	Uni-ZAP XR	86	1009	1	1009	276	276	186	1			8
77	HMSCD68	209086 05/29/97	Uni-ZAP XR	87	1367	1	1367		254	187	1			19
78	HMWDZ81	209086 05/29/97	Uni-Zap XR	88	1088	1	883	214	214	188	1	22	23	30
79	HMWQG73	209086 05/29/97	Uni-Zap XR	89	1861	875	1861		1160	189	1	15	16	18
80	HOECN31	209086 05/29/97	Uni-ZAP XR	90	1259	34	1259	338	338	190	1	28	29	32
81	HPTRF90	209086 05/29/97	pBluescript	91	1566	450	1552	593	593	191	1	28	29	83
82	HSRDH01	209086 05/29/97	Uni-ZAP XR	92	1593	107	1593	379	379	192	1	22	23	122
83	HSAWD74	209126 06/19/97	Uni-ZAP XR	93	970	106	970	142	142	193	1	26	27	142

Gene No.	cDNA Clone ID	ATCC Deposit Nr and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
83	HSTBE27	209086 05/29/97	Uni-ZAP XR	110	646	117	646	122	122	210	1	31	32	46
84	HTEJO12	209086 05/29/97	Uni-ZAP XR	94	934	1	934	202	202	194	1	20	21	50
85	HTLAB43	209086 05/29/97	Uni-ZAP XR	95	1392	199	1392	384	384	195	1	17	18	221
86	HTWCT03	209086 05/29/97	pSport1	96	1963	1	1963	334	334	196	1	26	27	101

Table 1 summarizes the information corresponding to each "Gene No." described above. The nucleotide sequence identified as "NT SEQ ID NO:X" was assembled from partially homologous ("overlapping") sequences obtained from the "cDNA clone ID" identified in Table 1 and, in some cases, from additional related DNA clones. The overlapping sequences were assembled into a single contiguous sequence of high redundancy (usually three to five overlapping sequences at each nucleotide position), resulting in a final sequence identified as SEQ ID NO:X.

The cDNA Clone ID was deposited on the date and given the corresponding deposit number listed in "ATCC Deposit No:Z and Date." Some of the deposits contain multiple different clones corresponding to the same gene. "Vector" refers to the type of vector contained in the cDNA Clone ID.

"Total NT Seq." refers to the total number of nucleotides in the contig identified by "Gene No." The deposited clone may contain all or most of these sequences, reflected by the nucleotide position indicated as "5' NT of Clone Seq." and the "3' NT of Clone Seq." of SEQ ID NO:X. The nucleotide position of SEQ ID NO:X of the putative start codon (methionine) is identified as "5' NT of Start Codon." Similarly, the nucleotide position of SEQ ID NO:X of the predicted signal sequence is identified as "5' NT of First AA of Signal Pep."

The translated amino acid sequence, beginning with the methionine, is identified as "AA SEQ ID NO:Y," although other reading frames can also be easily translated using known molecular biology techniques. The polypeptides produced by these alternative open reading frames are specifically contemplated by the present invention.

The first and last amino acid position of SEQ ID NO:Y of the predicted signal peptide is identified as "First AA of Sig Pep" and "Last AA of Sig Pep." The predicted first amino acid position of SEQ ID NO:Y of the secreted portion is identified as "Predicted First AA of Secreted Portion." Finally, the amino acid position of SEQ ID NO:Y of the last amino acid in the open reading frame is identified as "Last AA of ORF."

SEQ ID NO:X and the translated SEQ ID NO:Y are sufficiently accurate and otherwise suitable for a variety of uses well known in the art and described further below. For instance, SEQ ID NO:X is useful for designing nucleic acid hybridization probes that will detect nucleic acid sequences contained in SEQ ID NO:X or the cDNA contained in the deposited clone. These probes will also hybridize to nucleic acid molecules in biological samples, thereby enabling a variety of forensic and diagnostic methods of the invention. Similarly, polypeptides identified from SEQ ID NO:Y may be used to generate antibodies which bind specifically to the secreted proteins encoded by the cDNA clones identified in Table 1.

Nevertheless, DNA sequences generated by sequencing reactions can contain sequencing errors. The errors exist as misidentified nucleotides, or as insertions or deletions of nucleotides in the generated DNA sequence. The erroneously inserted or deleted nucleotides cause frame shifts in the reading frames of the predicted amino acid sequence. In these cases, the predicted amino acid sequence diverges from the actual amino acid sequence, even though the generated DNA sequence may be greater than 99.9% identical to the actual DNA sequence (for example, one base insertion or deletion in an open reading frame of over 1000 bases).

Accordingly, for those applications requiring precision in the nucleotide sequence or the amino acid sequence, the present invention provides not only the generated nucleotide sequence identified as SEQ ID NO:X and the predicted translated amino acid sequence identified as SEQ ID NO:Y, but also a sample of plasmid DNA containing a human cDNA of the invention deposited with the ATCC, as set forth in Table 1. The nucleotide sequence of each deposited clone can readily be determined by sequencing the deposited clone in accordance with known methods. The predicted amino acid sequence can then be verified from such deposits. Moreover, the amino acid sequence of the protein encoded by a particular clone can also be directly determined by peptide sequencing or by expressing the protein in a suitable host cell containing the deposited human cDNA, collecting the protein, and determining its sequence.

The present invention also relates to the genes corresponding to SEQ ID NO:X, SEQ ID NO:Y, or the deposited clone. The corresponding gene can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include preparing probes or primers from the disclosed sequence and identifying or amplifying the corresponding gene from appropriate sources of genomic material.

Also provided in the present invention are species homologs. Species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source for the desired homologue.

The polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

The polypeptides may be in the form of the secreted protein, including the mature form, or may be a part of a larger protein, such as a fusion protein (see below).

It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification, such as multiple histidine residues, or an additional sequence for stability during recombinant production.

- 5 The polypeptides of the present invention are preferably provided in an isolated form, and preferably are substantially purified. A recombinantly produced version of a polypeptide, including the secreted polypeptide, can be substantially purified by the one-step method described in Smith and Johnson, *Gene* 67:31-40 (1988). Polypeptides of the invention also can be purified from natural or recombinant sources using antibodies of the invention raised against the secreted protein in methods which are well known in the art.
- 10

Signal Sequences

- Methods for predicting whether a protein has a signal sequence, as well as the cleavage point for that sequence, are available. For instance, the method of McGeoch, *Virus Res.* 3:271-286 (1985), uses the information from a short N-terminal charged region and a subsequent uncharged region of the complete (uncleaved) protein. The method of von Heinje, *Nucleic Acids Res.* 14:4683-4690 (1986) uses the information from the residues surrounding the cleavage site, typically residues -13 to +2, where +1 indicates the amino terminus of the secreted protein. The accuracy of predicting the cleavage points of known mammalian secretory proteins for each of these methods is in the range of 75-80%. (von Heinje, *supra.*) However, the two methods do not always produce the same predicted cleavage point(s) for a given protein.
- 15
- 20

- In the present case, the deduced amino acid sequence of the secreted polypeptide was analyzed by a computer program called SignalP (Henrik Nielsen et al., *Protein Engineering* 10:1-6 (1997)), which predicts the cellular location of a protein based on the amino acid sequence. As part of this computational prediction of localization, the methods of McGeoch and von Heinje are incorporated. The analysis of the amino acid sequences of the secreted proteins described herein by this program provided the results shown in Table 1.
- 25
- 30

- As one of ordinary skill would appreciate, however, cleavage sites sometimes vary from organism to organism and cannot be predicted with absolute certainty. Accordingly, the present invention provides secreted polypeptides having a sequence shown in SEQ ID NO:Y which have an N-terminus beginning within 5 residues (i.e., + or - 5 residues) of the predicted cleavage point. Similarly, it is also recognized that in some cases, cleavage of the signal sequence from a secreted protein is not entirely
- 35

uniform, resulting in more than one secreted species. These polypeptides, and the polynucleotides encoding such polypeptides, are contemplated by the present invention.

Moreover, the signal sequence identified by the above analysis may not necessarily predict the naturally occurring signal sequence. For example, the naturally occurring signal sequence may be further upstream from the predicted signal sequence. However, it is likely that the predicted signal sequence will be capable of directing the secreted protein to the ER. These polypeptides, and the polynucleotides encoding such polypeptides, are contemplated by the present invention.

10 **Polynucleotide and Polypeptide Variants**

"Variant" refers to a polynucleotide or polypeptide differing from the polynucleotide or polypeptide of the present invention, but retaining essential properties thereof. Generally, variants are overall closely similar, and, in many regions, identical to the polynucleotide or polypeptide of the present invention.

By a polynucleotide having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence of the present invention, it is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the polypeptide. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. The query sequence may be an entire sequence shown in Table 1, the ORF (open reading frame), or any fragment specified as described herein.

As a practical matter, whether any particular nucleic acid molecule or polypeptide is at least 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleotide sequence of the present invention can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. (1990) 6:237-245). In a sequence alignment the query and subject sequences are both DNA sequences. An RNA sequence can be compared by converting U's to T's. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB alignment of DNA sequences to calculate percent identity are: Matrix=Unitary, k-tuple=4, Mismatch Penalty=1, Joining Penalty=30, Randomization

Group Length=0, Cutoff Score=1, Gap Penalty=5, Gap Size Penalty 0.05, Window Size=500 or the length of the subject nucleotide sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence because of 5' or 3' deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for 5' and 3' truncations of the subject sequence when calculating percent identity. For subject sequences truncated at the 5' or 3' ends, relative to the query sequence, the percent identity is corrected by calculating the number of bases of the query sequence that are 5' and 3' of the subject sequence, which are not matched/aligned, as a percent of the total bases of the query sequence. Whether a nucleotide is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This corrected score is what is used for the purposes of the present invention. Only bases outside the 5' and 3' bases of the subject sequence, as displayed by the FASTDB alignment, which are not matched/aligned with the query sequence, are calculated for the purposes of manually adjusting the percent identity score.

For example, a 90 base subject sequence is aligned to a 100 base query sequence to determine percent identity. The deletions occur at the 5' end of the subject sequence and therefore, the FASTDB alignment does not show a matched/alignment of the first 10 bases at 5' end. The 10 unpaired bases represent 10% of the sequence (number of bases at the 5' and 3' ends not matched/total number of bases in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 bases were perfectly matched the final percent identity would be 90%. In another example, a 90 base subject sequence is compared with a 100 base query sequence. This time the deletions are internal deletions so that there are no bases on the 5' or 3' of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only bases 5' and 3' of the subject sequence which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a query amino acid sequence of the present invention, it is intended that the amino acid sequence of the subject polypeptide is identical to the query sequence except that the subject polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the query amino acid sequence. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a query

amino acid sequence, up to 5% of the amino acid residues in the subject sequence may be inserted, deleted, (indels) or substituted with another amino acid. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions,
5 interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular polypeptide is at least 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequences shown in Table 1 or to the amino acid sequence encoded by deposited DNA clone can be
10 determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. (1990) 6:237-245). In a sequence alignment the query and
15 subject sequences are either both nucleotide sequences or both amino acid sequences. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.05, Window
20 Size=500 or the length of the subject amino acid sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity.
25 For subject sequences truncated at the N- and C-termini, relative to the the query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are N- and C-terminal of the subject sequence, which are not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. Whether a residue is matched/aligned is determined by results of
30 the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of the present invention. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the query sequence, are
35 considered for the purposes of manually adjusting the percent identity score. That is, only query residue positions outside the farthest N- and C-terminal residues of the subject sequence.

For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and C- termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100 residue query sequence. This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

The variants may contain alterations in the coding regions, non-coding regions, or both. Especially preferred are polynucleotide variants containing alterations which produce silent substitutions, additions, or deletions, but do not alter the properties or activities of the encoded polypeptide. Nucleotide variants produced by silent substitutions due to the degeneracy of the genetic code are preferred. Moreover, variants in which 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination are also preferred. Polynucleotide variants can be produced for a variety of reasons, e.g., to optimize codon expression for a particular host (change codons in the human mRNA to those preferred by a bacterial host such as *E. coli*).

Naturally occurring variants are called "allelic variants," and refer to one of several alternate forms of a gene occupying a given locus on a chromosome of an organism. (Genes II, Lewin, B., ed., John Wiley & Sons, New York (1985).) These allelic variants can vary at either the polynucleotide and/or polypeptide level. Alternatively, non-naturally occurring variants may be produced by mutagenesis techniques or by direct synthesis.

Using known methods of protein engineering and recombinant DNA technology, variants may be generated to improve or alter the characteristics of the polypeptides of the present invention. For instance, one or more amino acids can be deleted from the N-terminus or C-terminus of the secreted protein without substantial loss of biological function. The authors of Ron et al., *J. Biol. Chem.* 268: 2984-2988 (1993), reported variant KGF proteins having heparin binding activity even after

deleting 3, 8, or 27 amino-terminal amino acid residues. Similarly, Interferon gamma exhibited up to ten times higher activity after deleting 8-10 amino acid residues from the carboxy terminus of this protein. (Dobeli et al., J. Biotechnology 7:199-216 (1988).)

Moreover, ample evidence demonstrates that variants often retain a biological activity similar to that of the naturally occurring protein. For example, Gayle and coworkers (J. Biol. Chem 268:22105-22111 (1993)) conducted extensive mutational analysis of human cytokine IL-1a. They used random mutagenesis to generate over 3,500 individual IL-1a mutants that averaged 2.5 amino acid changes per variant over the entire length of the molecule. Multiple mutations were examined at every possible amino acid position. The investigators found that "[m]ost of the molecule could be altered with little effect on either [binding or biological activity]." (See, Abstract.) In fact, only 23 unique amino acid sequences, out of more than 3,500 nucleotide sequences examined, produced a protein that significantly differed in activity from wild-type.

Furthermore, even if deleting one or more amino acids from the N-terminus or C-terminus of a polypeptide results in modification or loss of one or more biological functions, other biological activities may still be retained. For example, the ability of a deletion variant to induce and/or to bind antibodies which recognize the secreted form will likely be retained when less than the majority of the residues of the secreted form are removed from the N-terminus or C-terminus. Whether a particular polypeptide lacking N- or C-terminal residues of a protein retains such immunogenic activities can readily be determined by routine methods described herein and otherwise known in the art.

Thus, the invention further includes polypeptide variants which show substantial biological activity. Such variants include deletions, insertions, inversions, repeats, and substitutions selected according to general rules known in the art so as to have little effect on activity. For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie, J. U. et al., Science 247:1306-1310 (1990), wherein the authors indicate that there are two main strategies for studying the tolerance of an amino acid sequence to change.

The first strategy exploits the tolerance of amino acid substitutions by natural selection during the process of evolution. By comparing amino acid sequences in different species, conserved amino acids can be identified. These conserved amino acids are likely important for protein function. In contrast, the amino acid positions where substitutions have been tolerated by natural selection indicates that these positions are not critical for protein function. Thus, positions tolerating amino acid substitution could be modified while still maintaining biological activity of the protein.

The second strategy uses genetic engineering to introduce amino acid changes at specific positions of a cloned gene to identify regions critical for protein function. For example, site directed mutagenesis or alanine-scanning mutagenesis (introduction of single alanine mutations at every residue in the molecule) can be used. (Cunningham and Wells, Science 244:1081-1085 (1989).) The resulting mutant molecules can then be tested for biological activity.

As the authors state, these two strategies have revealed that proteins are surprisingly tolerant of amino acid substitutions. The authors further indicate which amino acid changes are likely to be permissive at certain amino acid positions in the protein. For example, most buried (within the tertiary structure of the protein) amino acid residues require nonpolar side chains, whereas few features of surface side chains are generally conserved. Moreover, tolerated conservative amino acid substitutions involve replacement of the aliphatic or hydrophobic amino acids Ala, Val, Leu and Ile; replacement of the hydroxyl residues Ser and Thr; replacement of the acidic residues Asp and Glu; replacement of the amide residues Asn and Gln, replacement of the basic residues Lys, Arg, and His; replacement of the aromatic residues Phe, Tyr, and Trp, and replacement of the small-sized amino acids Ala, Ser, Thr, Met, and Gly.

Besides conservative amino acid substitution, variants of the present invention include (i) substitutions with one or more of the non-conserved amino acid residues, where the substituted amino acid residues may or may not be one encoded by the genetic code, or (ii) substitution with one or more of amino acid residues having a substituent group, or (iii) fusion of the mature polypeptide with another compound, such as a compound to increase the stability and/or solubility of the polypeptide (for example, polyethylene glycol), or (iv) fusion of the polypeptide with additional amino acids, such as an IgG Fc fusion region peptide, or leader or secretory sequence, or a sequence facilitating purification. Such variant polypeptides are deemed to be within the scope of those skilled in the art from the teachings herein.

For example, polypeptide variants containing amino acid substitutions of charged amino acids with other charged or neutral amino acids may produce proteins with improved characteristics, such as less aggregation. Aggregation of pharmaceutical formulations both reduces activity and increases clearance due to the aggregate's immunogenic activity. (Pinckard et al., Clin. Exp. Immunol. 2:331-340 (1967); Robbins et al., Diabetes 36: 838-845 (1987); Cleland et al., Crit. Rev. Therapeutic Drug Carrier Systems 10:307-377 (1993).)

Polynucleotide and Polypeptide Fragments

In the present invention, a "polynucleotide fragment" refers to a short polynucleotide having a nucleic acid sequence contained in the deposited clone or shown in SEQ ID NO:X. The short nucleotide fragments are preferably at least about 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably, at least about 40 nt in length. A fragment "at least 20 nt in length," for example, is intended to include 20 or more contiguous bases from the cDNA sequence contained in the deposited clone or the nucleotide sequence shown in SEQ ID NO:X. These nucleotide fragments are useful as diagnostic probes and primers as discussed herein. Of course, larger fragments (e.g., 50, 150, 500, 600, 2000 nucleotides) are preferred.

Moreover, representative examples of polynucleotide fragments of the invention, include, for example, fragments having a sequence from about nucleotide number 1-50, 51-100, 101-150, 151-200, 201-250, 251-300, 301-350, 351-400, 401-450, 451-500, 501-550, 551-600, 651-700, 701-750, 751-800, 800-850, 851-900, 901-950, 951-1000, 1001-1050, 1051-1100, 1101-1150, 1151-1200, 1201-1250, 1251-1300, 1301-1350, 1351-1400, 1401-1450, 1451-1500, 1501-1550, 1551-1600, 1601-1650, 1651-1700, 1701-1750, 1751-1800, 1801-1850, 1851-1900, 1901-1950, 1951-2000, or 2001 to the end of SEQ ID NO:X or the cDNA contained in the deposited clone. In this context "about" includes the particularly recited ranges, larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. Preferably, these fragments encode a polypeptide which has biological activity. More preferably, these polynucleotides can be used as probes or primers as discussed herein.

In the present invention, a "polypeptide fragment" refers to a short amino acid sequence contained in SEQ ID NO:Y or encoded by the cDNA contained in the deposited clone. Protein fragments may be "free-standing," or comprised within a larger polypeptide of which the fragment forms a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, 102-120, 121-140, 141-160, or 161 to the end of the coding region. Moreover, polypeptide fragments can be about 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, or 150 amino acids in length. In this context "about" includes the particularly recited ranges, larger or smaller by several (5, 4, 3, 2, or 1) amino acids, at either extreme or at both extremes.

Preferred polypeptide fragments include the secreted protein as well as the mature form. Further preferred polypeptide fragments include the secreted protein or the mature form having a continuous series of deleted residues from the amino or the

carboxy terminus, or both. For example, any number of amino acids, ranging from 1-60, can be deleted from the amino terminus of either the secreted polypeptide or the mature form. Similarly, any number of amino acids, ranging from 1-30, can be deleted from the carboxy terminus of the secreted protein or mature form. Furthermore, any combination of the above amino and carboxy terminus deletions are preferred. Similarly, polynucleotide fragments encoding these polypeptide fragments are also preferred.

Particularly, N-terminal deletions of the polypeptide of the present invention can be described by the general formula m-p, where p is the total number of amino acids in the polypeptide and m is an integer from 2 to (p-1), and where both of these integers (m & p) correspond to the position of the amino acid residue identified in SEQ ID NO:Y.

Moreover, C-terminal deletions of the polypeptide of the present invention can also be described by the general formula 1-n, where n is an integer from 2 to (p-1), and again where these integers (n & p) correspond to the position of the amino acid residue identified in SEQ ID NO:Y.

The invention also provides polypeptides having one or more amino acids deleted from both the amino and the carboxyl termini, which may be described generally as having residues m-n of SEQ ID NO:Y, where m and n are integers as described above.

Also preferred are polypeptide and polynucleotide fragments characterized by structural or functional domains, such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions. Polypeptide fragments of SEQ ID NO:Y falling within conserved domains are specifically contemplated by the present invention. Moreover, polynucleotide fragments encoding these domains are also contemplated.

Other preferred fragments are biologically active fragments. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the polypeptide of the present invention. The biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity.

Epitopes & Antibodies

In the present invention, "epitopes" refer to polypeptide fragments having antigenic or immunogenic activity in an animal, especially in a human. A preferred embodiment of the present invention relates to a polypeptide fragment comprising an

epitope, as well as the polynucleotide encoding this fragment. A region of a protein molecule to which an antibody can bind is defined as an "antigenic epitope." In contrast, an "immunogenic epitope" is defined as a part of a protein that elicits an antibody response. (See, for instance, Geysen et al., Proc. Natl. Acad. Sci. USA 81:3998-4002 (1983).)

Fragments which function as epitopes may be produced by any conventional means. (See, e.g., Houghten, R. A., Proc. Natl. Acad. Sci. USA 82:5131-5135 (1985) further described in U.S. Patent No. 4,631,211.)

In the present invention, antigenic epitopes preferably contain a sequence of at least seven, more preferably at least nine, and most preferably between about 15 to about 30 amino acids. Antigenic epitopes are useful to raise antibodies, including monoclonal antibodies, that specifically bind the epitope. (See, for instance, Wilson et al., Cell 37:767-778 (1984); Sutcliffe, J. G. et al., Science 219:660-666 (1983).)

Similarly, immunogenic epitopes can be used to induce antibodies according to methods well known in the art. (See, for instance, Sutcliffe et al., supra; Wilson et al., supra; Chow, M. et al., Proc. Natl. Acad. Sci. USA 82:910-914; and Bittle, F. J. et al., J. Gen. Virol. 66:2347-2354 (1985).) A preferred immunogenic epitope includes the secreted protein. The immunogenic epitopes may be presented together with a carrier protein, such as an albumin, to an animal system (such as rabbit or mouse) or, if it is long enough (at least about 25 amino acids), without a carrier. However, immunogenic epitopes comprising as few as 8 to 10 amino acids have been shown to be sufficient to raise antibodies capable of binding to, at the very least, linear epitopes in a denatured polypeptide (e.g., in Western blotting.)

As used herein, the term "antibody" (Ab) or "monoclonal antibody" (Mab) is meant to include intact molecules as well as antibody fragments (such as, for example, Fab and F(ab')₂ fragments) which are capable of specifically binding to protein. Fab and F(ab')₂ fragments lack the Fc fragment of intact antibody, clear more rapidly from the circulation, and may have less non-specific tissue binding than an intact antibody. (Wahl et al., J. Nucl. Med. 24:316-325 (1983).) Thus, these fragments are preferred, as well as the products of a FAB or other immunoglobulin expression library. Moreover, antibodies of the present invention include chimeric, single chain, and humanized antibodies.

Fusion Proteins

Any polypeptide of the present invention can be used to generate fusion proteins. For example, the polypeptide of the present invention, when fused to a second protein, can be used as an antigenic tag. Antibodies raised against the

polypeptide of the present invention can be used to indirectly detect the second protein by binding to the polypeptide. Moreover, because secreted proteins target cellular locations based on trafficking signals, the polypeptides of the present invention can be used as targeting molecules once fused to other proteins.

5 Examples of domains that can be fused to polypeptides of the present invention include not only heterologous signal sequences, but also other heterologous functional regions. The fusion does not necessarily need to be direct, but may occur through linker sequences.

10 Moreover, fusion proteins may also be engineered to improve characteristics of the polypeptide of the present invention. For instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence during purification from the host cell or subsequent handling and storage. Also, peptide moieties may be added to the polypeptide to facilitate purification. Such regions may be removed prior to final
15 preparation of the polypeptide. The addition of peptide moieties to facilitate handling of polypeptides are familiar and routine techniques in the art.

 Moreover, polypeptides of the present invention, including fragments, and specifically epitopes, can be combined with parts of the constant domain of immunoglobulins (IgG), resulting in chimeric polypeptides. These fusion proteins
20 facilitate purification and show an increased half-life in vivo. One reported example describes chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. (EP A 394,827; Traunecker et al., Nature 331:84-86 (1988).) Fusion proteins having disulfide-linked dimeric structures (due to the IgG)
25 can also be more efficient in binding and neutralizing other molecules, than the monomeric secreted protein or protein fragment alone. (Fountoulakis et al., J. Biochem. 270:3958-3964 (1995).)

 Similarly, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules
30 together with another human protein or part thereof. In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and thus can result in, for example, improved pharmacokinetic properties. (EP-A 0232 262.) Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified, would be desired. For example, the Fc portion may hinder therapy and diagnosis if the
35 fusion protein is used as an antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. (See, D.

Bennett et al., J. Molecular Recognition 8:52-58 (1995); K. Johanson et al., J. Biol. Chem. 270:9459-9471 (1995).)

- Moreover, the polypeptides of the present invention can be fused to marker sequences, such as a peptide which facilitates purification of the fused polypeptide. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., Proc. Natl. Acad. Sci. USA 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein.
- Another peptide tag useful for purification, the "HA" tag, corresponds to an epitope derived from the influenza hemagglutinin protein. (Wilson et al., Cell 37:767 (1984).)

Thus, any of these above fusions can be engineered using the polynucleotides or the polypeptides of the present invention.

15 **Vectors, Host Cells, and Protein Production**

- The present invention also relates to vectors containing the polynucleotide of the present invention, host cells, and the production of polypeptides by recombinant techniques. The vector may be, for example, a phage, plasmid, viral, or retroviral vector. Retroviral vectors may be replication competent or replication defective. In the latter case, viral propagation generally will occur only in complementing host cells.

- The polynucleotides may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the vector is a virus, it may be packaged in vitro using an appropriate packaging cell line and then transduced into host cells.

- The polynucleotide insert should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the E. coli lac, trp, phoA and tac promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan. The expression constructs will further contain sites for transcription initiation, termination, and, in the transcribed region, a ribosome binding site for translation. The coding portion of the transcripts expressed by the constructs will preferably include a translation initiating codon at the beginning and a termination codon (UAA, UGA or UAG) appropriately positioned at the end of the polypeptide to be translated.

- As indicated, the expression vectors will preferably include at least one selectable marker. Such markers include dihydrofolate reductase, G418 or neomycin resistance for eukaryotic cell culture and tetracycline, kanamycin or ampicillin resistance

genes for culturing in *E. coli* and other bacteria. Representative examples of appropriate hosts include, but are not limited to, bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells; insect cells such as *Drosophila* S2 and *Spodoptera Sf9* cells; animal cells such as CHO, COS, 293, and Bowes melanoma cells; and plant cells. Appropriate culture mediums and conditions for the above-described host cells are known in the art.

Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from QIAGEN, Inc.; pBluescript vectors, Phagescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene Cloning Systems, Inc.; and ptc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia Biotech, Inc. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Other suitable vectors will be readily apparent to the skilled artisan.

Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, or other methods. Such methods are described in many standard laboratory manuals, such as Davis et al., *Basic Methods In Molecular Biology* (1986). It is specifically contemplated that the polypeptides of the present invention may in fact be expressed by a host cell lacking a recombinant vector.

A polypeptide of this invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification.

Polypeptides of the present invention, and preferably the secreted form, can also be recovered from: products purified from natural sources, including bodily fluids, tissues and cells, whether directly isolated or cultured; products of chemical synthetic procedures; and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect, and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-mediated processes. Thus, it is well known in the art that the N-terminal methionine encoded by the translation initiation codon generally is removed with high efficiency from any protein

after translation in all eukaryotic cells. While the N-terminal methionine on most proteins also is efficiently removed in most prokaryotes, for some proteins, this prokaryotic removal process is inefficient, depending on the nature of the amino acid to which the N-terminal methionine is covalently linked.

5

Uses of the Polynucleotides

Each of the polynucleotides identified herein can be used in numerous ways as reagents. The following description should be considered exemplary and utilizes known techniques.

10

The polynucleotides of the present invention are useful for chromosome identification. There exists an ongoing need to identify new chromosome markers, since few chromosome marking reagents, based on actual sequence data (repeat polymorphisms), are presently available. Each polynucleotide of the present invention can be used as a chromosome marker.

15

Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the sequences shown in SEQ ID NO:X. Primers can be selected using computer analysis so that primers do not span more than one predicted exon in the genomic DNA. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the SEQ ID NO:X will yield an amplified fragment.

20

Similarly, somatic hybrids provide a rapid method of PCR mapping the polynucleotides to particular chromosomes. Three or more clones can be assigned per day using a single thermal cycler. Moreover, sublocalization of the polynucleotides can be achieved with panels of specific chromosome fragments. Other gene mapping strategies that can be used include in situ hybridization, prescreening with labeled flow-sorted chromosomes, and preselection by hybridization to construct chromosome specific-cDNA libraries.

25

Precise chromosomal location of the polynucleotides can also be achieved using fluorescence in situ hybridization (FISH) of a metaphase chromosomal spread. This technique uses polynucleotides as short as 500 or 600 bases; however, polynucleotides 2,000-4,000 bp are preferred. For a review of this technique, see Verma et al., "Human Chromosomes: a Manual of Basic Techniques," Pergamon Press, New York (1988).

30

For chromosome mapping, the polynucleotides can be used individually (to mark a single chromosome or a single site on that chromosome) or in panels (for marking multiple sites and/or multiple chromosomes). Preferred polynucleotides correspond to the noncoding regions of the cDNAs because the coding sequences are

35

more likely conserved within gene families, thus increasing the chance of cross hybridization during chromosomal mapping.

Once a polynucleotide has been mapped to a precise chromosomal location, the physical position of the polynucleotide can be used in linkage analysis. Linkage analysis establishes coinheritance between a chromosomal location and presentation of a particular disease. (Disease mapping data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library) .) Assuming 1 megabase mapping resolution and one gene per 20 kb, a cDNA precisely localized to a chromosomal region associated with the disease could be one of 50-500 potential causative genes.

Thus, once coinheritance is established, differences in the polynucleotide and the corresponding gene between affected and unaffected individuals can be examined. First, visible structural alterations in the chromosomes, such as deletions or translocations, are examined in chromosome spreads or by PCR. If no structural alterations exist, the presence of point mutations are ascertained. Mutations observed in some or all affected individuals, but not in normal individuals, indicates that the mutation may cause the disease. However, complete sequencing of the polypeptide and the corresponding gene from several normal individuals is required to distinguish the mutation from a polymorphism. If a new polymorphism is identified, this polymorphic polypeptide can be used for further linkage analysis.

Furthermore, increased or decreased expression of the gene in affected individuals as compared to unaffected individuals can be assessed using polynucleotides of the present invention. Any of these alterations (altered expression, chromosomal rearrangement, or mutation) can be used as a diagnostic or prognostic marker.

In addition to the foregoing, a polynucleotide can be used to control gene expression through triple helix formation or antisense DNA or RNA. Both methods rely on binding of the polynucleotide to DNA or RNA. For these techniques, preferred polynucleotides are usually 20 to 40 bases in length and complementary to either the region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res. 6:3073 (1979); Cooney et al., Science 241:456 (1988); and Dervan et al., Science 251:1360 (1991)) or to the mRNA itself (antisense - Okano, J. Neurochem. 56:560 (1991); Oligodeoxy-nucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988).) Triple helix formation optimally results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. Both techniques are effective in model

systems, and the information disclosed herein can be used to design antisense or triple helix polynucleotides in an effort to treat disease.

Polynucleotides of the present invention are also useful in gene therapy. One goal of gene therapy is to insert a normal gene into an organism having a defective gene, in an effort to correct the genetic defect. The polynucleotides disclosed in the present invention offer a means of targeting such genetic defects in a highly accurate manner. Another goal is to insert a new gene that was not present in the host genome, thereby producing a new trait in the host cell.

The polynucleotides are also useful for identifying individuals from minute biological samples. The United States military, for example, is considering the use of restriction fragment length polymorphism (RFLP) for identification of its personnel. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identifying personnel. This method does not suffer from the current limitations of "Dog Tags" which can be lost, switched, or stolen, making positive identification difficult. The polynucleotides of the present invention can be used as additional DNA markers for RFLP.

The polynucleotides of the present invention can also be used as an alternative to RFLP, by determining the actual base-by-base DNA sequence of selected portions of an individual's genome. These sequences can be used to prepare PCR primers for amplifying and isolating such selected DNA, which can then be sequenced. Using this technique, individuals can be identified because each individual will have a unique set of DNA sequences. Once an unique ID database is established for an individual, positive identification of that individual, living or dead, can be made from extremely small tissue samples.

Forensic biology also benefits from using DNA-based identification techniques as disclosed herein. DNA sequences taken from very small biological samples such as tissues, e.g., hair or skin, or body fluids, e.g., blood, saliva, semen, etc., can be amplified using PCR. In one prior art technique, gene sequences amplified from polymorphic loci, such as DQa class II HLA gene, are used in forensic biology to identify individuals. (Erlich, H., PCR Technology, Freeman and Co. (1992).) Once these specific polymorphic loci are amplified, they are digested with one or more restriction enzymes, yielding an identifying set of bands on a Southern blot probed with DNA corresponding to the DQa class II HLA gene. Similarly, polynucleotides of the present invention can be used as polymorphic markers for forensic purposes.

There is also a need for reagents capable of identifying the source of a particular tissue. Such need arises, for example, in forensics when presented with tissue of

unknown origin. Appropriate reagents can comprise, for example, DNA probes or primers specific to particular tissue prepared from the sequences of the present invention. Panels of such reagents can identify tissue by species and/or by organ type. In a similar fashion, these reagents can be used to screen tissue cultures for contamination.

In the very least, the polynucleotides of the present invention can be used as molecular weight markers on Southern gels, as diagnostic probes for the presence of a specific mRNA in a particular cell type, as a probe to "subtract-out" known sequences in the process of discovering novel polynucleotides, for selecting and making oligomers for attachment to a "gene chip" or other support, to raise anti-DNA antibodies using DNA immunization techniques, and as an antigen to elicit an immune response.

Uses of the Polypeptides

Each of the polypeptides identified herein can be used in numerous ways. The following description should be considered exemplary and utilizes known techniques.

A polypeptide of the present invention can be used to assay protein levels in a biological sample using antibody-based techniques. For example, protein expression in tissues can be studied with classical immunohistological methods. (Jalkanen, M., et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, M., et al., J. Cell Biol. 105:3087-3096 (1987).) Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase, and radioisotopes, such as iodine (125I, 121I), carbon (14C), sulfur (35S), tritium (3H), indium (112In), and technetium (99mTc), and fluorescent labels, such as fluorescein and rhodamine, and biotin.

In addition to assaying secreted protein levels in a biological sample, proteins can also be detected in vivo by imaging. Antibody labels or markers for in vivo imaging of protein include those detectable by X-radiography, NMR or ESR. For X-radiography, suitable labels include radioisotopes such as barium or cesium, which emit detectable radiation but are not overtly harmful to the subject. Suitable markers for NMR and ESR include those with a detectable characteristic spin, such as deuterium, which may be incorporated into the antibody by labeling of nutrients for the relevant hybridoma.

A protein-specific antibody or antibody fragment which has been labeled with an appropriate detectable imaging moiety, such as a radioisotope (for example, 131I, 112In, 99mTc), a radio-opaque substance, or a material detectable by nuclear magnetic

resonance, is introduced (for example, parenterally, subcutaneously, or intraperitoneally) into the mammal. It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a human
5 subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of ^{99m}Tc. The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which contain the specific protein. In vivo tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments." (Chapter 13 in Tumor Imaging: The
10 Radiochemical Detection of Cancer, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982).)

Thus, the invention provides a diagnostic method of a disorder, which involves (a) assaying the expression of a polypeptide of the present invention in cells or body fluid of an individual; (b) comparing the level of gene expression with a standard gene
15 expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of a disorder.

Moreover, polypeptides of the present invention can be used to treat disease. For example, patients can be administered a polypeptide of the present invention in an effort to replace absent or decreased levels of the polypeptide (e.g., insulin), to
20 supplement absent or decreased levels of a different polypeptide (e.g., hemoglobin S for hemoglobin B), to inhibit the activity of a polypeptide (e.g., an oncogene), to activate the activity of a polypeptide (e.g., by binding to a receptor), to reduce the activity of a membrane bound receptor by competing with it for free ligand (e.g., soluble TNF receptors used in reducing inflammation), or to bring about a desired
25 response (e.g., blood vessel growth).

Similarly, antibodies directed to a polypeptide of the present invention can also be used to treat disease. For example, administration of an antibody directed to a polypeptide of the present invention can bind and reduce overproduction of the polypeptide. Similarly, administration of an antibody can activate the polypeptide, such
30 as by binding to a polypeptide bound to a membrane (receptor).

At the very least, the polypeptides of the present invention can be used as molecular weight markers on SDS-PAGE gels or on molecular sieve gel filtration columns using methods well known to those of skill in the art. Polypeptides can also be used to raise antibodies, which in turn are used to measure protein expression from a
35 recombinant cell, as a way of assessing transformation of the host cell. Moreover, the polypeptides of the present invention can be used to test the following biological activities.

Biological Activities

The polynucleotides and polypeptides of the present invention can be used in assays to test for one or more biological activities. If these polynucleotides and polypeptides do exhibit activity in a particular assay, it is likely that these molecules may be involved in the diseases associated with the biological activity. Thus, the polynucleotides and polypeptides could be used to treat the associated disease.

Immune Activity

A polypeptide or polynucleotide of the present invention may be useful in treating deficiencies or disorders of the immune system, by activating or inhibiting the proliferation, differentiation, or mobilization (chemotaxis) of immune cells. Immune cells develop through a process called hematopoiesis, producing myeloid (platelets, red blood cells, neutrophils, and macrophages) and lymphoid (B and T lymphocytes) cells from pluripotent stem cells. The etiology of these immune deficiencies or disorders may be genetic, somatic, such as cancer or some autoimmune disorders, acquired (e.g., by chemotherapy or toxins), or infectious. Moreover, a polynucleotide or polypeptide of the present invention can be used as a marker or detector of a particular immune system disease or disorder.

A polynucleotide or polypeptide of the present invention may be useful in treating or detecting deficiencies or disorders of hematopoietic cells. A polypeptide or polynucleotide of the present invention could be used to increase differentiation and proliferation of hematopoietic cells, including the pluripotent stem cells, in an effort to treat those disorders associated with a decrease in certain (or many) types hematopoietic cells. Examples of immunologic deficiency syndromes include, but are not limited to: blood protein disorders (e.g. agammaglobulinemia, dysgammaglobulinemia), ataxia telangiectasia, common variable immunodeficiency, Digeorge Syndrome, HIV infection, HTLV-BLV infection, leukocyte adhesion deficiency syndrome, lymphopenia, phagocyte bactericidal dysfunction, severe combined immunodeficiency (SCIDs), Wiskott-Aldrich Disorder, anemia, thrombocytopenia, or hemoglobinuria.

Moreover, a polypeptide or polynucleotide of the present invention could also be used to modulate hemostatic (the stopping of bleeding) or thrombolytic activity (clot formation). For example, by increasing hemostatic or thrombolytic activity, a polynucleotide or polypeptide of the present invention could be used to treat blood coagulation disorders (e.g., afibrinogenemia, factor deficiencies), blood platelet disorders (e.g. thrombocytopenia), or wounds resulting from trauma, surgery, or other causes. Alternatively, a polynucleotide or polypeptide of the present invention that can

decrease hemostatic or thrombolytic activity could be used to inhibit or dissolve clotting. These molecules could be important in the treatment of heart attacks (infarction), strokes, or scarring.

5 A polynucleotide or polypeptide of the present invention may also be useful in treating or detecting autoimmune disorders. Many autoimmune disorders result from inappropriate recognition of self as foreign material by immune cells. This inappropriate recognition results in an immune response leading to the destruction of the host tissue. Therefore, the administration of a polypeptide or polynucleotide of the present invention that inhibits an immune response, particularly the proliferation,
10 differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing autoimmune disorders.

Examples of autoimmune disorders that can be treated or detected by the present invention include, but are not limited to: Addison's Disease, hemolytic anemia, antiphospholipid syndrome, rheumatoid arthritis, dermatitis, allergic encephalomyelitis,
15 glomerulonephritis, Goodpasture's Syndrome, Graves' Disease, Multiple Sclerosis, Myasthenia Gravis, Neuritis, Ophthalmia, Bullous Pemphigoid, Pemphigus, Polyendocrinopathies, Purpura, Reiter's Disease, Stiff-Man Syndrome, Autoimmune Thyroiditis, Systemic Lupus Erythematosus, Autoimmune Pulmonary Inflammation, Guillain-Barre Syndrome, insulin dependent diabetes mellitus, and autoimmune
20 inflammatory eye disease.

Similarly, allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems, may also be treated by a polypeptide or polynucleotide of the present invention. Moreover, these molecules can be used to treat
anaphylaxis, hypersensitivity to an antigenic molecule, or blood group incompatibility.

25 A polynucleotide or polypeptide of the present invention may also be used to treat and/or prevent organ rejection or graft-versus-host disease (GVHD). Organ rejection occurs by host immune cell destruction of the transplanted tissue through an immune response. Similarly, an immune response is also involved in GVHD, but, in this case, the foreign transplanted immune cells destroy the host tissues. The
30 administration of a polypeptide or polynucleotide of the present invention that inhibits an immune response, particularly the proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing organ rejection or GVHD.

Similarly, a polypeptide or polynucleotide of the present invention may also be used to modulate inflammation. For example, the polypeptide or polynucleotide may
35 inhibit the proliferation and differentiation of cells involved in an inflammatory response. These molecules can be used to treat inflammatory conditions, both chronic and acute conditions, including inflammation associated with infection (e.g., septic

shock, sepsis, or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine induced lung injury, inflammatory bowel disease, Crohn's disease, or resulting from over production of cytokines (e.g., TNF or IL-1.)

Hyperproliferative Disorders

A polypeptide or polynucleotide can be used to treat or detect hyperproliferative disorders, including neoplasms. A polypeptide or polynucleotide of the present invention may inhibit the proliferation of the disorder through direct or indirect interactions. Alternatively, a polypeptide or polynucleotide of the present invention may proliferate other cells which can inhibit the hyperproliferative disorder.

For example, by increasing an immune response, particularly increasing antigenic qualities of the hyperproliferative disorder or by proliferating, differentiating, or mobilizing T-cells, hyperproliferative disorders can be treated. This immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, decreasing an immune response may also be a method of treating hyperproliferative disorders, such as a chemotherapeutic agent.

Examples of hyperproliferative disorders that can be treated or detected by a polynucleotide or polypeptide of the present invention include, but are not limited to neoplasms located in the: abdomen, bone, breast, digestive system, liver, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), eye, head and neck, nervous (central and peripheral), lymphatic system, pelvic, skin, soft tissue, spleen, thoracic, and urogenital.

Similarly, other hyperproliferative disorders can also be treated or detected by a polynucleotide or polypeptide of the present invention. Examples of such hyperproliferative disorders include, but are not limited to: hypergammaglobulinemia, lymphoproliferative disorders, paraproteinemias, purpura, sarcoidosis, Sezary Syndrome, Waldenstrom's Macroglobulinemia, Gaucher's Disease, histiocytosis, and any other hyperproliferative disease, besides neoplasia, located in an organ system listed above.

Infectious Disease

A polypeptide or polynucleotide of the present invention can be used to treat or detect infectious agents. For example, by increasing the immune response, particularly increasing the proliferation and differentiation of B and/or T cells, infectious diseases

may be treated. The immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, the polypeptide or polynucleotide of the present invention may also directly inhibit the infectious agent, without necessarily eliciting an immune response.

- 5 Viruses are one example of an infectious agent that can cause disease or symptoms that can be treated or detected by a polynucleotide or polypeptide of the present invention. Examples of viruses, include, but are not limited to the following DNA and RNA viral families: Arbovirus, Adenoviridae, Arenaviridae, Arterivirus, Birnaviridae, Bunyaviridae, Caliciviridae, Circoviridae, Coronaviridae, Flaviviridae, 10 Hepadnaviridae (Hepatitis), Herpesviridae (such as, Cytomegalovirus, Herpes Simplex, Herpes Zoster), Mononegavirus (e.g., Paramyxoviridae, Morbillivirus, Rhabdoviridae), Orthomyxoviridae (e.g., Influenza), Papovaviridae, Parvoviridae, Picornaviridae, Poxviridae (such as Smallpox or Vaccinia), Reoviridae (e.g., Rotavirus), Retroviridae (HTLV-I, HTLV-II, Lentivirus), and Togaviridae (e.g., 15 Rubivirus). Viruses falling within these families can cause a variety of diseases or symptoms, including, but not limited to: arthritis, bronchiolitis, encephalitis, eye infections (e.g., conjunctivitis, keratitis), chronic fatigue syndrome, hepatitis (A, B, C, E, Chronic Active, Delta), meningitis, opportunistic infections (e.g., AIDS), pneumonia, Burkitt's Lymphoma, chickenpox, hemorrhagic fever, Measles, Mumps, 20 Parainfluenza, Rabies, the common cold, Polio, leukemia, Rubella, sexually transmitted diseases, skin diseases (e.g., Kaposi's, warts), and viremia. A polypeptide or polynucleotide of the present invention can be used to treat or detect any of these symptoms or diseases.

- Similarly, bacterial or fungal agents that can cause disease or symptoms and that 25 can be treated or detected by a polynucleotide or polypeptide of the present invention include, but not limited to, the following Gram-Negative and Gram-positive bacterial families and fungi: Actinomycetales (e.g., Corynebacterium, Mycobacterium, Norcardia), Aspergillosis, Bacillaceae (e.g., Anthrax, Clostridium), Bacteroidaceae, Blastomycosis, Bordetella, Borrelia, Brucellosis, Candidiasis, Campylobacter, 30 Coccidioidomycosis, Cryptococcosis, Dermatocycoses, Enterobacteriaceae (Klebsiella, Salmonella, Serratia, Yersinia), Erysipelothrix, Helicobacter, Legionellosis, Leptospirosis, Listeria, Mycoplasmatales, Neisseriaceae (e.g., Acinetobacter, Gonorrhea, Meningococcal), Pasteurellaceae Infections (e.g., Actinobacillus, Haemophilus, Pasteurella), Pseudomonas, Rickettsiaceae, Chlamydiaceae, Syphilis, 35 and Staphylococcal. These bacterial or fungal families can cause the following diseases or symptoms, including, but not limited to: bacteremia, endocarditis, eye infections (conjunctivitis, tuberculosis, uveitis), gingivitis, opportunistic infections (e.g., AIDS

- related infections), paronychia, prosthesis-related infections, Reiter's Disease, respiratory tract infections, such as Whooping Cough or Empyema, sepsis, Lyme Disease, Cat-Scratch Disease, Dysentery, Paratyphoid Fever, food poisoning, Typhoid, pneumonia, Gonorrhea, meningitis, Chlamydia, Syphilis, Diphtheria,
- 5 Leprosy, Paratuberculosis, Tuberculosis, Lupus, Botulism, gangrene, tetanus, impetigo, Rheumatic Fever, Scarlet Fever, sexually transmitted diseases, skin diseases (e.g., cellulitis, dermatocycoses), toxemia, urinary tract infections, wound infections. A polypeptide or polynucleotide of the present invention can be used to treat or detect any of these symptoms or diseases.
- 10 Moreover, parasitic agents causing disease or symptoms that can be treated or detected by a polynucleotide or polypeptide of the present invention include, but not limited to, the following families: Amebiasis, Babesiosis, Coccidiosis, Cryptosporidiosis, Dientamoebiasis, Dourine, Ectoparasitic, Giardiasis, Helminthiasis, Leishmaniasis, Theileriasis, Toxoplasmosis, Trypanosomiasis, and Trichomonas.
- 15 These parasites can cause a variety of diseases or symptoms, including, but not limited to: Scabies, Trombiculiasis, eye infections, intestinal disease (e.g., dysentery, giardiasis), liver disease, lung disease, opportunistic infections (e.g., AIDS related), Malaria, pregnancy complications, and toxoplasmosis. A polypeptide or polynucleotide of the present invention can be used to treat or detect any of these symptoms or
- 20 diseases.
- Preferably, treatment using a polypeptide or polynucleotide of the present invention could either be by administering an effective amount of a polypeptide to the patient, or by removing cells from the patient, supplying the cells with a polynucleotide of the present invention, and returning the engineered cells to the patient (ex vivo
- 25 therapy). Moreover, the polypeptide or polynucleotide of the present invention can be used as an antigen in a vaccine to raise an immune response against infectious disease.

Regeneration

- A polynucleotide or polypeptide of the present invention can be used to
- 30 differentiate, proliferate, and attract cells, leading to the regeneration of tissues. (See, Science 276:59-87 (1997).) The regeneration of tissues could be used to repair, replace, or protect tissue damaged by congenital defects, trauma (wounds, burns, incisions, or ulcers), age, disease (e.g. osteoporosis, osteoarthritis, periodontal disease, liver failure), surgery, including cosmetic plastic surgery, fibrosis, reperfusion
- 35 injury, or systemic cytokine damage.

Tissues that could be regenerated using the present invention include organs (e.g., pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal

or cardiac), vascular (including vascular endothelium), nervous, hematopoietic, and skeletal (bone, cartilage, tendon, and ligament) tissue. Preferably, regeneration occurs without or decreased scarring. Regeneration also may include angiogenesis.

Moreover, a polynucleotide or polypeptide of the present invention may increase
5 regeneration of tissues difficult to heal. For example, increased tendon/ligament regeneration would quicken recovery time after damage. A polynucleotide or polypeptide of the present invention could also be used prophylactically in an effort to avoid damage. Specific diseases that could be treated include of tendinitis, carpal tunnel syndrome, and other tendon or ligament defects. A further example of tissue
10 regeneration of non-healing wounds includes pressure ulcers, ulcers associated with vascular insufficiency, surgical, and traumatic wounds.

Similarly, nerve and brain tissue could also be regenerated by using a polynucleotide or polypeptide of the present invention to proliferate and differentiate nerve cells. Diseases that could be treated using this method include central and
15 peripheral nervous system diseases, neuropathies, or mechanical and traumatic disorders (e.g., spinal cord disorders, head trauma, cerebrovascular disease, and stroke). Specifically, diseases associated with peripheral nerve injuries, peripheral neuropathy (e.g., resulting from chemotherapy or other medical therapies), localized neuropathies, and central nervous system diseases (e.g., Alzheimer's disease,
20 Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome), could all be treated using the polynucleotide or polypeptide of the present invention.

Chemotaxis

25 A polynucleotide or polypeptide of the present invention may have chemotaxis activity. A chemotactic molecule attracts or mobilizes cells (e.g., monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells) to a particular site in the body, such as inflammation, infection, or site of hyperproliferation. The mobilized cells can then fight off and/or heal the particular
30 trauma or abnormality.

A polynucleotide or polypeptide of the present invention may increase chemotactic activity of particular cells. These chemotactic molecules can then be used to treat inflammation, infection, hyperproliferative disorders, or any immune system disorder by increasing the number of cells targeted to a particular location in the body.
35 For example, chemotactic molecules can be used to treat wounds and other trauma to tissues by attracting immune cells to the injured location. Chemotactic molecules of the present invention can also attract fibroblasts, which can be used to treat wounds.

It is also contemplated that a polynucleotide or polypeptide of the present invention may inhibit chemotactic activity. These molecules could also be used to treat disorders. Thus, a polynucleotide or polypeptide of the present invention could be used as an inhibitor of chemotaxis.

5

Binding Activity

A polypeptide of the present invention may be used to screen for molecules that bind to the polypeptide or for molecules to which the polypeptide binds. The binding of the polypeptide and the molecule may activate (agonist), increase, inhibit
10 (antagonist), or decrease activity of the polypeptide or the molecule bound. Examples of such molecules include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

Preferably, the molecule is closely related to the natural ligand of the polypeptide, e.g., a fragment of the ligand, or a natural substrate, a ligand, a structural
15 or functional mimetic. (See, Coligan et al., Current Protocols in Immunology 1(2):Chapter 5 (1991).) Similarly, the molecule can be closely related to the natural receptor to which the polypeptide binds, or at least, a fragment of the receptor capable of being bound by the polypeptide (e.g., active site). In either case, the molecule can be rationally designed using known techniques.

Preferably, the screening for these molecules involves producing appropriate
20 cells which express the polypeptide, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, *Drosophila*, or *E. coli*. Cells expressing the polypeptide (or cell membrane containing the expressed polypeptide) are then preferably contacted with a test compound potentially containing
25 the molecule to observe binding, stimulation, or inhibition of activity of either the polypeptide or the molecule.

The assay may simply test binding of a candidate compound to the polypeptide, wherein binding is detected by a label, or in an assay involving competition with a labeled competitor. Further, the assay may test whether the candidate compound results
30 in a signal generated by binding to the polypeptide.

Alternatively, the assay can be carried out using cell-free preparations, polypeptide/molecule affixed to a solid support, chemical libraries, or natural product mixtures. The assay may also simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide, measuring polypeptide/molecule
35 activity or binding, and comparing the polypeptide/molecule activity or binding to a standard.

Preferably, an ELISA assay can measure polypeptide level or activity in a sample (e.g., biological sample) using a monoclonal or polyclonal antibody. The antibody can measure polypeptide level or activity by either binding, directly or indirectly, to the polypeptide or by competing with the polypeptide for a substrate.

5 All of these above assays can be used as diagnostic or prognostic markers. The molecules discovered using these assays can be used to treat disease or to bring about a particular result in a patient (e.g., blood vessel growth) by activating or inhibiting the polypeptide/molecule. Moreover, the assays can discover agents which may inhibit or enhance the production of the polypeptide from suitably manipulated cells or tissues.

10 Therefore, the invention includes a method of identifying compounds which bind to a polypeptide of the invention comprising the steps of: (a) incubating a candidate binding compound with a polypeptide of the invention; and (b) determining if binding has occurred. Moreover, the invention includes a method of identifying agonists/antagonists comprising the steps of: (a) incubating a candidate compound with
15 a polypeptide of the invention, (b) assaying a biological activity, and (b) determining if a biological activity of the polypeptide has been altered.

Other Activities

20 A polypeptide or polynucleotide of the present invention may also increase or decrease the differentiation or proliferation of embryonic stem cells, besides, as discussed above, hematopoietic lineage.

A polypeptide or polynucleotide of the present invention may also be used to modulate mammalian characteristics, such as body height, weight, hair color, eye color, skin, percentage of adipose tissue, pigmentation, size, and shape (e.g., cosmetic
25 surgery). Similarly, a polypeptide or polynucleotide of the present invention may be used to modulate mammalian metabolism affecting catabolism, anabolism, processing, utilization, and storage of energy.

A polypeptide or polynucleotide of the present invention may be used to change a mammal's mental state or physical state by influencing biorhythms, circadian
30 rhythms, depression (including depressive disorders), tendency for violence, tolerance for pain, reproductive capabilities (preferably by Activin or Inhibin-like activity), hormonal or endocrine levels, appetite, libido, memory, stress, or other cognitive qualities.

35 A polypeptide or polynucleotide of the present invention may also be used as a food additive or preservative, such as to increase or decrease storage capabilities, fat content, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional components.

Other Preferred Embodiments

5 Other preferred embodiments of the claimed invention include an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 50 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1.

10 Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of the Clone Sequence and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

15 Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of the Start Codon and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

20 Similarly preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of the First Amino Acid of the Signal Peptide and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

25 Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 150 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X.

Further preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 500 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X.

30 A further preferred embodiment is a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the nucleotide sequence of SEQ ID NO:X beginning with the nucleotide at about the position of the 5' Nucleotide of the First Amino Acid of the Signal Peptide and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the complete nucleotide sequence of SEQ ID NO:X.

5 Also preferred is an isolated nucleic acid molecule which hybridizes under stringent hybridization conditions to a nucleic acid molecule, wherein said nucleic acid molecule which hybridizes does not hybridize under stringent hybridization conditions to a nucleic acid molecule having a nucleotide sequence consisting of only A residues or of only T residues.

10 Also preferred is a composition of matter comprising a DNA molecule which comprises a human cDNA clone identified by a cDNA Clone Identifier in Table 1, which DNA molecule is contained in the material deposited with the American Type Culture Collection and given the ATCC Deposit Number shown in Table 1 for said cDNA Clone Identifier.

15 Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides in the nucleotide sequence of a human cDNA clone identified by a cDNA Clone Identifier in Table 1, which DNA molecule is contained in the deposit given the ATCC Deposit Number shown in Table 1.

20 Also preferred is an isolated nucleic acid molecule, wherein said sequence of at least 50 contiguous nucleotides is included in the nucleotide sequence of the complete open reading frame sequence encoded by said human cDNA clone.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 150 contiguous nucleotides in the nucleotide sequence encoded by said human cDNA clone.

25 A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 500 contiguous nucleotides in the nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the complete nucleotide sequence encoded by said human cDNA clone.

30 A further preferred embodiment is a method for detecting in a biological sample a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1; which method

comprises a step of comparing a nucleotide sequence of at least one nucleic acid molecule in said sample with a sequence selected from said group and determining whether the sequence of said nucleic acid molecule in said sample is at least 95% identical to said selected sequence.

5 Also preferred is the above method wherein said step of comparing sequences comprises determining the extent of nucleic acid hybridization between nucleic acid molecules in said sample and a nucleic acid molecule comprising said sequence selected from said group. Similarly, also preferred is the above method wherein said step of
10 comparing sequences is performed by comparing the nucleotide sequence determined from a nucleic acid molecule in said sample with said sequence selected from said group. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

A further preferred embodiment is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting nucleic acid molecules in said sample, if any, comprising a nucleotide sequence that is at least 95%
15 identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

20 The method for identifying the species, tissue or cell type of a biological sample can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

25 Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a gene encoding a secreted protein identified in Table 1, which method comprises a step of detecting in a biological sample obtained from said subject nucleic acid molecules, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous
30 nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

35 The method for diagnosing a pathological condition can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95%

identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

Also preferred is a composition of matter comprising isolated nucleic acid molecules wherein the nucleotide sequences of said nucleic acid molecules comprise a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1.

Also preferred is a polypeptide, wherein said sequence of contiguous amino acids is included in the amino acid sequence of SEQ ID NO:Y in the range of positions beginning with the residue at about the position of the First Amino Acid of the Secreted Portion and ending with the residue at about the Last Amino Acid of the Open Reading Frame as set forth for SEQ ID NO:Y in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the complete amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is a polypeptide wherein said sequence of contiguous amino acids is included in the amino acid sequence of a secreted portion of the secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is an isolated antibody which binds specifically to a polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is a method for detecting in a biological sample a polypeptide comprising an amino acid sequence which is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1; which method comprises a step of comparing an amino acid sequence of at least one polypeptide molecule in said sample with a sequence selected from said group and determining whether the sequence of said polypeptide molecule in said sample is at least 90% identical to said sequence of at least 10 contiguous amino acids.

Also preferred is the above method wherein said step of comparing an amino acid sequence of at least one polypeptide molecule in said sample with a sequence selected from said group comprises determining the extent of specific binding of polypeptides in said sample to an antibody which binds specifically to a polypeptide

comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is the above method wherein said step of comparing sequences is performed by comparing the amino acid sequence determined from a polypeptide molecule in said sample with said sequence selected from said group.

Also preferred is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting polypeptide molecules in said sample, if any, comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is the above method for identifying the species, tissue or cell type of a biological sample, which method comprises a step of detecting polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the above group.

Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a gene encoding a secreted protein identified in Table 1, which method comprises a step of detecting in a biological sample obtained from said subject polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

In any of these methods, the step of detecting said polypeptide molecules includes using an antibody.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a nucleotide sequence encoding a polypeptide wherein said polypeptide comprises an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated nucleic acid molecule, wherein said nucleotide sequence encoding a polypeptide has been optimized for expression of said polypeptide in a prokaryotic host.

Also preferred is an isolated nucleic acid molecule, wherein said polypeptide comprises an amino acid sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is a method of making a recombinant vector comprising inserting any of the above isolated nucleic acid molecule into a vector. Also preferred is the recombinant vector produced by this method. Also preferred is a method of making a recombinant host cell comprising introducing the vector into a host cell, as well as the recombinant host cell produced by this method.

Also preferred is a method of making an isolated polypeptide comprising culturing this recombinant host cell under conditions such that said polypeptide is expressed and recovering said polypeptide. Also preferred is this method of making an isolated polypeptide, wherein said recombinant host cell is a eukaryotic cell and said polypeptide is a secreted portion of a human secreted protein comprising an amino acid sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y beginning with the residue at the position of the First Amino Acid of the Secreted Portion of SEQ ID NO:Y wherein Y is an integer set forth in Table 1 and said position of the First Amino Acid of the Secreted Portion of SEQ ID NO:Y is defined in Table 1; and an amino acid sequence of a secreted portion of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1. The isolated polypeptide produced by this method is also preferred.

Also preferred is a method of treatment of an individual in need of an increased level of a secreted protein activity, which method comprises administering to such an individual a pharmaceutical composition comprising an amount of an isolated polypeptide, polynucleotide, or antibody of the claimed invention effective to increase the level of said protein activity in said individual.

Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

Examples

Example 1: Isolation of a Selected cDNA Clone From the Deposited Sample

Each cDNA clone in a cited ATCC deposit is contained in a plasmid vector. Table 1 identifies the vectors used to construct the cDNA library from which each clone was isolated. In many cases, the vector used to construct the library is a phage vector from which a plasmid has been excised. The table immediately below correlates the related plasmid for each phage vector used in constructing the cDNA library. For example, where a particular clone is identified in Table 1 as being isolated in the vector "Lambda Zap," the corresponding deposited clone is in "pBluescript."

<u>Vector Used to Construct Library</u>	<u>Corresponding Deposited Plasmid</u>
Lambda Zap	pBluescript (pBS)
Uni-Zap XR	pBluescript (pBS)
Zap Express	pBK
lafmid BA	plafmid BA
pSport1	pSport1
pCMVSPORT 2.0	pCMVSPORT 2.0
pCMVSPORT 3.0	pCMVSPORT 3.0
pCR [®] 2.1	pCR [®] 2.1

Vectors Lambda Zap (U.S. Patent Nos. 5,128,256 and 5,286,636), Uni-Zap XR (U.S. Patent Nos. 5,128, 256 and 5,286,636), Zap Express (U.S. Patent Nos. 5,128,256 and 5,286,636), pBluescript (pBS) (Short, J. M. et al., Nucleic Acids Res. 16:7583-7600 (1988); Alting-Mees, M. A. and Short, J. M., Nucleic Acids Res. 17:9494 (1989)) and pBK (Alting-Mees, M. A. et al., Strategies 5:58-61 (1992)) are commercially available from Stratagene Cloning Systems, Inc., 11011 N. Torrey Pines Road, La Jolla, CA, 92037. pBS contains an ampicillin resistance gene and pBK contains a neomycin resistance gene. Both can be transformed into E. coli strain XL-1

Blue, also available from Stratagene. pBS comes in 4 forms SK+, SK-, KS+ and KS-. The S and K refers to the orientation of the polylinker to the T7 and T3 primer sequences which flank the polylinker region ("S" is for SacI and "K" is for KpnI which are the first sites on each respective end of the linker). "+" or "-" refer to the orientation of the f1 origin of replication ("ori"), such that in one orientation, single stranded rescue initiated from the f1 ori generates sense strand DNA and in the other, antisense.

Vectors pSport1, pCMVSPORT 2.0 and pCMVSPORT 3.0, were obtained from Life Technologies, Inc., P. O. Box 6009, Gaithersburg, MD 20897. All Sport vectors contain an ampicillin resistance gene and may be transformed into E. coli strain DH10B, also available from Life Technologies. (See, for instance, Gruber, C. E., et al., Focus 15:59 (1993).) Vector lacmid BA (Bento Soares, Columbia University, NY) contains an ampicillin resistance gene and can be transformed into E. coli strain XL-1 Blue. Vector pCR[®]2.1, which is available from Invitrogen, 1600 Faraday Avenue, Carlsbad, CA 92008, contains an ampicillin resistance gene and may be transformed into E. coli strain DH10B, available from Life Technologies. (See, for instance, Clark, J. M., Nuc. Acids Res. 16:9677-9686 (1988) and Mead, D. et al., Bio/Technology 9: (1991).) Preferably, a polynucleotide of the present invention does not comprise the phage vector sequences identified for the particular clone in Table 1, as well as the corresponding plasmid vector sequences designated above.

The deposited material in the sample assigned the ATCC Deposit Number cited in Table 1 for any given cDNA clone also may contain one or more additional plasmids, each comprising a cDNA clone different from that given clone. Thus, deposits sharing the same ATCC Deposit Number contain at least a plasmid for each cDNA clone identified in Table 1. Typically, each ATCC deposit sample cited in Table 1 comprises a mixture of approximately equal amounts (by weight) of about 50 plasmid DNAs, each containing a different cDNA clone; but such a deposit sample may include plasmids for more or less than 50 cDNA clones, up to about 500 cDNA clones.

Two approaches can be used to isolate a particular clone from the deposited sample of plasmid DNAs cited for that clone in Table 1. First, a plasmid is directly isolated by screening the clones using a polynucleotide probe corresponding to SEQ ID NO:X.

Particularly, a specific polynucleotide with 30-40 nucleotides is synthesized using an Applied Biosystems DNA synthesizer according to the sequence reported. The oligonucleotide is labeled, for instance, with ³²P-γ-ATP using T4 polynucleotide kinase and purified according to routine methods. (E.g., Maniatis et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, Cold Spring, NY (1982).)

- The plasmid mixture is transformed into a suitable host, as indicated above (such as XL-1 Blue (Stratagene)) using techniques known to those of skill in the art, such as those provided by the vector supplier or in related publications or patents cited above. The transformants are plated on 1.5% agar plates (containing the appropriate selection agent, e.g., ampicillin) to a density of about 150 transformants (colonies) per plate. These plates are screened using Nylon membranes according to routine methods for bacterial colony screening (e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Edit., (1989), Cold Spring Harbor Laboratory Press, pages 1.93 to 1.104), or other techniques known to those of skill in the art.
- Alternatively, two primers of 17-20 nucleotides derived from both ends of the SEQ ID NO:X (i.e., within the region of SEQ ID NO:X bounded by the 5' NT and the 3' NT of the clone defined in Table 1) are synthesized and used to amplify the desired cDNA using the deposited cDNA plasmid as a template. The polymerase chain reaction is carried out under routine conditions, for instance, in 25 μ l of reaction mixture with 0.5 μ g of the above cDNA template. A convenient reaction mixture is 1.5-5 mM $MgCl_2$, 0.01% (w/v) gelatin, 20 μ M each of dATP, dCTP, dGTP, dTTP, 25 pmol of each primer and 0.25 Unit of Taq polymerase. Thirty five cycles of PCR (denaturation at 94°C for 1 min; annealing at 55°C for 1 min; elongation at 72°C for 1 min) are performed with a Perkin-Elmer Cetus automated thermal cycler. The amplified product is analyzed by agarose gel electrophoresis and the DNA band with expected molecular weight is excised and purified. The PCR product is verified to be the selected sequence by subcloning and sequencing the DNA product.
- Several methods are available for the identification of the 5' or 3' non-coding portions of a gene which may not be present in the deposited clone. These methods include but are not limited to, filter probing, clone enrichment using specific probes, and protocols similar or identical to 5' and 3' "RACE" protocols which are well known in the art. For instance, a method similar to 5' RACE is available for generating the missing 5' end of a desired full-length transcript. (Fromont-Racine et al., *Nucleic Acids Res.* 21(7):1683-1684 (1993).)
- Briefly, a specific RNA oligonucleotide is ligated to the 5' ends of a population of RNA presumably containing full-length gene RNA transcripts. A primer set containing a primer specific to the ligated RNA oligonucleotide and a primer specific to a known sequence of the gene of interest is used to PCR amplify the 5' portion of the desired full-length gene. This amplified product may then be sequenced and used to generate the full length gene.

This above method starts with total RNA isolated from the desired source, although poly-A+ RNA can be used. The RNA preparation can then be treated with phosphatase if necessary to eliminate 5' phosphate groups on degraded or damaged RNA which may interfere with the later RNA ligase step. The phosphatase should then
5 be inactivated and the RNA treated with tobacco acid pyrophosphatase in order to remove the cap structure present at the 5' ends of messenger RNAs. This reaction leaves a 5' phosphate group at the 5' end of the cap cleaved RNA which can then be ligated to an RNA oligonucleotide using T4 RNA ligase.

This modified RNA preparation is used as a template for first strand cDNA
10 synthesis using a gene specific oligonucleotide. The first strand synthesis reaction is used as a template for PCR amplification of the desired 5' end using a primer specific to the ligated RNA oligonucleotide and a primer specific to the known sequence of the gene of interest. The resultant product is then sequenced and analyzed to confirm that the 5' end sequence belongs to the desired gene.

15
Example 2: Isolation of Genomic Clones Corresponding to a Polynucleotide

A human genomic P1 library (Genomic Systems, Inc.) is screened by PCR using primers selected for the cDNA sequence corresponding to SEQ ID NO:X.,
20 according to the method described in Example 1. (See also, Sambrook.)

Example 3: Tissue Distribution of Polypeptide

Tissue distribution of mRNA expression of polynucleotides of the present invention is determined using protocols for Northern blot analysis, described by,
25 among others, Sambrook et al. For example, a cDNA probe produced by the method described in Example 1 is labeled with P³² using the rediprime™ DNA labeling system (Amersham Life Science), according to manufacturer's instructions. After labeling, the probe is purified using CHROMA SPIN-100™ column (Clontech Laboratories, Inc.), according to manufacturer's protocol number PT1200-1. The purified labeled probe is
30 then used to examine various human tissues for mRNA expression.

Multiple Tissue Northern (MTN) blots containing various human tissues (H) or human immune system tissues (IM) (Clontech) are examined with the labeled probe using ExpressHyb™ hybridization solution (Clontech) according to manufacturer's protocol number PT1190-1. Following hybridization and washing, the blots are
35 mounted and exposed to film at -70°C overnight, and the films developed according to standard procedures.

Example 4: Chromosomal Mapping of the Polynucleotides

An oligonucleotide primer set is designed according to the sequence at the 5' end of SEQ ID NO:X. This primer preferably spans about 100 nucleotides. This primer set is then used in a polymerase chain reaction under the following set of conditions : 30 seconds, 95°C; 1 minute, 56°C; 1 minute, 70°C. This cycle is repeated 32 times followed by one 5 minute cycle at 70°C. Human, mouse, and hamster DNA is used as template in addition to a somatic cell hybrid panel containing individual chromosomes or chromosome fragments (Bios, Inc). The reactions is analyzed on either 8% polyacrylamide gels or 3.5 % agarose gels. Chromosome mapping is determined by the presence of an approximately 100 bp PCR fragment in the particular somatic cell hybrid.

Example 5: Bacterial Expression of a Polypeptide

A polynucleotide encoding a polypeptide of the present invention is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' ends of the DNA sequence, as outlined in Example 1, to synthesize insertion fragments. The primers used to amplify the cDNA insert should preferably contain restriction sites, such as BamHI and XbaI, at the 5' end of the primers in order to clone the amplified product into the expression vector. For example, BamHI and XbaI correspond to the restriction enzyme sites on the bacterial expression vector pQE-9. (Qiagen, Inc., Chatsworth, CA). This plasmid vector encodes antibiotic resistance (Amp^r), a bacterial origin of replication (ori), an IPTG-regulatable promoter/operator (P/O), a ribosome binding site (RBS), a 6-histidine tag (6-His), and restriction enzyme cloning sites.

The pQE-9 vector is digested with BamHI and XbaI and the amplified fragment is ligated into the pQE-9 vector maintaining the reading frame initiated at the bacterial RBS. The ligation mixture is then used to transform the E. coli strain M15/rep4 (Qiagen, Inc.) which contains multiple copies of the plasmid pREP4, which expresses the lacI repressor and also confers kanamycin resistance (Kan^r). Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies are selected. Plasmid DNA is isolated and confirmed by restriction analysis.

Clones containing the desired constructs are grown overnight (O/N) in liquid culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells are grown to an optical density 600 (O.D.⁶⁰⁰) of between 0.4 and 0.6. IPTG

(Isopropyl-B-D-thiogalacto pyranoside) is then added to a final concentration of 1 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to increased gene expression.

Cells are grown for an extra 3 to 4 hours. Cells are then harvested by centrifugation (20 mins at 6000Xg). The cell pellet is solubilized in the chaotropic agent 6 Molar Guanidine HCl by stirring for 3-4 hours at 4°C. The cell debris is removed by centrifugation, and the supernatant containing the polypeptide is loaded onto a nickel-nitrilo-tri-acetic acid ("Ni-NTA") affinity resin column (available from QIAGEN, Inc., *supra*). Proteins with a 6 x His tag bind to the Ni-NTA resin with high affinity and can be purified in a simple one-step procedure (for details see: The QIAexpressionist (1995) QIAGEN, Inc., *supra*).

Briefly, the supernatant is loaded onto the column in 6 M guanidine-HCl, pH 8, the column is first washed with 10 volumes of 6 M guanidine-HCl, pH 8, then washed with 10 volumes of 6 M guanidine-HCl pH 6, and finally the polypeptide is eluted with 6 M guanidine-HCl, pH 5.

The purified protein is then renatured by dialyzing it against phosphate-buffered saline (PBS) or 50 mM Na-acetate, pH 6 buffer plus 200 mM NaCl. Alternatively, the protein can be successfully refolded while immobilized on the Ni-NTA column. The recommended conditions are as follows: renature using a linear 6M-1M urea gradient in 500 mM NaCl, 20% glycerol, 20 mM Tris/HCl pH 7.4, containing protease inhibitors. The renaturation should be performed over a period of 1.5 hours or more. After renaturation the proteins are eluted by the addition of 250 mM imidazole. Imidazole is removed by a final dialyzing step against PBS or 50 mM sodium acetate pH 6 buffer plus 200 mM NaCl. The purified protein is stored at 4°C or frozen at -80°C.

In addition to the above expression vector, the present invention further includes an expression vector comprising phage operator and promoter elements operatively linked to a polynucleotide of the present invention, called pHE4a. (ATCC Accession Number 209645, deposited on February 25, 1998.) This vector contains: 1) a neomycinphosphotransferase gene as a selection marker, 2) an E. coli origin of replication, 3) a T5 phage promoter sequence, 4) two lac operator sequences, 5) a Shine-Delgarno sequence, and 6) the lactose operon repressor gene (lacIq). The origin of replication (oriC) is derived from pUC19 (LTI, Gaithersburg, MD). The promoter sequence and operator sequences are made synthetically.

DNA can be inserted into the pHEa by restricting the vector with NdeI and XbaI, BamHI, XhoI, or Asp718, running the restricted product on a gel, and isolating the larger fragment (the stuffer fragment should be about 310 base pairs). The DNA

insert is generated according to the PCR protocol described in Example 1, using PCR primers having restriction sites for NdeI (5' primer) and XbaI, BamHI, XhoI, or Asp718 (3' primer). The PCR insert is gel purified and restricted with compatible enzymes. The insert and vector are ligated according to standard protocols.

- 5 The engineered vector could easily be substituted in the above protocol to express protein in a bacterial system.

Example 6: Purification of a Polypeptide from an Inclusion Body

- 10 The following alternative method can be used to purify a polypeptide expressed in *E. coli* when it is present in the form of inclusion bodies. Unless otherwise specified, all of the following steps are conducted at 4-10°C.

- Upon completion of the production phase of the *E. coli* fermentation, the cell culture is cooled to 4-10°C and the cells harvested by continuous centrifugation at 15,000 rpm (Heraeus Sepatech). On the basis of the expected yield of protein per unit weight of cell paste and the amount of purified protein required, an appropriate amount of cell paste, by weight, is suspended in a buffer solution containing 100 mM Tris, 50 mM EDTA, pH 7.4. The cells are dispersed to a homogeneous suspension using a high shear mixer.

- 20 The cells are then lysed by passing the solution through a microfluidizer (Microfluidics, Corp. or APV Gaulin, Inc.) twice at 4000-6000 psi. The homogenate is then mixed with NaCl solution to a final concentration of 0.5 M NaCl, followed by centrifugation at 7000 xg for 15 min. The resultant pellet is washed again using 0.5M NaCl, 100 mM Tris, 50 mM EDTA, pH 7.4.

- 25 The resulting washed inclusion bodies are solubilized with 1.5 M guanidine hydrochloride (GuHCl) for 2-4 hours. After 7000 xg centrifugation for 15 min., the pellet is discarded and the polypeptide containing supernatant is incubated at 4°C overnight to allow further GuHCl extraction.

- 30 Following high speed centrifugation (30,000 xg) to remove insoluble particles, the GuHCl solubilized protein is refolded by quickly mixing the GuHCl extract with 20 volumes of buffer containing 50 mM sodium, pH 4.5, 150 mM NaCl, 2 mM EDTA by vigorous stirring. The refolded diluted protein solution is kept at 4°C without mixing for 12 hours prior to further purification steps.

 To clarify the refolded polypeptide solution, a previously prepared tangential filtration unit equipped with 0.16 µm membrane filter with appropriate surface area

(e.g., Filtron), equilibrated with 40 mM sodium acetate, pH 6.0 is employed. The filtered sample is loaded onto a cation exchange resin (e.g., Poros HS-50, Perseptive Biosystems). The column is washed with 40 mM sodium acetate, pH 6.0 and eluted with 250 mM, 500 mM, 1000 mM, and 1500 mM NaCl in the same buffer, in a stepwise manner. The absorbance at 280 nm of the effluent is continuously monitored. Fractions are collected and further analyzed by SDS-PAGE.

Fractions containing the polypeptide are then pooled and mixed with 4 volumes of water. The diluted sample is then loaded onto a previously prepared set of tandem columns of strong anion (Poros HQ-50, Perseptive Biosystems) and weak anion (Poros CM-20, Perseptive Biosystems) exchange resins. The columns are equilibrated with 40 mM sodium acetate, pH 6.0. Both columns are washed with 40 mM sodium acetate, pH 6.0, 200 mM NaCl. The CM-20 column is then eluted using a 10 column volume linear gradient ranging from 0.2 M NaCl, 50 mM sodium acetate, pH 6.0 to 1.0 M NaCl, 50 mM sodium acetate, pH 6.5. Fractions are collected under constant A_{280} monitoring of the effluent. Fractions containing the polypeptide (determined, for instance, by 16% SDS-PAGE) are then pooled.

The resultant polypeptide should exhibit greater than 95% purity after the above refolding and purification steps. No major contaminant bands should be observed from Commassie blue stained 16% SDS-PAGE gel when 5 μ g of purified protein is loaded. The purified protein can also be tested for endotoxin/LPS contamination, and typically the LPS content is less than 0.1 ng/ml according to LAL assays.

Example 7: Cloning and Expression of a Polypeptide in a Baculovirus

Expression System

In this example, the plasmid shuttle vector pA2 is used to insert a polynucleotide into a baculovirus to express a polypeptide. This expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by convenient restriction sites such as BamHI, Xba I and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the plasmid contains the beta-galactosidase gene from *E. coli* under control of a weak *Drosophila* promoter in the same orientation, followed by the polyadenylation signal of the polyhedrin gene. The inserted genes are flanked on both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate a viable virus that express the cloned polynucleotide.

Many other baculovirus vectors can be used in place of the vector above, such as pAc373, pVL941, and pAcIM1, as one skilled in the art would readily appreciate, as long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as required. Such vectors are described, for instance, in Luckow et al., Virology 170:31-39 (1989).

Specifically, the cDNA sequence contained in the deposited clone, including the AUG initiation codon and the naturally associated leader sequence identified in Table 1, is amplified using the PCR protocol described in Example 1. If the naturally occurring signal sequence is used to produce the secreted protein, the pA2 vector does not need a second signal peptide. Alternatively, the vector can be modified (pA2 GP) to include a baculovirus leader sequence, using the standard methods described in Summers et al., "A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures," Texas Agricultural Experimental Station Bulletin No. 1555 (1987).

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

The plasmid is digested with the corresponding restriction enzymes and optionally, can be dephosphorylated using calf intestinal phosphatase, using routine procedures known in the art. The DNA is then isolated from a 1% agarose gel using a commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, Ca.).

The fragment and the dephosphorylated plasmid are ligated together with T4 DNA ligase. *E. coli* HB101 or other suitable *E. coli* hosts such as XL-1 Blue (Stratagene Cloning Systems, La Jolla, CA) cells are transformed with the ligation mixture and spread on culture plates. Bacteria containing the plasmid are identified by digesting DNA from individual colonies and analyzing the digestion product by gel electrophoresis. The sequence of the cloned fragment is confirmed by DNA sequencing.

Five µg of a plasmid containing the polynucleotide is co-transfected with 1.0 µg of a commercially available linearized baculovirus DNA ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA), using the lipofection method described by Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413-7417 (1987). One µg of BaculoGold™ virus DNA and 5 µg of the plasmid are mixed in a sterile well of a microtiter plate containing 50 µl of serum-free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards, 10 µl Lipofectin plus 90 µl Grace's medium are added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm

tissue culture plate with 1 ml Grace's medium without serum. The plate is then incubated for 5 hours at 27° C. The transfection solution is then removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. Cultivation is then continued at 27° C for four days.

- 5 After four days the supernatant is collected and a plaque assay is performed, as described by Summers and Smith, *supra*. An agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) is used to allow easy identification and isolation of gal-expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10.)
- 10 After appropriate incubation, blue stained plaques are picked with the tip of a micropipettor (e.g., Eppendorf). The agar containing the recombinant viruses is then resuspended in a microcentrifuge tube containing 200 µl of Grace's medium and the suspension containing the recombinant baculovirus is used to infect Sf9 cells seeded in
- 15 35 mm dishes. Four days later the supernatants of these culture dishes are harvested and then they are stored at 4° C.

- To verify the expression of the polypeptide, Sf9 cells are grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells are infected with the recombinant baculovirus containing the polynucleotide at a multiplicity of infection
- 20 ("MOI") of about 2. If radiolabeled proteins are desired, 6 hours later the medium is removed and is replaced with SF900 II medium minus methionine and cysteine (available from Life Technologies Inc., Rockville, MD). After 42 hours, 5 µCi of ³⁵S-methionine and 5 µCi ³⁵S-cysteine (available from Amersham) are added. The cells are further incubated for 16 hours and then are harvested by centrifugation. The proteins
- 25 in the supernatant as well as the intracellular proteins are analyzed by SDS-PAGE followed by autoradiography (if radiolabeled).

Microsequencing of the amino acid sequence of the amino terminus of purified protein may be used to determine the amino terminal sequence of the produced protein.

30 **Example 8: Expression of a Polypeptide in Mammalian Cells**

The polypeptide of the present invention can be expressed in a mammalian cell. A typical mammalian expression vector contains a promoter element, which mediates

the initiation of transcription of mRNA, a protein coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription is achieved with the early and late promoters from SV40, the long terminal repeats (LTRs) from Retroviruses, e.g., RSV, HTLVI, HIVI and the early promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter).

Suitable expression vectors for use in practicing the present invention include, for example, vectors such as pSVL and pMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146), pBC12MI (ATCC 67109), pCMVSPORT 2.0, and pCMVSPORT 3.0. Mammalian host cells that could be used include, human Hela, 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

Alternatively, the polypeptide can be expressed in stable cell lines containing the polynucleotide integrated into a chromosome. The co-transfection with a selectable marker such as dhfr, gpt, neomycin, hygromycin allows the identification and isolation of the transfected cells.

The transfected gene can also be amplified to express large amounts of the encoded protein. The DHFR (dihydrofolate reductase) marker is useful in developing cell lines that carry several hundred or even several thousand copies of the gene of interest. (See, e.g., Alt, F. W., et al., J. Biol. Chem. 253:1357-1370 (1978); Hamlin, J. L. and Ma, C., Biochem. et Biophys. Acta, 1097:107-143 (1990); Page, M. J. and Sydenham, M. A., Biotechnology 9:64-68 (1991).) Another useful selection marker is the enzyme glutamine synthase (GS) (Murphy et al., Biochem J. 227:277-279 (1991); Bebbington et al., Bio/Technology 10:169-175 (1992). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the production of proteins.

Derivatives of the plasmid pSV2-dhfr (ATCC Accession No. 37146), the expression vectors pC4 (ATCC Accession No. 209646) and pC6 (ATCC Accession No. 209647) contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen et al., Molecular and Cellular Biology, 438-447 (March, 1985)) plus a fragment of the CMV-enhancer (Boshart et al., Cell 41:521-530 (1985).) Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the cloning of the gene of interest. The vectors also contain the 3' intron, the

polyadenylation and termination signal of the rat preproinsulin gene, and the mouse DHFR gene under control of the SV40 early promoter.

Specifically, the plasmid pC6, for example, is digested with appropriate restriction enzymes and then dephosphorylated using calf intestinal phosphates by procedures known in the art. The vector is then isolated from a 1% agarose gel.

A polynucleotide of the present invention is amplified according to the protocol outlined in Example 1. If the naturally occurring signal sequence is used to produce the secreted protein, the vector does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., WO 96/34891.)

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("Geneclean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

The amplified fragment is then digested with the same restriction enzyme and purified on a 1% agarose gel. The isolated fragment and the dephosphorylated vector are then ligated with T4 DNA ligase. *E. coli* HB101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid pC6 using, for instance, restriction enzyme analysis.

Chinese hamster ovary cells lacking an active DHFR gene is used for transfection. Five μ g of the expression plasmid pC6 is cotransfected with 0.5 μ g of the plasmid pSVneo using lipofectin (Felgner et al., *supra*). The plasmid pSV2-neo contains a dominant selectable marker, the *neo* gene from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of methotrexate plus 1 mg/ml G418. After about 10-14 days single clones are trypsinized and then seeded in 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1 μ M, 2 μ M, 5 μ M, 10 mM, 20 mM). The same procedure is repeated until clones are obtained which grow at a concentration of 100 - 200 μ M. Expression of the desired gene product is analyzed, for instance, by SDS-PAGE and Western blot or by reversed phase HPLC analysis.

Example 9: Protein Fusions

The polypeptides of the present invention are preferably fused to other proteins. These fusion proteins can be used for a variety of applications. For example, fusion of the present polypeptides to His-tag, HA-tag, protein A, IgG domains, and maltose binding protein facilitates purification. (See Example 5; see also EP A 394,827; Traunecker, et al., Nature 331:84-86 (1988).) Similarly, fusion to IgG-1, IgG-3, and albumin increases the half-life time in vivo. Nuclear localization signals fused to the polypeptides of the present invention can target the protein to a specific subcellular localization, while covalent heterodimer or homodimers can increase or decrease the activity of a fusion protein. Fusion proteins can also create chimeric molecules having more than one function. Finally, fusion proteins can increase solubility and/or stability of the fused protein compared to the non-fused protein. All of the types of fusion proteins described above can be made by modifying the following protocol, which outlines the fusion of a polypeptide to an IgG molecule, or the protocol described in Example 5.

Briefly, the human Fc portion of the IgG molecule can be PCR amplified, using primers that span the 5' and 3' ends of the sequence described below. These primers also should have convenient restriction enzyme sites that will facilitate cloning into an expression vector, preferably a mammalian expression vector.

For example, if pC4 (Accession No. 209646) is used, the human Fc portion can be ligated into the BamHI cloning site. Note that the 3' BamHI site should be destroyed. Next, the vector containing the human Fc portion is re-restricted with BamHI, linearizing the vector, and a polynucleotide of the present invention, isolated by the PCR protocol described in Example 1, is ligated into this BamHI site. Note that the polynucleotide is cloned without a stop codon, otherwise a fusion protein will not be produced.

If the naturally occurring signal sequence is used to produce the secreted protein, pC4 does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., WO 96/34891.)

Human IgG Fc region:

```
GGGATCCGGAGCCCAAATCTTCTGACAAAACTCACACATGCCCACCGTGCC
CAGCACCTGAATTGAGGGTGACCGTCAGTCTTCCTCTTCCCCCAAACC
35 CAAGGACACCCTCATGATCTCCCGGACTCCTGAGGTCACATGCGTGGTGGT
GGACGTAAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACG
GCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAAC
```

AGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTG
 AATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCAACCCCC
 ATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGT
 GTACACCCTGCCCCCATCCCGGGATGAGCTGACCAAGAACCAGGTCAGCCT
 5 GACCTGCCTGGTCAAAGGCTTCTATCCAAGCGACATCGCCGTGGAGTGGGA
 GAGCAATGGGCAGCCGGAGAACAACACTACAAGACCACGCCTCCCGTGCTGG
 ACTCCGACGGCTCCTTCTCCTCTACAGCAAGCTCACCGTGGACAAGAGCA
 GGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGC
 ACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAATGAGTGC
 10 GACGGCCGCGACTCTAGAGGAT (SEQ ID NO:1)

Example 10: Production of an Antibody from a Polypeptide

The antibodies of the present invention can be prepared by a variety of methods. (See, Current Protocols, Chapter 2.) For example, cells expressing a polypeptide of the present invention is administered to an animal to induce the production of sera containing polyclonal antibodies. In a preferred method, a preparation of the secreted protein is prepared and purified to render it substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

20 In the most preferred method, the antibodies of the present invention are monoclonal antibodies (or protein binding fragments thereof). Such monoclonal antibodies can be prepared using hybridoma technology. (Köhler et al., Nature 256:495 (1975); Köhler et al., Eur. J. Immunol. 6:511 (1976); Köhler et al., Eur. J. Immunol. 6:292 (1976); Hammerling et al., in: Monoclonal Antibodies and T-Cell
 25 Hybridomas, Elsevier, N.Y., pp. 563-681 (1981).) In general, such procedures involve immunizing an animal (preferably a mouse) with polypeptide or, more preferably, with a secreted polypeptide-expressing cell. Such cells may be cultured in any suitable tissue culture medium; however, it is preferable to culture cells in Earle's modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at
 30 about 56°C), and supplemented with about 10 g/l of nonessential amino acids, about 1,000 U/ml of penicillin, and about 100 µg/ml of streptomycin.

The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line
 35 (SP2O), available from the ATCC. After fusion, the resulting hybridoma cells are selectively maintained in HAT medium, and then cloned by limiting dilution as

described by Wands et al. (Gastroenterology 80:225-232 (1981).) The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the polypeptide.

Alternatively, additional antibodies capable of binding to the polypeptide can be produced in a two-step procedure using anti-idiotypic antibodies. Such a method makes use of the fact that antibodies are themselves antigens, and therefore, it is possible to obtain an antibody which binds to a second antibody. In accordance with this method, protein specific antibodies are used to immunize an animal, preferably a mouse. The splenocytes of such an animal are then used to produce hybridoma cells, and the hybridoma cells are screened to identify clones which produce an antibody whose ability to bind to the protein-specific antibody can be blocked by the polypeptide. Such antibodies comprise anti-idiotypic antibodies to the protein-specific antibody and can be used to immunize an animal to induce formation of further protein-specific antibodies.

It will be appreciated that Fab and F(ab')₂ and other fragments of the antibodies of the present invention may be used according to the methods disclosed herein. Such fragments are typically produced by proteolytic cleavage, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂ fragments). Alternatively, secreted protein-binding fragments can be produced through the application of recombinant DNA technology or through synthetic chemistry.

For in vivo use of antibodies in humans, it may be preferable to use "humanized" chimeric monoclonal antibodies. Such antibodies can be produced using genetic constructs derived from hybridoma cells producing the monoclonal antibodies described above. Methods for producing chimeric antibodies are known in the art. (See, for review, Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., WO 8702671; Boulianne et al., Nature 312:643 (1984); Neuberger et al., Nature 314:268 (1985).)

Example 11: Production Of Secreted Protein For High-Throughput Screening Assays

The following protocol produces a supernatant containing a polypeptide to be tested. This supernatant can then be used in the Screening Assays described in Examples 13-20.

First, dilute Poly-D-Lysine (644 587 Boehringer-Mannheim) stock solution (1mg/ml in PBS) 1:20 in PBS (w/o calcium or magnesium 17-516F Biowhittaker) for a

working solution of 50ug/ml. Add 200 ul of this solution to each well (24 well plates) and incubate at RT for 20 minutes. Be sure to distribute the solution over each well (note: a 12-channel pipetter may be used with tips on every other channel). Aspirate off the Poly-D-Lysine solution and rinse with 1ml PBS (Phosphate Buffered Saline). The
 5 PBS should remain in the well until just prior to plating the cells and plates may be poly-lysine coated in advance for up to two weeks.

Plate 293T cells (do not carry cells past P+20) at 2×10^5 cells/well in .5ml DMEM(Dulbecco's Modified Eagle Medium)(with 4.5 G/L glucose and L-glutamine (12-604F Biowhittaker))/10% heat inactivated FBS(14-503F Biowhittaker)/1x
 10 Penstrep(17-602E Biowhittaker). Let the cells grow overnight.

The next day, mix together in a sterile solution basin: 300 ul Lipofectamine (18324-012 Gibco/BRL) and 5ml Optimem I (31985070 Gibco/BRL)/96-well plate. With a small volume multi-channel pipetter, aliquot approximately 2ug of an expression vector containing a polynucleotide insert, produced by the methods described in
 15 Examples 8 or 9, into an appropriately labeled 96-well round bottom plate. With a multi-channel pipetter, add 50ul of the Lipofectamine/Optimem I mixture to each well. Pipette up and down gently to mix. Incubate at RT 15-45 minutes. After about 20 minutes, use a multi-channel pipetter to add 150ul Optimem I to each well. As a control, one plate of vector DNA lacking an insert should be transfected with each set of
 20 transfections.

Preferably, the transfection should be performed by tag-teaming the following tasks. By tag-teaming, hands on time is cut in half, and the cells do not spend too much time on PBS. First, person A aspirates off the media from four 24-well plates of cells, and then person B rinses each well with .5-1ml PBS. Person A then aspirates off
 25 PBS rinse, and person B, using a 12-channel pipetter with tips on every other channel, adds the 200ul of DNA/Lipofectamine/Optimem I complex to the odd wells first, then to the even wells, to each row on the 24-well plates. Incubate at 37°C for 6 hours.

While cells are incubating, prepare appropriate media, either 1%BSA in DMEM with 1x penstrep, or CHO-5 media (116.6 mg/L of CaCl₂ (anhyd); 0.00130 mg/L
 30 CuSO₄·5H₂O; 0.050 mg/L of Fe(NO₃)₃·9H₂O; 0.417 mg/L of FeSO₄·7H₂O; 311.80 mg/L of KCl; 28.64 mg/L of MgCl₂; 48.84 mg/L of MgSO₄; 6995.50 mg/L of NaCl; 2400.0 mg/L of NaHCO₃; 62.50 mg/L of NaH₂PO₄·H₂O; 71.02 mg/L of Na₂HPO₄; .4320 mg/L of ZnSO₄·7H₂O; .002 mg/L of Arachidonic Acid ; 1.022 mg/L of Cholesterol; .070 mg/L of DL-alpha-Tocopherol-Acetate; 0.0520 mg/L of Linoleic
 35 Acid; 0.010 mg/L of Linolenic Acid; 0.010 mg/L of Myristic Acid; 0.010 mg/L of Oleic Acid; 0.010 mg/L of Palmitic Acid; 0.010 mg/L of Palmitic Acid; 100 mg/L of

Pluronic F-68; 0.010 mg/L of Stearic Acid; 2.20 mg/L of Tween 80; 4551 mg/L of D-Glucose; 130.85 mg/ml of L- Alanine; 147.50 mg/ml of L-Arginine-HCL; 7.50 mg/ml of L-Asparagine-H₂O; 6.65 mg/ml of L-Aspartic Acid; 29.56 mg/ml of L-Cystine-2HCL-H₂O; 31.29 mg/ml of L-Cystine-2HCL; 7.35 mg/ml of L-Glutamic Acid; 365.0 mg/ml of L-Glutamine; 18.75 mg/ml of Glycine; 52.48 mg/ml of L-Histidine-HCL-H₂O; 106.97 mg/ml of L-Isoleucine; 111.45 mg/ml of L-Leucine; 163.75 mg/ml of L-Lysine HCL; 32.34 mg/ml of L-Methionine; 68.48 mg/ml of L-Phenylalanine; 40.0 mg/ml of L-Proline; 26.25 mg/ml of L-Serine; 101.05 mg/ml of L-Threonine; 19.22 mg/ml of L-Tryptophan; 91.79 mg/ml of L-Tyrosine-2Na-2H₂O; 99.65 mg/ml of L-Valine; 0.0035 mg/L of Biotin; 3.24 mg/L of D-Ca Pantothenate; 11.78 mg/L of Choline Chloride; 4.65 mg/L of Folic Acid; 15.60 mg/L of i-Inositol; 3.02 mg/L of Niacinamide; 3.00 mg/L of Pyridoxal HCL; 0.031 mg/L of Pyridoxine HCL; 0.319 mg/L of Riboflavin; 3.17 mg/L of Thiamine HCL; 0.365 mg/L of Thymidine; and 0.680 mg/L of Vitamin B₁₂; 25 mM of HEPES Buffer; 2.39 mg/L of Na Hypoxanthine; 0.105 mg/L of Lipoic Acid; 0.081 mg/L of Sodium Putrescine-2HCL; 55.0 mg/L of Sodium Pyruvate; 0.0067 mg/L of Sodium Selenite; 20uM of Ethanolamine; 0.122 mg/L of Ferric Citrate; 41.70 mg/L of Methyl-B-Cyclodextrin complexed with Linoleic Acid; 33.33 mg/L of Methyl-B-Cyclodextrin complexed with Oleic Acid; and 10 mg/L of Methyl-B-Cyclodextrin complexed with Retinal) with 2mm glutamine and 1x penstrep. (BSA (81-068-3 Bayer) 100gm dissolved in 1L DMEM for a 10% BSA stock solution). Filter the media and collect 50 ul for endotoxin assay in 15ml polystyrene conical.

The transfection reaction is terminated, preferably by tag-teaming, at the end of the incubation period. Person A aspirates off the transfection media, while person B adds 1.5ml appropriate media to each well. Incubate at 37°C for 45 or 72 hours depending on the media used: 1%BSA for 45 hours or CHO-5 for 72 hours.

On day four, using a 300ul multichannel pipetter, aliquot 600ul in one 1ml deep well plate and the remaining supernatant into a 2ml deep well. The supernatants from each well can then be used in the assays described in Examples 13-20.

It is specifically understood that when activity is obtained in any of the assays described below using a supernatant, the activity originates from either the polypeptide directly (e.g., as a secreted protein) or by the polypeptide inducing expression of other proteins, which are then secreted into the supernatant. Thus, the invention further provides a method of identifying the protein in the supernatant characterized by an activity in a particular assay.

Example 12: Construction of GAS Reporter Construct

One signal transduction pathway involved in the differentiation and proliferation of cells is called the Jaks-STATs pathway. Activated proteins in the Jaks-STATs pathway bind to gamma activation site "GAS" elements or interferon-sensitive responsive element ("ISRE"), located in the promoter of many genes. The binding of a protein to these elements alter the expression of the associated gene.

GAS and ISRE elements are recognized by a class of transcription factors called Signal Transducers and Activators of Transcription, or "STATs." There are six members of the STATs family. Stat1 and Stat3 are present in many cell types, as is Stat2 (as response to IFN-alpha is widespread). Stat4 is more restricted and is not in many cell types though it has been found in T helper class I, cells after treatment with IL-12. Stat5 was originally called mammary growth factor, but has been found at higher concentrations in other cells including myeloid cells. It can be activated in tissue culture cells by many cytokines.

The STATs are activated to translocate from the cytoplasm to the nucleus upon tyrosine phosphorylation by a set of kinases known as the Janus Kinase ("Jaks") family. Jaks represent a distinct family of soluble tyrosine kinases and include Tyk2, Jak1, Jak2, and Jak3. These kinases display significant sequence similarity and are generally catalytically inactive in resting cells.

The Jaks are activated by a wide range of receptors summarized in the Table below. (Adapted from review by Schidler and Darnell, Ann. Rev. Biochem. 64:621-51 (1995).) A cytokine receptor family, capable of activating Jaks, is divided into two groups: (a) Class 1 includes receptors for IL-2, IL-3, IL-4, IL-6, IL-7, IL-9, IL-11, IL-12, IL-15, Epo, PRL, GH, G-CSF, GM-CSF, LIF, CNTF, and thrombopoietin; and (b) Class 2 includes IFN-a, IFN-g, and IL-10. The Class 1 receptors share a conserved cysteine motif (a set of four conserved cysteines and one tryptophan) and a WSXWS motif (a membrane proximal region encoding Trp-Ser-Xxx-Trp-Ser (SEQ ID NO:2)).

Thus, on binding of a ligand to a receptor, Jaks are activated, which in turn activate STATs, which then translocate and bind to GAS elements. This entire process is encompassed in the Jaks-STATs signal transduction pathway.

Therefore, activation of the Jaks-STATs pathway, reflected by the binding of the GAS or the ISRE element, can be used to indicate proteins involved in the proliferation and differentiation of cells. For example, growth factors and cytokines are known to activate the Jaks-STATs pathway. (See Table below.) Thus, by using GAS elements linked to reporter molecules, activators of the Jaks-STATs pathway can be identified.

	<u>Ligand</u>	<u>tyk2</u>	<u>JAKs</u> <u>Jak1</u>	<u>Jak2</u>	<u>Jak3</u>	<u>STATS</u>	<u>GAS(elements) or ISRE</u>
<u>IFN family</u>							
5	IFN-a/B	+	+	-	-	1,2,3	ISRE
	IFN-g		+	+	-	1	GAS (IRF1>Lys6>IFP)
	Il-10	+	?	?	-	1,3	
<u>gp130 family</u>							
10	IL-6 (Pleiotrohic)	+	+	+	?	1,3	GAS (IRF1>Lys6>IFP)
	Il-11(Pleiotrohic)	?	+	?	?	1,3	
	OnM(Pleiotrohic)	?	+	+	?	1,3	
	LIF(Pleiotrohic)	?	+	+	?	1,3	
	CNTF(Pleiotrohic)	-/+	+	+	?	1,3	
15	G-CSF(Pleiotrohic)	?	+	?	?	1,3	
	IL-12(Pleiotrohic)	+	-	+	+	1,3	
<u>g-C family</u>							
20	IL-2 (lymphocytes)	-	+	-	+	1,3,5	GAS
	IL-4 (lymph/myeloid)	-	+	-	+	6	GAS (IRF1 = IFP >>Ly6)(IgH)
	IL-7 (lymphocytes)	-	+	-	+	5	GAS
	IL-9 (lymphocytes)	-	+	-	+	5	GAS
	IL-13 (lymphocyte)	-	+	?	?	6	GAS
	IL-15	?	+	?	+	5	GAS
25	<u>gp140 family</u>						
	IL-3 (myeloid)	-	-	+	-	5	GAS (IRF1>IFP>>Ly6)
	IL-5 (myeloid)	-	-	+	-	5	GAS
	GM-CSF (myeloid)	-	-	+	-	5	GAS
30	<u>Growth hormone family</u>						
	GH	?	-	+	-	5	
	PRL	?	+/-	+	-	1,3,5	
	EPO	?	-	+	-	5	GAS(B-CAS>IRF1=IFP>>Ly6)
35	<u>Receptor Tyrosine Kinases</u>						
	EGF	?	+	+	-	1,3	GAS (IRF1)
	PDGF	?	+	+	-	1,3	
	CSF-1	?	+	+	-	1,3	GAS (not IRF1)
40							

To construct a synthetic GAS containing promoter element, which is used in the Biological Assays described in Examples 13-14, a PCR based strategy is employed to generate a GAS-SV40 promoter sequence. The 5' primer contains four tandem copies of the GAS binding site found in the IRF1 promoter and previously demonstrated to bind STATs upon induction with a range of cytokines (Rothman et al., Immunity 1:457-468 (1994).), although other GAS or ISRE elements can be used instead. The 5' primer also contains 18bp of sequence complementary to the SV40 early promoter sequence and is flanked with an XhoI site. The sequence of the 5' primer is:
 5':GCGCCTCGAGATTCCCCGAAATCTAGATTCCCCGAAATGATTCCCCG
 10 AAATGATTCCCCGAAATATCTGCCATCTCAATTAG:3' (SEQ ID NO:3)

The downstream primer is complementary to the SV40 promoter and is flanked with a Hind III site: 5':GCGGCAAGCTTTTGGCAAAGCCTAGGC:3' (SEQ ID NO:4)

PCR amplification is performed using the SV40 promoter template present in the B-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI/Hind III and subcloned into BLSK2-. (Stratagene.) Sequencing with forward and reverse primers confirms that the insert contains the following sequence:
 5':CTCGAGATTCCCCGAAATCTAGATTCCCCGAAATGATTCCCCGAAATG
 20 ATTTCCCCGAAATATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCGCCC
 CTAAGTCCGCCCATCCCGCCCCTAACTCCGCCCAGTCCGCCCATTCTCCGC
 CCCATGGCTGACTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCTCGGC
 CTCTGAGCTATTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTT
 TGCAAAAAGCTT:3' (SEQ ID NO:5)

With this GAS promoter element linked to the SV40 promoter, a GAS:SEAP2 reporter construct is next engineered. Here, the reporter molecule is a secreted alkaline phosphatase, or "SEAP." Clearly, however, any reporter molecule can be instead of SEAP, in this or in any of the other Examples. Well known reporter molecules that can be used instead of SEAP include chloramphenicol acetyltransferase (CAT), luciferase, alkaline phosphatase, B-galactosidase, green fluorescent protein (GFP), or any protein detectable by an antibody.

The above sequence confirmed synthetic GAS-SV40 promoter element is subcloned into the pSEAP-Promoter vector obtained from Clontech using HindIII and XhoI, effectively replacing the SV40 promoter with the amplified GAS:SV40 promoter element, to create the GAS-SEAP vector. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

Thus, in order to generate mammalian stable cell lines expressing the GAS-SEAP reporter, the GAS-SEAP cassette is removed from the GAS-SEAP vector using SalI and NotI, and inserted into a backbone vector containing the neomycin resistance gene, such as pGFP-1 (Clontech), using these restriction sites in the multiple cloning site, to create the GAS-SEAP/Neo vector. Once this vector is transfected into mammalian cells, this vector can then be used as a reporter molecule for GAS binding as described in Examples 13-14.

Other constructs can be made using the above description and replacing GAS with a different promoter sequence. For example, construction of reporter molecules containing NFK-B and EGR promoter sequences are described in Examples 15 and 16. However, many other promoters can be substituted using the protocols described in these Examples. For instance, SRE, IL-2, NFAT, or Osteocalcin promoters can be substituted, alone or in combination (e.g., GAS/NF-KB/EGR, GAS/NF-KB, IL-2/NFAT, or NF-KB/GAS). Similarly, other cell lines can be used to test reporter construct activity, such as HELA (epithelial), HUVEC (endothelial), Reh (B-cell), Saos-2 (osteoblast), HUVAC (aortic), or Cardiomyocyte.

Example 13: High-Throughput Screening Assay for T-cell Activity.

The following protocol is used to assess T-cell activity by identifying factors, such as growth factors and cytokines, that may proliferate or differentiate T-cells. T-cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 12. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The T-cell used in this assay is Jurkat T-cells (ATCC Accession No. TIB-152), although Molt-3 cells (ATCC Accession No. CRL-1552) and Molt-4 cells (ATCC Accession No. CRL-1582) cells can also be used.

Jurkat T-cells are lymphoblastic CD4+ Th1 helper cells. In order to generate stable cell lines, approximately 2 million Jurkat cells are transfected with the GAS-SEAP/neo vector using DMRIE-C (Life Technologies)(transfection procedure described below). The transfected cells are seeded to a density of approximately 20,000 cells per well and transfectants resistant to 1 mg/ml gentamicin selected. Resistant colonies are expanded and then tested for their response to increasing concentrations of interferon gamma. The dose response of a selected clone is demonstrated.

Specifically, the following protocol will yield sufficient cells for 75 wells containing 200 ul of cells. Thus, it is either scaled up, or performed in multiple to generate sufficient cells for multiple 96 well plates. Jurkat cells are maintained in RPMI + 10% serum with 1%Pen-Strep. Combine 2.5 mls of OPTI-MEM (Life Technologies)

with 10 ug of plasmid DNA in a T25 flask. Add 2.5 ml OPTI-MEM containing 50 ul of DMRIE-C and incubate at room temperature for 15-45 mins.

- During the incubation period, count cell concentration, spin down the required number of cells (10^7 per transfection), and resuspend in OPTI-MEM to a final concentration of 10^7 cells/ml. Then add 1ml of 1×10^7 cells in OPTI-MEM to T25 flask and incubate at 37°C for 6 hrs. After the incubation, add 10 ml of RPMI + 15% serum.

The Jurkat:GAS-SEAP stable reporter lines are maintained in RPMI + 10% serum, 1 mg/ml Gentamicin, and 1% Pen-Strep. These cells are treated with supernatants containing a polypeptide as produced by the protocol described in Example 11.

- On the day of treatment with the supernatant, the cells should be washed and resuspended in fresh RPMI + 10% serum to a density of 500,000 cells per ml. The exact number of cells required will depend on the number of supernatants being screened. For one 96 well plate, approximately 10 million cells (for 10 plates, 100 million cells) are required.

- Transfer the cells to a triangular reservoir boat, in order to dispense the cells into a 96 well dish, using a 12 channel pipette. Using a 12 channel pipette, transfer 200 ul of cells into each well (therefore adding 100, 000 cells per well).

- After all the plates have been seeded, 50 ul of the supernatants are transferred directly from the 96 well plate containing the supernatants into each well using a 12 channel pipette. In addition, a dose of exogenous interferon gamma (0.1, 1.0, 10 ng) is added to wells H9, H10, and H11 to serve as additional positive controls for the assay.

- The 96 well dishes containing Jurkat cells treated with supernatants are placed in an incubator for 48 hrs (note: this time is variable between 48-72 hrs). 35 ul samples from each well are then transferred to an opaque 96 well plate using a 12 channel pipette. The opaque plates should be covered (using sellophane covers) and stored at -20°C until SEAP assays are performed according to Example 17. The plates containing the remaining treated cells are placed at 4°C and serve as a source of material for repeating the assay on a specific well if desired.

- As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate Jurkat T cells. Over 30 fold induction is typically observed in the positive control wells.

Example 14: High-Throughput Screening Assay Identifying Myeloid Activity

The following protocol is used to assess myeloid activity by identifying factors, such as growth factors and cytokines, that may proliferate or differentiate myeloid cells.

- 5 Myeloid cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 12. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The myeloid cell used in this assay is U937, a pre-monocyte cell line, although TF-1, HL60, or KG1 can be used.

- 10 To transiently transfect U937 cells with the GAS/SEAP/Neo construct produced in Example 12, a DEAE-Dextran method (Kharbanda et. al., 1994, Cell Growth & Differentiation, 5:259-265) is used. First, harvest 2×10^7 U937 cells and wash with PBS. The U937 cells are usually grown in RPMI 1640 medium containing 10% heat-inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 mg/ml streptomycin.

- 15 Next, suspend the cells in 1 ml of 20 mM Tris-HCl (pH 7.4) buffer containing 0.5 mg/ml DEAE-Dextran, 8 ug GAS-SEAP2 plasmid DNA, 140 mM NaCl, 5 mM KCl, 375 uM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 1 mM MgCl_2 , and 675 uM CaCl_2 . Incubate at 37°C for 45 min.

- 20 Wash the cells with RPMI 1640 medium containing 10% FBS and then resuspend in 10 ml complete medium and incubate at 37°C for 36 hr.

The GAS-SEAP/U937 stable cells are obtained by growing the cells in 400 ug/ml G418. The G418-free medium is used for routine growth but every one to two months, the cells should be re-grown in 400 ug/ml G418 for couple of passages.

- 25 These cells are tested by harvesting 1×10^8 cells (this is enough for ten 96-well plates assay) and wash with PBS. Suspend the cells in 200 ml above described growth medium, with a final density of 5×10^5 cells/ml. Plate 200 ul cells per well in the 96-well plate (or 1×10^5 cells/well).

- 30 Add 50 ul of the supernatant prepared by the protocol described in Example 11. Incubate at 37°C for 48 to 72 hr. As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate U937 cells. Over 30 fold induction is typically observed in the positive control wells. SEAP assay the supernatant according to the protocol described in Example 17.

Example 15: High-Throughput Screening Assay Identifying Neuronal Activity.

When cells undergo differentiation and proliferation, a group of genes are activated through many different signal transduction pathways. One of these genes,
5 EGR1 (early growth response gene 1), is induced in various tissues and cell types upon activation. The promoter of EGR1 is responsible for such induction. Using the EGR1 promoter linked to reporter molecules, activation of cells can be assessed.

Particularly, the following protocol is used to assess neuronal activity in PC12 cell lines. PC12 cells (rat phenochromocytoma cells) are known to proliferate and/or
10 differentiate by activation with a number of mitogens, such as TPA (tetradecanoyl phorbol acetate), NGF (nerve growth factor), and EGF (epidermal growth factor). The EGR1 gene expression is activated during this treatment. Thus, by stably transfecting PC12 cells with a construct containing an EGR promoter linked to SEAP reporter, activation of PC12 cells can be assessed.

15 The EGR/SEAP reporter construct can be assembled by the following protocol. The EGR-1 promoter sequence (-633 to +1)(Sakamoto K et al., Oncogene 6:867-871 (1991)) can be PCR amplified from human genomic DNA using the following primers:

5' GCGCTCGAGGGATGACAGCGATAGAACCCCGG -3' (SEQ ID NO:6)

5' GCGAAGCTTCGCGACTCCCCGGATCCGCCTC-3' (SEQ ID NO:7)

20 Using the GAS:SEAP/Neo vector produced in Example 12, EGR1 amplified product can then be inserted into this vector. Linearize the GAS:SEAP/Neo vector using restriction enzymes XhoI/HindIII, removing the GAS/SV40 stuffer. Restrict the EGR1 amplified product with these same enzymes. Ligate the vector and the EGR1 promoter.

25 To prepare 96 well-plates for cell culture, two mls of a coating solution (1:30 dilution of collagen type I (Upstate Biotech Inc. Cat#08-115) in 30% ethanol (filter sterilized)) is added per one 10 cm plate or 50 ml per well of the 96-well plate, and allowed to air dry for 2 hr.

PC12 cells are routinely grown in RPMI-1640 medium (Bio Whittaker)
30 containing 10% horse serum (JRH BIOSCIENCES, Cat. # 12449-78P), 5% heat-inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 ug/ml streptomycin on a precoated 10 cm tissue culture dish. One to four split is done every three to four days. Cells are removed from the plates by scraping and resuspended with pipetting up and down for more than 15 times.

35 Transfect the EGR/SEAP/Neo construct into PC12 using the Lipofectamine protocol described in Example 11. EGR-SEAP/PC12 stable cells are obtained by growing the cells in 300 ug/ml G418. The G418-free medium is used for routine

growth but every one to two months, the cells should be re-grown in 300 ug/ml G418 for couple of passages.

To assay for neuronal activity, a 10 cm plate with cells around 70 to 80% confluent is screened by removing the old medium. Wash the cells once with PBS (Phosphate buffered saline). Then starve the cells in low serum medium (RPMI-1640 containing 1% horse serum and 0.5% FBS with antibiotics) overnight.

The next morning, remove the medium and wash the cells with PBS. Scrape off the cells from the plate, suspend the cells well in 2 ml low serum medium. Count the cell number and add more low serum medium to reach final cell density as 5×10^5 cells/ml.

Add 200 ul of the cell suspension to each well of 96-well plate (equivalent to 1×10^5 cells/well). Add 50 ul supernatant produced by Example 11, 37°C for 48 to 72 hr. As a positive control, a growth factor known to activate PC12 cells through EGR can be used, such as 50 ng/ul of Neuronal Growth Factor (NGF). Over fifty-fold induction of SEAP is typically seen in the positive control wells. SEAP assay the supernatant according to Example 17.

Example 16: High-Throughput Screening Assay for T-cell Activity

NF- κ B (Nuclear Factor κ B) is a transcription factor activated by a wide variety of agents including the inflammatory cytokines IL-1 and TNF, CD30 and CD40, lymphotoxin-alpha and lymphotoxin-beta, by exposure to LPS or thrombin, and by expression of certain viral gene products. As a transcription factor, NF- κ B regulates the expression of genes involved in immune cell activation, control of apoptosis (NF- κ B appears to shield cells from apoptosis), B and T-cell development, anti-viral and antimicrobial responses, and multiple stress responses.

In non-stimulated conditions, NF- κ B is retained in the cytoplasm with I- κ B (Inhibitor κ B). However, upon stimulation, I- κ B is phosphorylated and degraded, causing NF- κ B to shuttle to the nucleus, thereby activating transcription of target genes. Target genes activated by NF- κ B include IL-2, IL-6, GM-CSF, ICAM-1 and class 1 MHC.

Due to its central role and ability to respond to a range of stimuli, reporter constructs utilizing the NF- κ B promoter element are used to screen the supernatants produced in Example 11. Activators or inhibitors of NF- κ B would be useful in treating

diseases. For example, inhibitors of NF- κ B could be used to treat those diseases related to the acute or chronic activation of NF- κ B, such as rheumatoid arthritis.

To construct a vector containing the NF- κ B promoter element, a PCR based strategy is employed. The upstream primer contains four tandem copies of the NF- κ B binding site (GGGGACTTTCCC) (SEQ ID NO:8), 18 bp of sequence complementary to the 5' end of the SV40 early promoter sequence, and is flanked with an XhoI site:
5':GCGGCCTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGAC
TTCCATCCTGCCATCTCAATTAG:3' (SEQ ID NO:9)

The downstream primer is complementary to the 3' end of the SV40 promoter and is flanked with a Hind III site:
5':GCGGCAAGCTTTTTGCAAAGCCTAGGC:3' (SEQ ID NO:4)

PCR amplification is performed using the SV40 promoter template present in the pB-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI and Hind III and subcloned into BLSK2-. (Stratagene)
15 Sequencing with the T7 and T3 primers confirms the insert contains the following sequence:

5':CTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGACTTTCC
ATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCGCCCCCTAACTCCGCCCA
20 TCCCGCCCCCTAACTCCGCCCAGTTCCGCCCATTTCTCCGCCCCATGGCTGACT
AATTTTTTTTTATTTATGCAGAGGCCGAGGCCGCCTCGGCCTCTGAGCTATTC
CAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAAGCTT:
3' (SEQ ID NO:10)

25 Next, replace the SV40 minimal promoter element present in the pSEAP2- promoter plasmid (Clontech) with this NF- κ B/SV40 fragment using XhoI and HindIII. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

In order to generate stable mammalian cell lines, the NF- κ B/SV40/SEAP
30 cassette is removed from the above NF- κ B/SEAP vector using restriction enzymes SalI and NotI, and inserted into a vector containing neomycin resistance. Particularly, the NF- κ B/SV40/SEAP cassette was inserted into pGFP-1 (Clontech), replacing the GFP gene, after restricting pGFP-1 with SalI and NotI.

Once NF- κ B/SV40/SEAP/Neo vector is created, stable Jurkat T-cells are created and maintained according to the protocol described in Example 13. Similarly, the method for assaying supernatants with these stable Jurkat T-cells is also described in Example 13. As a positive control, exogenous TNF alpha (0.1, 1, 10 ng) is added to wells H9, H10, and H11, with a 5-10 fold activation typically observed.

Example 17: Assay for SEAP Activity

As a reporter molecule for the assays described in Examples 13-16, SEAP activity is assayed using the Tropix Phospho-light Kit (Cat. BP-400) according to the following general procedure. The Tropix Phospho-light Kit supplies the Dilution, Assay, and Reaction Buffers used below.

Prime a dispenser with the 2.5x Dilution Buffer and dispense 15 μ l of 2.5x dilution buffer into Optiplates containing 35 μ l of a supernatant. Seal the plates with a plastic sealer and incubate at 65°C for 30 min. Separate the Optiplates to avoid uneven heating.

Cool the samples to room temperature for 15 minutes. Empty the dispenser and prime with the Assay Buffer. Add 50 μ l Assay Buffer and incubate at room temperature 5 min. Empty the dispenser and prime with the Reaction Buffer (see the table below). Add 50 μ l Reaction Buffer and incubate at room temperature for 20 minutes. Since the intensity of the chemiluminescent signal is time dependent, and it takes about 10 minutes to read 5 plates on luminometer, one should treat 5 plates at each time and start the second set 10 minutes later.

Read the relative light unit in the luminometer. Set H12 as blank, and print the results. An increase in chemiluminescence indicates reporter activity.

Reaction Buffer Formulation:

# of plates	Rxn buffer diluent (ml)	CSPD (ml)
10	60	3
11	65	3.25
12	70	3.5
13	75	3.75
14	80	4
15	85	4.25
16	90	4.5
17	95	4.75
18	100	5
19	105	5.25
20	110	5.5
21	115	5.75
22	120	6

23	125	6.25
24	130	6.5
25	135	6.75
26	140	7
27	145	7.25
28	150	7.5
29	155	7.75
30	160	8
31	165	8.25
32	170	8.5
33	175	8.75
34	180	9
35	185	9.25
36	190	9.5
37	195	9.75
38	200	10
39	205	10.25
40	210	10.5
41	215	10.75
42	220	11
43	225	11.25
44	230	11.5
45	235	11.75
46	240	12
47	245	12.25
48	250	12.5
49	255	12.75
50	260	13

Example 18: High-Throughput Screening Assay Identifying Changes in Small Molecule Concentration and Membrane Permeability

Binding of a ligand to a receptor is known to alter intracellular levels of small molecules, such as calcium, potassium, sodium, and pH, as well as alter membrane potential. These alterations can be measured in an assay to identify supernatants which bind to receptors of a particular cell. Although the following protocol describes an assay for calcium, this protocol can easily be modified to detect changes in potassium, sodium, pH, membrane potential, or any other small molecule which is detectable by a fluorescent probe.

The following assay uses Fluorometric Imaging Plate Reader ("FLIPR") to measure changes in fluorescent molecules (Molecular Probes) that bind small molecules. Clearly, any fluorescent molecule detecting a small molecule can be used instead of the calcium fluorescent molecule, fluo-3, used here.

For adherent cells, seed the cells at 10,000 -20,000 cells/well in a Co-star black 96-well plate with clear bottom. The plate is incubated in a CO₂ incubator for 20 hours. The adherent cells are washed two times in Biotek washer with 200 ul of HBSS (Hank's Balanced Salt Solution) leaving 100 ul of buffer after the final wash.

A stock solution of 1 mg/ml fluo-3 is made in 10% pluronic acid DMSO. To load the cells with fluo-3, 50 ul of 12 ug/ml fluo-3 is added to each well. The plate is incubated at 37°C in a CO₂ incubator for 60 min. The plate is washed four times in the Biotek washer with HBSS leaving 100 ul of buffer.

- 5 For non-adherent cells, the cells are spun down from culture media. Cells are re-suspended to 2-5x10⁶ cells/ml with HBSS in a 50-ml conical tube. 4 ul of 1 mg/ml fluo-3 solution in 10% pluronic acid DMSO is added to each ml of cell suspension. The tube is then placed in a 37°C water bath for 30-60 min. The cells are washed twice with HBSS, resuspended to 1x10⁶ cells/ml, and dispensed into a microplate, 100
10 ul/well. The plate is centrifuged at 1000 rpm for 5 min. The plate is then washed once in Denley CellWash with 200 ul, followed by an aspiration step to 100 ul final volume.

For a non-cell based assay, each well contains a fluorescent molecule, such as fluo-3. The supernatant is added to the well, and a change in fluorescence is detected.

- To measure the fluorescence of intracellular calcium, the FLIPR is set for the
15 following parameters: (1) System gain is 300-800 mW; (2) Exposure time is 0.4 second; (3) Camera F/stop is F/2; (4) Excitation is 488 nm; (5) Emission is 530 nm; and (6) Sample addition is 50 ul. Increased emission at 530 nm indicates an extracellular signaling event which has resulted in an increase in the intracellular Ca⁺⁺ concentration.

20

Example 19: High-Throughput Screening Assay Identifying Tyrosine Kinase Activity

- The Protein Tyrosine Kinases (PTK) represent a diverse group of transmembrane and cytoplasmic kinases. Within the Receptor Protein Tyrosine Kinase
25 RPTK) group are receptors for a range of mitogenic and metabolic growth factors including the PDGF, FGF, EGF, NGF, HGF and Insulin receptor subfamilies. In addition there are a large family of RPTKs for which the corresponding ligand is unknown. Ligands for RPTKs include mainly secreted small proteins, but also membrane-bound and extracellular matrix proteins.

- 30 Activation of RPTK by ligands involves ligand-mediated receptor dimerization, resulting in transphosphorylation of the receptor subunits and activation of the cytoplasmic tyrosine kinases. The cytoplasmic tyrosine kinases include receptor associated tyrosine kinases of the src-family (e.g., src, yes, lck, lyn, fyn) and non-receptor linked and cytosolic protein tyrosine kinases, such as the Jak family, members
35 of which mediate signal transduction triggered by the cytokine superfamily of receptors (e.g., the Interleukins, Interferons, GM-CSF, and Leptin).

Because of the wide range of known factors capable of stimulating tyrosine kinase activity, the identification of novel human secreted proteins capable of activating tyrosine kinase signal transduction pathways are of interest. Therefore, the following protocol is designed to identify those novel human secreted proteins capable of activating the tyrosine kinase signal transduction pathways.

Seed target cells (e.g., primary keratinocytes) at a density of approximately 25,000 cells per well in a 96 well Loprodyne Silent Screen Plates purchased from Nalge Nunc (Naperville, IL). The plates are sterilized with two 30 minute rinses with 100% ethanol, rinsed with water and dried overnight. Some plates are coated for 2 hr with 100 µl of cell culture grade type I collagen (50 mg/ml), gelatin (2%) or polylysine (50 mg/ml), all of which can be purchased from Sigma Chemicals (St. Louis, MO) or 10% Matrigel purchased from Becton Dickinson (Bedford, MA), or calf serum, rinsed with PBS and stored at 4°C. Cell growth on these plates is assayed by seeding 5,000 cells/well in growth medium and indirect quantitation of cell number through use of alamarBlue as described by the manufacturer Alamar Biosciences, Inc. (Sacramento, CA) after 48 hr. Falcon plate covers #3071 from Becton Dickinson (Bedford, MA) are used to cover the Loprodyne Silent Screen Plates. Falcon Microtest III cell culture plates can also be used in some proliferation experiments.

To prepare extracts, A431 cells are seeded onto the nylon membranes of Loprodyne plates (20,000/200µl/well) and cultured overnight in complete medium. Cells are quiesced by incubation in serum-free basal medium for 24 hr. After 5-20 minutes treatment with EGF (60ng/ml) or 50 µl of the supernatant produced in Example 11, the medium was removed and 100 µl of extraction buffer ((20 mM HEPES pH 7.5, 0.15 M NaCl, 1% Triton X-100, 0.1% SDS, 2 mM Na₃VO₄, 2 mM Na₄P₂O₇ and a cocktail of protease inhibitors (# 1836170) obtained from Boehringer Mannheim (Indianapolis, IN) is added to each well and the plate is shaken on a rotating shaker for 5 minutes at 4°C. The plate is then placed in a vacuum transfer manifold and the extract filtered through the 0.45 µm membrane bottoms of each well using house vacuum. Extracts are collected in a 96-well catch/assay plate in the bottom of the vacuum manifold and immediately placed on ice. To obtain extracts clarified by centrifugation, the content of each well, after detergent solubilization for 5 minutes, is removed and centrifuged for 15 minutes at 4°C at 16,000 x g.

Test the filtered extracts for levels of tyrosine kinase activity. Although many methods of detecting tyrosine kinase activity are known, one method is described here.

Generally, the tyrosine kinase activity of a supernatant is evaluated by determining its ability to phosphorylate a tyrosine residue on a specific substrate (a

biotinylated peptide). Biotinylated peptides that can be used for this purpose include PSK1 (corresponding to amino acids 6-20 of the cell division kinase cdc2-p34) and PSK2 (corresponding to amino acids 1-17 of gastrin). Both peptides are substrates for a range of tyrosine kinases and are available from Boehringer Mannheim.

- 5 The tyrosine kinase reaction is set up by adding the following components in order. First, add 10ul of 5uM Biotinylated Peptide, then 10ul ATP/Mg₂₊ (5mM ATP/50mM MgCl₂), then 10ul of 5x Assay Buffer (40mM imidazole hydrochloride, pH7.3, 40 mM beta-glycerophosphate, 1mM EGTA, 100mM MgCl₂, 5 mM MnCl₂, 0.5 mg/ml BSA), then 5ul of Sodium Vanadate(1mM), and then 5ul of water. Mix the
- 10 components gently and preincubate the reaction mix at 30°C for 2 min. Initiate the reaction by adding 10ul of the control enzyme or the filtered supernatant.

The tyrosine kinase assay reaction is then terminated by adding 10 ul of 120mM EDTA and place the reactions on ice.

- 15 Tyrosine kinase activity is determined by transferring 50 ul aliquot of reaction mixture to a microtiter plate (MTP) module and incubating at 37°C for 20 min. This allows the streptavidin coated 96 well plate to associate with the biotinylated peptide. Wash the MTP module with 300ul/well of PBS four times. Next add 75 ul of anti-phosphotyrosine antibody conjugated to horse radish peroxidase(anti-P-Tyr-POD(0.5u/ml)) to each well and incubate at 37°C for one hour. Wash the well as
- 20 above.

- Next add 100ul of peroxidase substrate solution (Boehringer Mannheim) and incubate at room temperature for at least 5 mins (up to 30 min). Measure the absorbance of the sample at 405 nm by using ELISA reader. The level of bound peroxidase activity is quantitated using an ELISA reader and reflects the level of
- 25 tyrosine kinase activity.

Example 20: High-Throughput Screening Assay Identifying Phosphorylation Activity

- 30 As a potential alternative and/or complement to the assay of protein tyrosine kinase activity described in Example 19, an assay which detects activation (phosphorylation) of major intracellular signal transduction intermediates can also be used. For example, as described below one particular assay can detect tyrosine phosphorylation of the Erk-1 and Erk-2 kinases. However, phosphorylation of other molecules, such as Raf, JNK, p38 MAP, Map kinase kinase (MEK), MEK kinase,
- 35 Src, Muscle specific kinase (MuSK), IRAK, Tec, and Janus, as well as any other

phosphoserine, phosphotyrosine, or phosphothreonine molecule, can be detected by substituting these molecules for Erk-1 or Erk-2 in the following assay.

Specifically, assay plates are made by coating the wells of a 96-well ELISA plate with 0.1ml of protein G (1ug/ml) for 2 hr at room temp, (RT). The plates are then
5 rinsed with PBS and blocked with 3% BSA/PBS for 1 hr at RT. The protein G plates are then treated with 2 commercial monoclonal antibodies (100ng/well) against Erk-1 and Erk-2 (1 hr at RT) (Santa Cruz Biotechnology). (To detect other molecules, this step can easily be modified by substituting a monoclonal antibody detecting any of the above described molecules.) After 3-5 rinses with PBS, the plates are stored at 4°C
10 until use.

A431 cells are seeded at 20,000/well in a 96-well Loprodyne filterplate and cultured overnight in growth medium. The cells are then starved for 48 hr in basal medium (DMEM) and then treated with EGF (6ng/well) or 50 ul of the supernatants obtained in Example 11 for 5-20 minutes. The cells are then solubilized and extracts
15 filtered directly into the assay plate.

After incubation with the extract for 1 hr at RT, the wells are again rinsed. As a positive control, a commercial preparation of MAP kinase (10ng/well) is used in place of A431 extract. Plates are then treated with a commercial polyclonal (rabbit) antibody (1ug/ml) which specifically recognizes the phosphorylated epitope of the Erk-1 and
20 Erk-2 kinases (1 hr at RT). This antibody is biotinylated by standard procedures. The bound polyclonal antibody is then quantitated by successive incubations with Europium-streptavidin and Europium fluorescence enhancing reagent in the Wallac DELFIA instrument (time-resolved fluorescence). An increased fluorescent signal over background indicates a phosphorylation.

25

Example 21: Method of Determining Alterations in a Gene Corresponding to a Polynucleotide

RNA isolated from entire families or individual patients presenting with a phenotype of interest (such as a disease) is be isolated. cDNA is then generated from
30 these RNA samples using protocols known in the art. (See, Sambrook.) The cDNA is then used as a template for PCR, employing primers surrounding regions of interest in SEQ ID NO:X. Suggested PCR conditions consist of 35 cycles at 95°C for 30 seconds; 60-120 seconds at 52-58°C; and 60-120 seconds at 70°C, using buffer solutions described in Sidransky, D., et al., Science 252:706 (1991).

35 PCR products are then sequenced using primers labeled at their 5' end with T4 polynucleotide kinase, employing SequiTherm Polymerase. (Epicentre Technologies).

The intron-exon borders of selected exons is also determined and genomic PCR products analyzed to confirm the results. PCR products harboring suspected mutations is then cloned and sequenced to validate the results of the direct sequencing.

5 PCR products is cloned into T-tailed vectors as described in Holton, T.A. and Graham, M.W., Nucleic Acids Research, 19:1156 (1991) and sequenced with T7 polymerase (United States Biochemical). Affected individuals are identified by mutations not present in unaffected individuals.

10 Genomic rearrangements are also observed as a method of determining alterations in a gene corresponding to a polynucleotide. Genomic clones isolated according to Example 2 are nick-translated with digoxigenindeoxy-uridine 5'-triphosphate (Boehringer Mannheim), and FISH performed as described in Johnson, Cg. et al., Methods Cell Biol. 35:73-99 (1991). Hybridization with the labeled probe is carried out using a vast excess of human cot-1 DNA for specific hybridization to the corresponding genomic locus.

15 Chromosomes are counterstained with 4,6-diamino-2-phenylidole and propidium iodide, producing a combination of C- and R-bands. Aligned images for precise mapping are obtained using a triple-band filter set (Chroma Technology, Brattleboro, VT) in combination with a cooled charge-coupled device camera (Photometrics, Tucson, AZ) and variable excitation wavelength filters. (Johnson, Cv. et al., Genet. Anal. Tech. Appl., 8:75 (1991).) Image collection, analysis and chromosomal fractional length measurements are performed using the ISee Graphical Program System. (Inovision Corporation, Durham, NC.) Chromosome alterations of the genomic region hybridized by the probe are identified as insertions, deletions, and translocations. These alterations are used as a diagnostic marker for an associated
25 disease.

Example 22: Method of Detecting Abnormal Levels of a Polypeptide in a Biological Sample

30 A polypeptide of the present invention can be detected in a biological sample, and if an increased or decreased level of the polypeptide is detected, this polypeptide is a marker for a particular phenotype. Methods of detection are numerous, and thus, it is understood that one skilled in the art can modify the following assay to fit their particular needs.

35 For example, antibody-sandwich ELISAs are used to detect polypeptides in a sample, preferably a biological sample. Wells of a microtiter plate are coated with specific antibodies, at a final concentration of 0.2 to 10 ug/ml. The antibodies are either monoclonal or polyclonal and are produced by the method described in Example 10.

The wells are blocked so that non-specific binding of the polypeptide to the well is reduced.

The coated wells are then incubated for > 2 hours at RT with a sample containing the polypeptide. Preferably, serial dilutions of the sample should be used to validate results. The plates are then washed three times with deionized or distilled water to remove unbounded polypeptide.

Next, 50 ul of specific antibody-alkaline phosphatase conjugate, at a concentration of 25-400 ng, is added and incubated for 2 hours at room temperature. The plates are again washed three times with deionized or distilled water to remove unbounded conjugate.

Add 75 ul of 4-methylumbelliferyl phosphate (MUP) or p-nitrophenyl phosphate (NPP) substrate solution to each well and incubate 1 hour at room temperature. Measure the reaction by a microtiter plate reader. Prepare a standard curve, using serial dilutions of a control sample, and plot polypeptide concentration on the X-axis (log scale) and fluorescence or absorbance of the Y-axis (linear scale). Interpolate the concentration of the polypeptide in the sample using the standard curve.

Example 23: Formulating a Polypeptide

The secreted polypeptide composition will be formulated and dosed in a fashion consistent with good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with the secreted polypeptide alone), the site of delivery, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" for purposes herein is thus determined by such considerations.

As a general proposition, the total pharmaceutically effective amount of secreted polypeptide administered parenterally per dose will be in the range of about 1 µg/kg/day to 10 mg/kg/day of patient body weight, although, as noted above, this will be subject to therapeutic discretion. More preferably, this dose is at least 0.01 mg/kg/day, and most preferably for humans between about 0.01 and 1 mg/kg/day for the hormone. If given continuously, the secreted polypeptide is typically administered at a dose rate of about 1 µg/kg/hour to about 50 µg/kg/hour, either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears to vary depending on the desired effect.

Pharmaceutical compositions containing the secreted protein of the invention are administered orally, rectally, parenterally, intracisternally, intravaginally,

intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

The secreted polypeptide is also suitably administered by sustained-release systems. Suitable examples of sustained-release compositions include semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules. Sustained-release matrices include polylactides (U.S. Pat. No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman, U. et al., *Biopolymers* 22:547-556 (1983)), poly (2-hydroxyethyl methacrylate) (R. Langer et al., *J. Biomed. Mater. Res.* 15:167-277 (1981), and R. Langer, *Chem. Tech.* 12:98-105 (1982)), ethylene vinyl acetate (R. Langer et al.) or poly-D-(-)-3-hydroxybutyric acid (EP 133,988). Sustained-release compositions also include liposomally entrapped polypeptides. Liposomes containing the secreted polypeptide are prepared by methods known per se: DE 3,218,121; Epstein et al., *Proc. Natl. Acad. Sci. USA* 82:3688-3692 (1985); Hwang et al., *Proc. Natl. Acad. Sci. USA* 77:4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U.S. Pat. Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the optimal secreted polypeptide therapy.

For parenteral administration, in one embodiment, the secreted polypeptide is formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation. For example, the formulation preferably does not include oxidizing agents and other compounds that are known to be deleterious to polypeptides.

Generally, the formulations are prepared by contacting the polypeptide uniformly and intimately with liquid carriers or finely divided solid carriers or both. Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten residues) polypeptides, e.g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, manose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic surfactants such as polysorbates, poloxamers, or PEG.

The secreted polypeptide is typically formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml, preferably 1-10 mg/ml, at a pH of about 3 to 8. It will be understood that the use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of polypeptide salts.

Any polypeptide to be used for therapeutic administration can be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (e.g., 0.2 micron membranes). Therapeutic polypeptide compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

Polypeptides ordinarily will be stored in unit or multi-dose containers, for example, sealed ampoules or vials, as an aqueous solution or as a lyophilized formulation for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1% (w/v) aqueous polypeptide solution, and the resulting mixture is lyophilized. The infusion solution is prepared by reconstituting the lyophilized polypeptide using bacteriostatic Water-for-Injection.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, the polypeptides of the present invention may be employed in conjunction with other therapeutic compounds.

Example 24: Method of Treating Decreased Levels of the Polypeptide

It will be appreciated that conditions caused by a decrease in the standard or normal expression level of a secreted protein in an individual can be treated by administering the polypeptide of the present invention, preferably in the secreted form.

- 5 Thus, the invention also provides a method of treatment of an individual in need of an increased level of the polypeptide comprising administering to such an individual a pharmaceutical composition comprising an amount of the polypeptide to increase the activity level of the polypeptide in such an individual.

- 10 For example, a patient with decreased levels of a polypeptide receives a daily dose 0.1-100 ug/kg of the polypeptide for six consecutive days. Preferably, the polypeptide is in the secreted form. The exact details of the dosing scheme, based on administration and formulation, are provided in Example 23.

Example 25: Method of Treating Increased Levels of the Polypeptide

- 15 Antisense technology is used to inhibit production of a polypeptide of the present invention. This technology is one example of a method of decreasing levels of a polypeptide, preferably a secreted form, due to a variety of etiologies, such as cancer.

- For example, a patient diagnosed with abnormally increased levels of a polypeptide is administered intravenously antisense polynucleotides at 0.5, 1.0, 1.5, 20 2.0 and 3.0 mg/kg day for 21 days. This treatment is repeated after a 7-day rest period if the treatment was well tolerated. The formulation of the antisense polynucleotide is provided in Example 23.

Example 26: Method of Treatment Using Gene Therapy

- 25 One method of gene therapy transplants fibroblasts, which are capable of expressing a polypeptide, onto a patient. Generally, fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The flask is 30 turned upside down, closed tight and left at room temperature over night. After 24 hours at room temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e.g., Ham's F12 media, with 10% FBS, penicillin and streptomycin) is added. The flasks are then incubated at 37°C for approximately one week.

At this time, fresh media is added and subsequently changed every several days. After an additional two weeks in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks.

5 pMV-7 (Kirschmeier, P.T. et al., DNA, 7:219-25 (1988)), flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf intestinal phosphatase. The linear vector is fractionated on agarose gel and purified, using glass beads.

10 The cDNA encoding a polypeptide of the present invention can be amplified using PCR primers which correspond to the 5' and 3' end sequences respectively as set forth in Example 1. Preferably, the 5' primer contains an EcoRI site and the 3' primer includes a HindIII site. Equal quantities of the Moloney murine sarcoma virus linear backbone and the amplified EcoRI and HindIII fragment are added together, in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is then used to
15 transform bacteria HB101, which are then plated onto agar containing kanamycin for the purpose of confirming that the vector has the gene of interest properly inserted.

The amphotropic pA317 or GP+am12 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the gene is
20 then added to the media and the packaging cells transduced with the vector. The packaging cells now produce infectious viral particles containing the gene (the packaging cells are now referred to as producer cells).

Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media,
25 containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is removed from a sub-confluent plate of fibroblasts and quickly replaced with the media from the producer cells. This media is removed and replaced with fresh media. If the titer of virus is high, then virtually all fibroblasts will be infected and no selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a
30 selectable marker, such as neo or his. Once the fibroblasts have been efficiently infected, the fibroblasts are analyzed to determine whether protein is produced.

The engineered fibroblasts are then transplanted onto the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads.

Example 27: Method of Treatment Using Gene Therapy - In Vivo

Another aspect of the present invention is using *in vivo* gene therapy methods to treat disorders, diseases and conditions. The gene therapy method relates to the introduction of naked nucleic acid (DNA, RNA, and antisense DNA or RNA) sequences into an animal to increase or decrease the expression of the polypeptide of the present invention. A polynucleotide of the present invention may be operatively linked to a promoter or any other genetic elements necessary for the expression of the encoded polypeptide by the target tissue. Such gene therapy and delivery techniques and methods are known in the art, see, for example, WO90/11092, WO98/11779; U.S. Patent NO. 5693622, 5705151, 5580859; Tabata H. et al. (1997) Cardiovasc. Res. 35(3):470-479, Chao J et al. (1997) Pharmacol. Res. 35(6):517-522, Wolff J.A. (1997) Neuromuscul. Disord. 7(5):314-318, Schwartz B. et al. (1996) Gene Ther. 3(5):405-411, Tsurumi Y. et al. (1996) Circulation 94(12):3281-3290 (incorporated herein by reference).

The polynucleotide constructs of the present invention may be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, intestine and the like). These polynucleotide constructs can be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

The term "naked" polynucleotide, DNA or RNA, refers to sequences that are free from any delivery vehicle that acts to assist, promote, or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the polynucleotides may also be delivered in liposome formulations (such as those taught in Felgner P.L. et al. (1995) Ann. NY Acad. Sci. 772:126-139 and Abdallah B. et al. (1995) Biol. Cell 85(1):1-7) which can be prepared by methods well known to those skilled in the art.

The polynucleotide vector constructs of the present invention used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Any strong promoter known to those skilled in the art can be used for driving the expression of DNA. Unlike other gene therapies techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

The polynucleotide construct of the present invention can be delivered to the interstitial space of tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

For the naked polynucleotide injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 g/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration. The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked polynucleotide constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

The dose response effects of injected polynucleotide in muscle *in vivo* is determined as follows. Suitable template DNA for production of mRNA coding for the polypeptide of the present invention is prepared in accordance with a standard recombinant DNA methodology. The template DNA, which may be either circular or linear, is either used as naked DNA or complexed with

liposomes. The quadriceps muscles of mice are then injected with various amounts of the template DNA.

Five to six week old female and male Balb/C mice are anesthetized by intraperitoneal injection with 0.3 ml of 2.5% Avertin. A 1.5 cm incision is made on the anterior thigh, and the quadriceps muscle is directly visualized. The template DNA is injected in 0.1 ml of carrier in a 1 cc syringe through a 27 gauge needle over one minute, approximately 0.5 cm from the distal insertion site of the muscle into the knee and about 0.2 cm deep. A suture is placed over the injection site for future localization, and the skin is closed with stainless steel clips.

After an appropriate incubation time (e.g., 7 days) muscle extracts are prepared by excising the entire quadriceps. Every fifth 15 um cross-section of the individual quadriceps muscles is histochemically stained for protein expression. A time course for protein expression may be done in a similar fashion except that quadriceps from different mice are harvested at different times. Persistence of DNA in muscle following injection may be determined by Southern blot analysis after preparing total cellular DNA and HIRT supernatants from injected and control mice. The results of the above experimentation in mice can be use to extrapolate proper dosages and other treatment parameters in humans and other animals using naked DNA of the present invention.

It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples. Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

The entire disclosure of each document cited (including patents, patent applications, journal articles, abstracts, laboratory manuals, books, or other disclosures) in the Background of the Invention, Detailed Description, and Examples is hereby incorporated herein by reference.

Sequence Listing

(1) GENERAL INFORMATION:

- 5 (i) APPLICANT: Rosen et al.
- (ii) TITLE OF INVENTION: 86 Human Secreted Proteins
- 10 (iii) NUMBER OF SEQUENCES: 318
- (iv) CORRESPONDENCE ADDRESS:
- 15 (A) ADDRESSEE: Human Genome Sciences, Inc.
- (B) STREET: 9410 Key West Avenue
- (C) CITY: Rockville
- 20 (D) STATE: Maryland
- (E) COUNTRY: USA
- 25 (F) ZIP: 20850
- (v) COMPUTER READABLE FORM:
- 30 (A) MEDIUM TYPE: Diskette, 3.50 inch, 1.4Mb storage
- (B) COMPUTER: HP Vectra 486/33
- 35 (C) OPERATING SYSTEM: MSDOS version 6.2
- (D) SOFTWARE: ASCII Text
- 40 (vi) CURRENT APPLICATION DATA:
- (A) APPLICATION NUMBER:
- 45 (B) FILING DATE: June 11, 1998
- (C) CLASSIFICATION:
- 50 (vii) PRIOR APPLICATION DATA:
- (A) APPLICATION NUMBER:
- 55 (B) FILING DATE:

(viii) ATTORNEY/AGENT INFORMATION:

- 5 (A) NAME: A. Anders Brookes
(B) REGISTRATION NUMBER: 36,373
10 (C) REFERENCE/DOCKET NUMBER: PZ008PCT

(vi) TELECOMMUNICATION INFORMATION:

- 15 (A) TELEPHONE: (301) 309-8504
(B) TELEFAX: (301) 309-8439

20

(2) INFORMATION FOR SEQ ID NO: 1:

- 25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 733 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

GGGATCCGGA GCCCAAATCT TCTGACAAAA CTCACACATG CCCACCGTGC CCAGCACCTG 60
AATTCGAGGG TGCACCGTCA GTCTTCCTCT TCCCCCAAAA ACCCAAGGAC ACCCTCATGA 120
35 TCTCCCGGAC TCCTGAGGTC ACATGCGTGG TGGTGGACGT AAGCCACGAA GACCCTGAGG 180
TCAAGTTCAA CTGGTACGTG GACGGCGTGG AGGTGCATAA TGCCAAGACA AAGCCGCGGG 240
40 AGGAGCAGTA CAACAGCACG TACCGTGTGG TCAGCGTCCT CACCGTCCTG CACCAGGACT 300
GGCTGAATGG CAAGGAGTAC AAGTGCAAGG TCTCCAACAA AGCCCTCCCA ACCCCCATCG 360
AGAAAACCAT CTCCAAGCC AAAGGCGAGC CCCGAGAACC ACAGGTGTAC ACCCTGCCCC 420
45 CATCCCGGGA TGAGCTGACC AAGAACCAGG TCAGCCTGAC CTGCCTGGTC AAAGGCTTCT 480
ATCCAAGCGA CATCGCCGTG GAGTGGGAGA GCAATGGGCA GCCGGAGAAC AACTACAAGA 540
50 CCACGCCTCC CGTGCTGGAC TCCGACGGCT CCTTCTTCCT CTACAGCAAG CTCACCGTGG 600
ACAAGAGCAG GTGGCAGCAG GGGAACGTCT TCTCATGCTC CGTGATGCAT GAGGCTCTGC 660
ACAACCACTA CACGAGAAG AGCCTCTCCC TGTCTCCGGG TAAATGAGTG CGACGGCCGC 720
55 GACTCTAGAG GAT 733

(2) INFORMATION FOR SEQ ID NO: 2:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 5 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Trp Ser Xaa Trp Ser
1 5

15

(2) INFORMATION FOR SEQ ID NO: 3:

- 20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 86 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
25 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GCGCCTCGAG ATTTCCCGGA AATCTAGATT TCCCGAAAT GATTTCCTCG AAATGATTTT 60

30 CCCGAAATAT CTGCCATCTC AATTAG 86

35 (2) INFORMATION FOR SEQ ID NO: 4:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 27 base pairs
(B) TYPE: nucleic acid
40 (C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

45 GCGCAAGCT TTTTGCAAAG CCTAGGC 27

50 (2) INFORMATION FOR SEQ ID NO: 5:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 271 base pairs
(B) TYPE: nucleic acid
55 (C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

60 CTCGAGATTT CCCCGAAATC TAGATTTCCC CGAAATGATT TCCCGAAAT GATTTCCTCG 60

10
5
15
20
25
30
35
40
45
50
55
60

AAATATCTGC CATCTCAATT AGTCAGCAAC CATAGTCCCG CCCCTAACTC CGCCCATCCC 120
GCCCCTAACT CCGCCCAGTT CGCCCCATTC TCCGCCCAT GGCTGACTAA TTTT TTTTAT 180
TTATGCAGAG GCCGAGGCCG CCTCGGCCTC TGAGCTATTC CAGAAGTAGT GAGGAGGCTT 240
TTTTGGAGGC CTAGGCTTTT GCAAAAAGCT T 271

(2) INFORMATION FOR SEQ ID NO: 6:

15
20
25

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 32 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

GCGCTCGAGG GATGACAGCG ATAGAACCCC GG 32

(2) INFORMATION FOR SEQ ID NO: 7:

30
35
40

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 31 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

GCGAAGCTTC GCGACTCCCC GGATCCGCCT C 31

(2) INFORMATION FOR SEQ ID NO: 8:

45
50
55

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 12 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

GGGGACTTTC CC 12

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 73 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

5 GCGGCCTCGA GGGGACTTTC CCGGGGACTT TCCGGGGACT TTCCGGGACT TTCCATCCTG 60
10 CCATCTCAAT TAG 73

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 256 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

15 CTCGAGGGGA CTTTCCCGGG GACTTTCCGG GGACTTTCCG GGACTTTCCA TCTGCCATCT 60
20 CAATTAGTCA GCAACCATAG TCCCGCCCCT AACTCCGCCC ATCCCCGCCC TAACTCCGCC 120
30 CAGTCCGCC CATTCCTCCG CCCATGGCTG ACTAATTTT TTTATTTATG CAGAGGCCGA 180
GGCCGCTCG GCCTCTGAGC TATTCCAGAA GTAGTGAGGA GGCTTTTTTG GAGGCCTAGG 240
35 CTTTTCAGAA AAGCTT 256

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1220 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

50 CATGAATGGC TCGCACAAGG ACCCCCTCCT CCCCTTTCCT GCTTCTGGA GAACTCCCTC 60
CCTCCCTCCA GCTCCGCCAG CCCAGGCGCC CCTTCCCTGG AAGCCGAGCG GCTTCGCTCG 120
CATTTACCG CCGCCGCTC TCGCAATATT GCAATATAGG GGAAAAGCAG ACCATGGTGA 180
55 ATCCGGGAG CAGCTCGCAG CCGCCCCCGG TGACGGCCGG CTCCCTCTCC TGGAAGCGGT 240
GCGCAGGCTG CGGGGGCAAG ATTGCGGACC GCTTCTGCT CTATGCCATG GACAGCTATT 300
60 GGCACAGCCG GTGCCTCAAG TGCTCCTGCT GCCAGGCGCA NTGGGCGACA TCGGCACGTC 360

	CTGTTACACC AAAAGTGGCA TGATCCTTTG CAGAAATGAC TACATTAGGT TATTTGGAAA	420
	TAGCGGTGCT TGCAGCGCTT GCGGACAGTC GATTCCCTGCG AGTGAACCTG TCATGAGGGC	480
5	GCAAGGCAAT GTGTATCATC TTAAGTGTIT TACATGCTCT ACCTGCCCGA ATCGCCTGGT	540
	CCCGGGAGAT CGGTTTCACT ACATCAATGG CAGTTTATTT TGTGAACATG ATAGACCTAC	600
10	AGCTCTCATC AATGGCCAAT TGAATTCAC TCAAGCAAT CCACTACTGC CAGACCAGAA	660
	GGTCTGCTAA AAGGTCAGAG TAATGCAGAA TCGGTGCCTT CATCTCAGAT TTGTTTCATCA	720
	CAGGTGGATC CCATGTTTCT TCAGTAGACA AGTCACCTTT GTAGCTAGCA CCAGTGCCAG	780
15	CTCCATGCCA TTGCACCTTC TTTAGTCTTG ATTGCCCTTC CCGCATTTWT TGGTGTATTA	840
	AAATGACTRA TKAAGCTAAT TAAAGAAGC ATTCAAATCT GCTTCTTACC CTCATTAACA	900
20	ATTAGCAGGG CACTGGCCAG AGTTTGTACC CTGTGTTTTC CCTTAACAAC ATTCTATTG	960
	CTCTTTGTAT ATTTAAGTGT TGAAGGAAA CGTGTTTCAA TCAAACTGA CCATGAGATA	1020
	AAGGAAAGAG ATGTGGCTTT TGTGATATTC TATCACAAAC ACTTATTGTA TCTCTGTAAA	1080
25	ATACAATGTA TGTATGCATG TAAGTGTITT TGTCCTAATG TTGCTACTCC CATGGCAAAG	1140
	AAAAAAAAA GAATGAAAAA AARAAAAAAA AAAAAAAAAA AAAAAAAAAA CTCGAGGGGG	1200
30	GGCCCCGTACC CAATCGCCCT	1220

(2) INFORMATION FOR SEQ ID NO: 12:

35

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1939 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

40

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

45	GAACACAAAC ATGCAGTCTG TAGCAGATGG TAATAGGCTG AYATATTACA CTTGTTGATG	60
	TAAATCTGAT AGGTTTCTTT CTCTCCAAGG ACAGCTTTTT AAATATTAA CAGTATCAAT	120
	AATTTTTCAG TTTCTGTGAG AATTTTATAA TTTATAATTT GCAGACTTAA TGTATAATCT	180
50	ATTTTGTCTT AACAAATACA AATATATTTT TTATTTTCAGA TTRTATATAT TCCTACCAGA	240
	TGGAGATAAT TACAGCTTTA AAAATTTTTA TTTTTTCATT TTATTTTACA CATTGACATT	300
55	AAATTTTTAT GGACACATAA TAACTGTACA TATATATGGG GTAGAAATGT ATGTTTTAAT	360
	ACATGTACTC AATGTGTAAT GATCAAATCA GGGTAAATTT CATAATGATT TTTCTGTAGG	420
	GAGAAAATTC AAAATCTACT CTTCTGGCTA TTTTCAAATA TATAATATGT TATTGTTAAC	480
60	TATACTCATC CTAATATGCA ATAGGACACC AGAACTTATT CCTGGGTTCT ACATCCGTTA	540

	AGGCAACCAA GGATTGAAA TATGGAAAA AAAAATGCG TCTGTACTGA ACATGTACAG	600
5	ACTTTTCT TGTCTTAT CTTCACAA TATAGTACAA TAACTATTG CATGACATTT	660
	ACATCGGATA TTATGAGTGA TCTAGAGTTG ATATGAAGTA TATGGGAGGA TGTGCAAAGG	720
	TGATGTGCAA ATACTATGTC ATTTTATATC AGGGACTTGA GTATCCTTTG TTAYCCTCAG	780
10	GAGATCCTGA AACYAGTCCC CCATGGATAC TGAGGGCTGA CTGTATAGTC CTATCCTCAC	840
	GGAACTTTCA TTCTAATGRG GGAAGACTGA CTATAACAA AATATATGTA ATAGGTGGTG	900
15	GTAAGTACCG TCGAGAAGTA ACAAATGGGG CAAAGTGAGT TATACAGCTC CATYCTTAGA	960
	AACCTTGGAG TACTTTTCTT AGTTTATACT CGTGGTGGTT TCCTTTTGTC TCCTTTATTA	1020
	CATGGGACTC TGACATGTGC CCATAGCTAG GGTGGCAGTA GGATCTACCC GAAAAGCGTC	1080
20	CTGCTGATAC AGGACCAAAG CATCCTGTTG TTCTCGAGCC TATAAAAAGA GCTAATGGTC	1140
	TTGCTTCTCT TAACGTGGC CTCCTACACT GTGTTTGGGA TGATTGGTGA TGTCTTGGAT	1200
25	ATTCTGTTTC TTTGGAACCT TGAATATACA ACACCTTACT AGGGAATTAG CAATGGAAGC	1260
	AGAGCAAAGA TGTACAGAGG AAACAATGCR TAACTCTGAT GGAATTGAAG TCATGAGGCA	1320
	GCAGAGAGCT TAAATTASAG CTTTAAAAAT TTTTATTTT TAGAGGGAAT TTAMTTGGGA	1380
30	GTAACAGCAG TAATAGTTAA CGGAGCCAGA ATGCTTGAGT CATATAATTG CAAAGCAGAG	1440
	TTGGGAGCAA CAGATGCTAA AGAGTAGTTG CTGTAGTTCC TCTTTGGGTC GTAGGAGCAG	1500
35	TTGTCATRTT MCTATAYAGC TACTGCATGA AGAAGAGTTC TTAGTGAGGC CTGGGTGAAC	1560
	AGCTCTTCTT AGTATCTCTG GTGACCCCAT TYGACCTTTT AACAAATCCC TAAGTAAATA	1620
	AATAGCCCCT MAGGWAACT AAGTTTCTCT CTGCTGTTTT TTTGCTTGAG AGAGCTATAA	1680
40	CTGTAATAGA CTTATATTTT TGAACATTTT AGTGCTTGCC AATATTGGT AATATTTATG	1740
	TTTCCTATAT TTGTAATGAA CATCTCTCTT CMGGTACAT TYTTGTAAAA TTATTGTTTS	1800
45	ATGSATAAAA GTTCACCTTT TATTGTATAA AATTGACTCA GATTAATTTA TACACATTGA	1860
	CAATGGGTAA ATAGAGTTTT TCAGATTATT AAAAGCTGAA GGATGCCCAT GTAAGCAAAA	1920
50	AAAAAAAAA AAAACTCGA	1939

(2) INFORMATION FOR SEQ ID NO: 13:

- 55 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 2602 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear
- 60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

	GGGTTCCTCG GGCACCTTTC CTTCCGGGT GTTCTGAAGC GGTTCCTCTG TAATCCTCAG	60
5	TGAGGAAACC CACCGTGAAT CGGATTGCCG TTCAGTCCCA CGGAAGCCTG GCTCGTTGGC	120
	CATGTINGGG ACGCATGTTT ATTAAGTTCA TTAAAATAAT TTCATTTGTC TTGGTTTGAA	180
10	GACTGCTTCA TTCTGCCTCT AGTACCAGCG GTTCTCTGT TCTGTGATCA ATGTGATTCA	240
	CAGGAACTCC TTAAGTAACA AACGAAATGA GCCAGGGGCG TGGAAAATAT GACTTCTATA	300
	TTGGTCTGGG ATTGGCTATG AGCTCCAGCA TTTTCATTGG AGGAAGTTC ATTTTGAAAA	360
15	AAAAGGCGCT CCTTCGACTT GCCAGGAAAG GCTCTATGAG AGCAGGTCAA GGTGGCCATG	420
	CATATCTTAA GGAATGGTTG TGGTGGGCTG GACTGCTGTC AATGGGAGCT GGTGAGGTGG	480
20	CCAACTTCGC TCGGTATGCG TTTGCACCAG CCACTCTAGT GACTCCACTA GGAGCTCTCA	540
	GCGTCTAGT AAGTGCCATT CTTTCTTCAT ACTTTCTCAA TGAAAGACTT AATCTTCATG	600
	GGAAAATTGG GTGTTTGCTA AGTATTCTAG GATCTACAGT TATGGTCATT CATGCTCCAA	660
25	AGGAAGAGGA GATTGAGACT TTAAATGAAA TGTCTCACA GCTAGGTGAT CCAGGTTTGT	720
	TGGTCTTTGC AACCTTGTG GTCATTGTGG CCTTGATATT AATCTTCGTG GTGGTCTCTC	780
30	GCCATGGACA GACAAACATT CTGTGTACA TAACAATCTG CTCTGTAATC GCGCGTTTT	840
	CAGTCTCCTG TGTGAAGGGC CTGGGCATTG CTATCAAGGA GCTGTTTGCA GGAAGCCTG	900
	TGCTGCGGCA TCCCTGGCT TGGATTCTGC TGTGAGCCT CATCGTCTGT GTGAGCACAC	960
35	AGATTAAITA CCTAAATAGG GCCCTGGATA TATTCAACAC TTCCATTGTG ACTCCAATAT	1020
	ATTATGTATT CTTTACAACA TCAGTTTAA CTTGTTGAGC TATCTTTTTT AAGGAGTGGC	1080
40	AAGATATGCC TGTGACGAT GTCATTGGTA CTTGAGTGG CTTCTTTACA ATCATTGTGG	1140
	GGATATTCTT GTTGCATGCC TTAAAGACG TCAGCTTTAG TCTAGCAAGT CTGCCTGIGT	1200
	CTTTTCGAAA AGACGAGAAA GCAATGAATG GCAATCTCTC TAATATGTAT GAAGTTCTTA	1260
45	ATAATAATGA AGAAAGCTTA ACCTGTGGAA TCGAACAACA CACTGGTGAA AATGTCTCCC	1320
	GAAGAAATGG AAATCTGACA GCTTTTAAAG AAAGGTGTAA TTAAAGGTTA ATCTGTGATT	1380
50	GTTATGAAGT GAATTTGAAT ATCATCAGAA TGTGTCTGAA AAAACATTGT CCTCAAATAA	1440
	TGTTCTTTAA AGGCAATCTT TTAAAGATT TCACTAATTT GGACCAAGAA ATTACTTTTC	1500
	TTGTATTTAA ACAACAATG GTAGCTCACT AAAATGACCT CAGCACATGA CGATTTCTAT	1560
55	TAACATTTTA TTGTTGTAGA AGTATTTTAC ATTTTCATCC CTCTCCAAA AGCCGAATGC	1620
	ACTAATGACA GTTTTAAGTC TATGAAAATG CTTTATTTTT TCATTGGTGA TGAAAGCTCG	1680
60	AAATGTGCAT TTGTATCCC CACTCCATCA ATCCCTGACC ATGTAAGGCT TTTTATTTTT	1740

5 AAAAAACAG AGTTATCCCA ATACATTATC CTGTGATTTA CTTTACCTAC AAAAGTGGCT 1800
 CCTGTTTGTT TGATGATGAT TGGTTTTATT TTTGAAATAT TTATTAAGGG AAAACTAAGT 1860
 TACTGAATGA AGGAACCTCT TTCTTACAAA ACAAAAAAAA GGGCAGAAAT CACCCCAAGG 1920
 AACGATTTCCT CAGGTTGAGA TGATCACCGT GAATCCGGCT TCCTCTGAGC ATTCGATGGC 1980
 10 CTTAGCACCT CATCAAGCCA GCACATCCTG CCTGCTGTTG CAGCCTGGCT GGGTTTATTC 2040
 TTCAGTTACC CTAATCCCAT GATGCCTGGA ACCTTGATTA CCGTTTTACA TCAGCTCTTG 2100
 TACTTTTCAG TATATTTTCA TAATGAGTTA TATTGTCATT TAGACTTTGA ACAGCTCTGG 2160
 15 GAAATAGAAG ACTAGGGTTG TTTCTTAAAT TTAGCTCATG TTATAATAAA AAGTTGAAAT 2220
 GAAGTTCTTA TTCTAAAAGT CTGAATGCTT AGAACAACT TAACATGTTT ATAGAATATG 2280
 GTCTCTTTGT ACCAAGTACT TTGCTTAAAG GCTCCTTTGG GCCACTACAT ATTTTGGTTT 2340
 20 CTAGAAAATG TTTGTTTATG AAGAAGTCGA TGGAAAACG CAAACATATG CAGAAAAGGT 2400
 AGAATAATAA AAAAGGTCTA ATGAACTCCA TTCAGCTTTG AACCTATCCA CTCATAACCA 2460
 25 TTGACTGGCC TTTTAAAAA AAGTATTGGG CAGAATTAAA TTCCACCTA GGTGATGGG 2520
 AAGGAAAGTG TTCGCCTGTN CCAGCCTGTG GTTCCTGCCT GGGNGGTTTA CCCAGTGGTG 2580
 GCGCCAGGCC AAGGTCCATT CA 2602
 30

(2) INFORMATION FOR SEQ ID NO: 14:

35

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 808 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

45 ACCCACGCGT CCGGTAAAC AAAGGGAATG ACGATATGGG AAAGAAAATA CATTTGGATG 60
 TTACAGATAT GTGTGTTCTT GGAGCCCAGG GCCAAGCCCT CCCTGGGGGA CTGGATTGG 120
 TGATCTCTCT CCTTGGCCCC AACCTGACAT CTTTCTTGT CTTTCTAGGA ATGTCTGATG 180
 50 GAAATTCCTC CTAACCTGGG GTCATACTCC ATTTCAATTCT CTGGGCTCAN TGAGAAGGAA 240
 AATTTTTTTT TAAGTAATTT ACTGAAAACC CAGATCACAC CATCATAAAT TCAGATAGGT 300
 GCAATTCTGC CCACAATGAA GGCAAAGTGT TACACTAATT TGAAAACAGT TTAGCCTCTT 360
 55 ATTCCCCCAA ACTTCATTCT TGAATTTTGT CATTTTTTGT GGGCAAGCTG TGGGAAAGGG 420
 GCACAAAAGT ATCACTGAAG TATTTTTTCA AAAAAGAAAA AAGGCAGTCT TCCTCTACTA 480
 60 ATGAGAATGC AAAATGTGTA ACAACTGTAA AATGTTTTCA CCCTGCTTTT AGACATAAAG 540

CTTTAAAAAA CTGTGAGGTC TTTTATCACT TCCCCATTGT ATATGTAATA TGGCTCCAGA 600
TAATTACTCT GCCACGGGGA GAAAATCTTC CATAACTCTC CCCTATATAT ATGTATACTC 660
5 CACCACCTTA TCTGTATTATG TCATGGTGGT GGGAGTATTT ATMCCACAGA AACAGGCAAA 720
TGATACAAAC CTGGGCGACA GAGCAAGACT CCACCTTCAA AAAAAAAAAA AAAAAAAAAA 780
10 AAAAAAAAAA AAAAAAAAAA GGGCGGCC 808

15 (2) INFORMATION FOR SEQ ID NO: 15:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 864 base pairs
(B) TYPE: nucleic acid
20 (C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

25 GGGTTTTTTG TTTTGTGTTT TTNAGGGGGG AGGGGGGGTT TCCCTCCTT TCCCCAGAC 60
TTCTCTTTGA ACACAAATGC ATTAGCCTTG TGGCTAGAAM ACCCTCTTCC TACCTCTGTC 120
TCCCTCACT TGTCATATGC TCTGACATGC TAACATTTCT TTTGTTTCATC CCTGTGCCCC 180
30 CCACAGAAAC ATCCCAGAAA AACCGGTCAG TGTTCCTTCC TCCCTGATCC TTAGGTTTCT 240
GAAATAGGGT TCTGTTACAT CCTCTTCGAT AGCCTGTTTA AAATGTTTAG AAGGTCTGGA 300
35 GCTCAAAAAT GCGTTCTTCC ACATTGATAA TTTAGTAAAC TGAGAACATT GACATCACTA 360
CAGGGCAGCA TAAGAGGTTG CTTACATGTG GTAGCAGCTC TGGTTTGATT CAAGTTGCTA 420
CCATGTACAT TGACAGCACA TATACCATAA CCAGCGTGTT GGGTTGAATT GCACTTTCTA 480
40 CCTTTGTATG AGATTTACAG ACTTCCTTC TGGGTTTGTA TCATGACCAG AGGGTACTA 540
TAGGGTTGGT TTATACTGCA ATATAGAGGA TCAGAAGCCA TTTGATTGTTG TAGGTGTGTC 600
45 AGAAGGAGA ATGATGGCAG ACGAACTGCT GGAAGAGGTC AGAAGATAGC CATGCTAAAA 660
TGCAATTATA TCCTCATGTT TATCCCAAAC TAATCTTGGA CTTTCCACT CATTAGCTTT 720
GTTTTGCCCT TGTTCCTT GAAGGTTTAA GTTCAACCAT ATTCTGTCAA CTGTTAGTT 780
50 TCAGTGAAT CTGTATTTC TGGTTCATTA TAACAAATTG TCGCTTAAA AAAAAAAAAA 840
AAAAGGGCG GCCGCTCTAG AGGG 864

55

(2) INFORMATION FOR SEQ ID NO: 16:

60 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2361 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

	GGCACGAGCT CGAGTTTTTT TTTTTTTTTT TTTCTATTTT TGCCAGACTC TTGATACTCT	60
10	TAAACTTGT TTGTGGTCAG CACAACAAGG AACAAAACAA AGCTTTGAAA AAACCTTAAAC	120
	ATGAAAAAAC GCACTGACAT TTTTTTTTAT TTAATATAGC CTGGACTTTA CCTGCGTATG	180
15	CACATGCTCA GAATTGTCTA CTAGGCTGAC TATGTATCAC CTCCTCAGCT TGGATCCAAT	240
	TGTGGATTTA TTACAAACA TCAAATGCCT TCAAGCCAAT CCTTTTGTCT GTATGTTTTG	300
	CAGCCTACTG TAGTAGATAC GCAACAGATA WTGTGGGAAA AAAAGAGATA AGAGGAGGAA	360
20	GCTAATAAGA GACTGTCAAG ATTGTATACC TTCTTGGTTT CTTTAAAGAA TTTGTTCCT	420
	TTCTACTATT ACAGCAAAGC AGCATTTTGT TACTGACTGC CTAAAATCAC TTAATCTCAG	480
25	GTGAACGCAT CACTTGCCAA ACTGTTGGAA TGCTATTTGT GTTTGTGTGC ACTGTTTTTT	540
	TCGTTTGTTC GTTTGTTTAT TTGGTTGGCT TTTTGGAGAG GGAAATTTGG AAACGGGACA	600
	TACACAAAAG TTACACACCC ACATTCCTTT TTTATCATGA CATACAAGAA GAAACTAGCA	660
30	GAGCTAAGAA TGGAGTGAAG AAAGGCAGTA TGGCAGGCAC CAGCAAAGAG TTGAGGGCTG	720
	TTGCTCTTAA AAATTATTTT TTTTATTATT ATTTTGAAAG TATGGAAGTT TTCCATTCAC	780
35	TGGGGAAGG AGGGAAAAGT GCATTTATTT TTATACAGAG TTAATTAAAT ACCTCCAAAA	840
	CACATATGTT GGAAATCGCT TTGCTGGTG CAAAGTATAT TAATGAGCAG GAATACATAC	900
	ATTGAGGTTA TGAATAGAGA GCTCAATTTG TACCTTTGCT GTCTTGCTCA AGCTTGGTAT	960
40	GGCATGAAAA CTCGACTTTA TTCCAAAAGT AACTTCAAAA TTTAAATAC TAGAACGTTT	1020
	GCTGCGATAA ATCTTTTGA TTTTGTGTT TTTCTAATGA GAATACTGTT TTTCAATTACC	1080
45	TAAAGAACAA TTTGCTAAAC ATGAGAAATC ACTCACTTGT ATTATGTATA GATTACATAG	1140
	GAAGAACAAT CACATCAGTA AGTTATAGTT TATATTAAAG GTAATTTTCT GTTGGCTCAT	1200
	AACAAATATA CCAGCATTCA TGATAGCATT TCAGCATTTT CCAAGGTACC AAGTGACTTT	1260
50	ATTTTGTGTT TGTGTGTTT GTTGTATTTT AGAAGGAATT CAGCTCTGAT GTTTTAAAG	1320
	AAAACCAGCA TCTCTGATGT TGCAACATAC GTGTAAAATG GGTGTTACAT CTATCCTGCC	1380
55	ATTTAACCCC ACAGTTAATA AAGTGGCTGA AAATAATAGT AGCTCTGGCT TGGTGCCTGA	1440
	CCTGGTTAAA TACTGTCTTA AAGCTCATAC AAAACAAATA GGCTTTTCCA TAAGTGGCCT	1500
	TTAAGAAAAC ATGGAAGACA ATTCATGTTT GACAAATGCT GACAGGGTGA AGAAAGCCCA	1560
60	GTGTAAAAAT GAATCGCGTT TTAAGTGATT CGGTAAAGA GTTTGGGCTC CCGTAGCAAA	1620

CTAATACTAG ATAATAAGGA AATGGGGGTG AAATATTTT TTATTGTTGA ATCATTTTGT 1680
 5 GAATGTCCCC CTCAAAAAA GCTAATGGAA TATTTGCCAT AAAGGGCATT TGGTGGTTTT 1740
 ATTTTGTGTT GAGGGGGWTT GTCAGAAAAT CCTTTTCTC TCTTACGYCT AACTGACTAG 1800
 GGAACAATTG TTGATATGCA TAGCATGGG AATACTTGTC ATTATATACT CTTACAAATA 1860
 10 ACACATGAAG CAAGAATGAC CAATATTCTG NATAATTGGG CACTGGGATC ACAAATGTG 1920
 ATAAAACTTT AAATGTATAA AACTTTATCA AATAAAGTTT TATTTTCCCC TTAAAAATGT 1980
 15 ATTTCTTTAG AGGCATTACT TTTTAAAAA TATTGGTCAA TTCCTGACAT AAGATGTGAG 2040
 GTTCACAGTT GTATTCCAGT ATTCAAGATA GATTCCTGAT TTTTCAATTA GGAAAAGTAA 2100
 AATCCAAAAT GTTAGCAAAA CAAAGTGCAA TATTAAATGT TTGCTTTATA GATTATATTC 2160
 20 TATGGCTGTT TGTAAATTCT CTTTTTTTCC TTTTATTTT GGTGCTGAAT ATGTCCTTGT 2220
 AGGCTCTGTT TTAAGAAAAC AATATGTGGG AAATGATTTA ATTTTTCCTA TTGCTCTTCC 2280
 25 TTGTGGAAAA TAAAGTGTTT TGTTTTTC TGTTTTGTA AAAAAAAAAA AAAAAAAAAA 2340
 AAAAAAAAAA AAGAANGAGA A 2361

30

(2) INFORMATION FOR SEQ ID NO: 17:

(i) SEQUENCE CHARACTERISTICS:
 35 (A) LENGTH: 803 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

40

CAGCTGCCCA CAAGGTGGGC TCCTGGGGGA GGGTCATCCC TCTGAGAAGA GGGCGGCACC 60
 AAGACCCACA CACCTGAAAA ATGTGGTACT TCATGTGCT GATCTCGATG GTCTTGCTGC 120
 45 TGTCCCATC CTGTTCTGAT TTATTGGTCA TTAGTGTCTT GAACCTGGAG CAAAGGAGAC 180
 AAAGCAAGGT GGGTTTGTAA CCTTTTACTT CACCACTGTG TGGCGNATGG CACCATCTGT 240
 50 CACCTGACCG GCTACCACAA GACGGAACAT TTTAAAAATT ACTGCTGTGC TCCTAAAATA 300
 ATTTTCAGCA AGTGCCATTT TACACCATCT TAGGAAGACA TCTGAGCTGA GCCCAATTCT 360
 GTCCCCACCA CCCACCCTAC AAGCGACCTG ACGCCTGTGG CCAGAATGCT GACTCTTCAT 420
 55 TCCAGGATAT TTATGTTTTC TAATAATAAA AGCAATAACT AGGCCAGAAA GAACACCACC 480
 TCAGAGCCCC CCTTCTCTGC TGCCCTGGGT CCACCCGTC TCATCCGCT GTGGGGCGAG 540
 60 TGGGGCTCTG CTGCAATGTG ACTGCAGTCT GAGGGGCAGA RGCTGCAGGK TACAGCCCCA 600

GCGAKTCACT CTCTGTCACC TGAATCTGA AACAAAGTGC TTCTGTGCCC CTGGGCTGGG 660
 AGTTTGTAT CTGAGGCTGC CTACCTGTTA GAACNTGTCA CCAGCAGGAC TTTATGTGCA 720
 5 TAAACAGCT TTCCTTCCAC CAAAAAAAAA AAAAAAAAAAC TCGAGGGGGG GCCCGGTACC 780
 CAATTCGCCC TATAGTGAGC GAT 803

10

(2) INFORMATION FOR SEQ ID NO: 18:

15 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1794 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

TTCTTTTTTG TTCATGGGAC ATGGTACCTA AGCAAATAGG AGTTGGGTTT GGTTTTTCTC 60
 CTAAAATAAT GCTCAATACT TACCTAATCA AATGGCATCC ATTTGAATAA AATGACAATA 120
 25 ACTAAAGCTA GTTAATGTCA GTGACATTAA ACTAACTCCA GGATTCAGGA GTTTTAATGT 180
 TAGAATTTAG ATTTAACAGA TAGAGTGTGG CTTCAATTGT CCATGGTAGC CCATCTCTCC 240
 30 TAAGACCTTT TCTAGTCTGT CTTCTGCCT TCGAACTTGA TGACAGTAAA ACCCTGTTTA 300
 GTATTCTCTT GTGCATTTGG TTGTGTGGTT AGCCGACTGT CTTGAACTA TTCATTTTGC 360
 TTCTAGTTTT ATTTTACAGA GGTAGCATTG GTGGTTTTTT TTTTTTTTTT CTGTCTCTGT 420
 35 GTTTGAAGTT TCAGTTTCTG TTTTCTAGGT AAGGCTTATT TTTGATTAGC AGTCAATGGC 480
 AAAGAAAAAG TAAATCAAAG ATGACTTCTT TTCAAATGT ATTGTTTAGC ACTTAACTCA 540
 40 GATGAATTTA TAAATTATTA ATCTTGATAC TAAGGATTTG TTACTTTTTT GCATATTAGG 600
 TTAATTTTTA CCTACATGT GAGAGTCTTA CCACTAAGCC ATTCTGTCTC TGTACTGTTC 660
 GGAAGTTTTG GAAACCCCTG CCAGTGATCT GGTGATGATC TGATGATTTA TTTAAAGAGC 720
 45 CGTTGATGCC TCCAGGAAAC TTAAGTATTT TATTAAATA TATATAGGAA TTTTTTTTTT 780
 TTTTGCTTTG TCTTCTCTC CCTTCTTTTA TCCTCATGTT CATCTTTCAA ACCAGTGTTC 840
 50 TGAAGTATG CATGCAGGCC TATAAATGAA AAACACAATT CTTTATGTGT ATAGCATGTG 900
 TATTAAATGTC TAACTACATA CGCAAAACT TCCTTTACAG AGGTCGGAC TAACATTTCA 960
 CATGCACATT TCAAAACAAG ATGTGTCATG AAAACAGCCC CTTTACCTGC CAAGACAAGC 1020
 55 AGGGCTATAT TTCAGTGACA GCTGATATTT GTTTTGAAAG TGAATCTCAT AATATATATA 1080
 TGTATTACAC ATTATTATGA CTAGAAGTAT GTAAGAAATG ATCAGAACAA AAGAAAATTT 1140
 60 CTATTTTCAT GCAAAATATTT TTCATCAGTC ATCACTCTCA AATATAAATT AAAATATAAC 1200

ACTCCTGAAT GCCTGAGGCA CGATCTGGAT TTAAATGTG TGGTATTCAT TGAAAAGAAG 1260
CTCTCCACCC ACTTGGTATT TCAAGAAAAT TTAAACGAT CCCAAGGAAA GATGATTGT 1320
5 ATGTTAAAGT GACTGCACAA GTAAAAGTCC AATGTTGTGT GCATGAAAAG GATTCCTTGG 1380
TTATGTGCAG GGAATCATCT CACATGCTGT TTTTCCTATT TGGTTTGAGA AACAGGCTGA 1440
10 CACTATTCTC TTTGATTAGA AAATAAATC ATAAACTCA TAATGTTGAT ATAATCAAGA 1500
TGTAACCACT ATAAATATGT AGAAGAGGAA GTTTTAAAG ACCTTAAGCT GGCATTGTGA 1560
AGGAACACCA TGGTAGACTC TTTTGTAAA TGTATTTTGT ATTAAATGAA ATGCAGTATA 1620
15 AAGGTTGGTG AAGTGTATA TAATTGTGTA AACAAATCCT GTTAATAGAG AGATGTACAG 1680
AATCGTTTGT TACTGTATCT TGAACCTGT GAAATAAAGA TTCCACCTCT GGTAAAAA 1740
20 AAAAAAAAAA AAYTCGGGGC CAGTCCCCC CCGCTATTT TAAAGGNAA AAAG 1794

25 (2) INFORMATION FOR SEQ ID NO: 19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1037 base pairs

(B) TYPE: nucleic acid

30 (C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

35 TCGAGTTTTT TTTTTTTTTT TGACAGAGTC TTGCTATGTT GCCCAGGCTG GAGTGCAGTG 60
GCAATCTTGG CTCAYTGCAA CCFYTGCTC CTGGGTCAA GCAATTYTCC TGCYTCAGCY 120
TCCYTAGTAG CTGGGACTAC AGGCACCTGC CACCATGCCA GGTAACTTT TTGTATTTTA 180
40 GTAGAGACAG AGTTTCACCA TGTGCCCCAC GCTGGTGTG AACTCCTGAG CTCAGGCAAT 240
CTGCCACCT TGGCTCCGA AAGTGCTAGG ATTACAGGCT TGAGCCACTG CACCCAGCCA 300
45 AGCTGTACTT TTTTTTTTTT TTMTAAAGCT TCAAACCTC AATATTTTAT TAAGAGTTAC 360
AGTTTGGTTT CAGTCATTCK GAGGAAATT AAGGAAGGG CTTGGCCCAW ACCTGGTAAA 420
AGAATGGAAG GAACCAATTT TTAACCATTT GGACCAGTGA TTYTCAATGG GAGTGCITTT 480
50 TGTCCCCCAG GAAACATCTR GAAAGGTATA WKGAGATATT TSTGGSTTGT CACAATTTGT 540
GATGGGGGAA AAAAGAACTA CCAGTATCAG GGGGATACAG GCCCGGTATC AGGTGGATAG 600
55 AGCCCTGGAA TATTGCTAAA CATCTACAG TGCAAAGACA SCCTTTMACA WACAGAACTA 660
TYTGGTCCAA AATGTCAATA GTGCTGAGGT TGAAGAACTC AATATTTTAT ATGTTTTCAG 720
GGAATTTCTA TGTGGGCTTG GGAAAGTTG AAGTCAATTG TCATTTGTAT ATTTAAAGGG 780
60

ATATATTTTA TCATTAGTCT ATAAATTCCA GTTGCAAAGT AGAGGCCCTG CACATTTGTG 840
 CACATATACA CACACCAGAA ATAAAYTMTK TKGCAATTAT CTTCTCTATC ATTGACAGGG 900
 5 CAATGACCTA TGAAAATTAT GTTATGTCTA ATAGTCCCTC ATTGTTATGT GCAAAACACC 960
 CAGCAAAGCT CAAGTTAAGR TTGTGGTCAC AAAGAAAAGA GCTATCATIG CTTTATGATG 1020
 10 TTGTCTGAAG TTAATGA 1037

15 (2) INFORMATION FOR SEQ ID NO: 20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1309 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

20 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

25 GGCACAGACT TTAAGAAATG CCAAATGCAA GGACCATTAA GAAATTTCTC CCCGAAATGA 60
 GGCTCCTCTA ACAAATGATG ATTANAACGC TCTCTCCTTG AGCAGTCACA TTCTAGAAAC 120
 ACGACATTCC ATGAGGCAGG AAGAGTTCAG TTAATTTGCT CCKGAAAAAG TGTGGTTCAG 180
 30 TGTTTGTGTG GCAATGTACG TGGGCAGAAG AGGCCGCTCA AGCTGTGTCC CCCCTGAGCA 240
 GGATTCAGGA AAGGGAAAAG AAGTTCTCTT CAACTCAGCC AAGGGGCCGT ACGATGGCCG 300
 35 ATGAGATTAT GTATTTAAAA GTTCTTTGTA AAGTGTAAC TAAAAACCTT AAATGTAAGA 360
 TGCTGTGTT ATTATTACTG TTGTGTGTGC TGTATGGAC ATGCCAAAAG GCCCTTGTTA 420
 GAAGACAGTT TTGCCTTTTC AATCTCATAG CAAGGAACTC AAGTCTGATG CTTCAAAAAG 480
 40 ATGAGAAGAA GGGCAAGAAG AGGGATAACT CCCAAGCTCA GAGGGAAAAA AAAGGTGGGG 540
 GAAAAGAGCC CCAGGGTGAC CTTCAGGAAA GGCCAGGACC AGGATGATCT AACCTTTCCC 600
 45 TTCACCAGAA ACAAAGCTAT TGCCAGACTG AACCTTAAAG TCAAGCAGTC ACCCACTGCC 660
 TTTGCTGGGA GCAGAAGCCC ATAGCAACAA GTGACCTGCC CCTCAGACTC AAGATCCCAG 720
 ATACCAGAGC TGGAGGAGTC ATAGGGCATT ACTGGTAGGC AGGAAAAC TG AGGGTCGAAC 780
 50 AAATGGAAGA ATGCGGTGAT CATAGACCAA AGACACACAG ATAATTAAAC CCATGTGTCC 840
 ACCCAGGCCA AAGTTCTTCC TGCTACCCCA CAGTGGATGT CCAGGCAGAT GGTCCCCACA 900
 55 TGATGGGGAA GCAGAGGGCA TAGTGTGGTT TTGTGGGACT TGTTCATGTT TTGTAGTGTG 960
 GGCTCAACAG TGCCAAAGGA AACACTAGGG AAAAGTTGGT GAAACATGCC AGCTAGCAGG 1020
 ACCAGTAAAG GCATAATCAG GCATTTGGCA AAGCTTGCTT TTCTAATTCA ATGATAGGTT 1080
 60 CTAATAGGAA ATTTTGAAG ATTTTAAAA ACAATGTTAT AGTGGCACTT CCCAGTATG 1140

GAATAAATAA CATGCATTCT TTTTTCATA TACTGTCATA TTCAGATGTC ATTAAAATAA 1200
 5 ATGGATGAGT CACAGAGGAG CTATCAGATG CTCTCATGAC TACCATAACT CAAAAAAAAA 1260
 AAAAAAAWA AAAGGGGGC CCGTACCCAT TTGCCCTAAA GGGATCGTA 1309

10

(2) INFORMATION FOR SEQ ID NO: 21:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1081 base pairs
 15 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

20 ACANATNITT TACTTAAATT TTATTTTATC TTATTTTATG GTGCTTTTAA TCTCAAAATT 60
 CTGAAAAGCG AATAGCACGT GTTTTCAGAA ACAAAATGTA AAGCAGTCAA ATTAAGTAGA 120
 25 TACTATTTAG AAATGTAAAA TACTCTCCAG ATCTACCATT AATAGAAAAT AACTAAACC 180
 TTATATTTTA TTTTTCGCAA AATATTTTAT TATAAAATAT GACCAAAATA TTTAAATGC 240
 ACAATGCTTT TAACTTAAAT GTGCTAACCC TGTTTCGTGC TGTTTTGTGC TGTACCTTTT 300
 30 CTGATTCMGA ATTATAGAAA ACTTGATAAA TACTTGATTT TAACCAATGA GACTACAGGC 360
 AGATGGGACT AAGTGTTTAT GGGACAATTA TGTACTATTT AACTTAAATA TATTTTGTIT 420
 35 AATAGGAAAT ATATAATAAT AGCATTTTAT GTAATAAAT ATGGGCAACG ATTATCTTGG 480
 AAATTAAAGA GTCAAAGCAA AGAAATGAAG GGCTGGTAAA ATGAATTTTG TAATATCCTC 540
 AGGATACTTT TATCTTAAAA GTATGTTGTT AAAGATTTTG TAAATGTAT TTCAACAATT 600
 40 TTAATGTGT TGAGCAAGTT GCAGTGCAA CACTGTCATT ATGTAGAGAG TTTATATGCA 660
 CATAATAACC TGTACCTATA AATCGTGCAA TAACCATATG CGACTATTTT GCCATGGAGA 720
 45 AATCTGACAG CATTGCAAAC AATAGTATTG TTTGATGTAG TTAACCTTAA GTTATTTTTC 780
 AGTAATTTCT TCACAAATCA AGATTCAAAC AGCTTTAAAC ACTTCCAATG AGATAAAATA 840
 TTTACTATTA TGCTTATTAG AACAAAAGGT GTTTAAGGAT GAACTAAATA TTTTAATTGA 900
 50 GCATTTATAT GGATAATCAT ACATTATGTA AGCCCATATG TATTTACATC CAGAGTCATA 960
 ATATTTTAAA TAAACAATCA TGCAGAACT TTTTATAGGG GTATACTATT GTTTTAATAT 1020
 55 CGTTGCCAAT TINGCTGACT TAAATATGT GACATTTTAA AATCAGGATT TTCCATATIN 1080
 G 1081

60

(2) INFORMATION FOR SEQ ID NO: 22:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 807 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

GAATTCGGCA CGAGCTCCTT CAGAAATGTC TTGGCTATTG TTGCTCTTTG CTCCTCTCTG 60
TAAATTTTAC CATAAACTTA RTTCCATAA TATATGACTG GAAATTTTAC AGAAGAGTTA 120
15 ATGTGTCTAA CTAGCAAACA CGAAGAAAAG CTCAGTGTTA GCAGTTAACT GAGGGAATGC 180
AAATCAAGAC CACAAGGAGA TAACAATTTG AGCCTATTGA CAAAAGTTCA GAAGTCTAAT 240
20 AATACTAAGT GTTGGAGAGG ATATGGCCCA GTATGATCTT ATCCACTGTT GGTGGGAGTA 300
TCAATTAGTA CAAACACTTT GAAAAATAAG ARGGAATTCT ATAATATCTA ACATTTGCAT 360
ATATCCATTT ATCTCTCTAG ATCTAGATCT TAGCCCTCTC CACCCTGCAC TGTGTTCTTG 420
25 GAAGGGGATC ATGAATGGTT TCCTTGCACT CTGCCTTCTG ATTTGGTTCA GCCAATGAGA 480
GACCATGGCA AGACATTTGT GAGAAGGGTA GAGAGTCAGG TCAAGGTTCT TAGTGAGATC 540
30 AACTCTTTCT CTGCCAGTTT GTTAACTGAA TTCTACTGAA AGCTAGAGCT CTGTTGAGTA 600
ATCTTTTAAA GCTGCAGCTA CCCTTTTGAG ATTAAGTAAT AGCTCCCTGT TTGTGCCTTG 660
TTAGGGCTAG GGATGTTTAA GGATCCTTGC CCTTGCTAGT CCTAGCATGT TTTGTTGTCC 720
35 CATAATAGTT CTTTTTTTAA ACTTTCCTCA ATTACACAAT TTGATCTTGT TCCTACCAGT 780
ACCNITGCTG GTACAACCTT AACTGCG 807

40

(2) INFORMATION FOR SEQ ID NO: 23:

(i) SEQUENCE CHARACTERISTICS:

- 45 (A) LENGTH: 632 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
50 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

GAATTCGGCA CGAGTCTAAC AGCATAAAGA AATAACAGCT GCATTCAAGA CCAGGATATG 60
55 TAAAATAATT TGTTTAGTTT CAGCCACTTT TTAAAGTCAA TTTTACACCC TGAAAGAAAG 120
GCAATCCTGA CTCCATTGTT CTTTCGCCAA TAAGGAGATC GGAATTACA ATAATAAATA 180
60 GAAGAAAGAA TGTGCTTTT CCTCACTGTA ATTAATTTTA TGGCTCTTGC GAAGATGAAT 240

TTTTGTGGTG ATTAAATAG TCCCTTGAC ATATTAGGTA CTCAGTAAGC ATTTGTGAAA 300
 TAGGGACTTT CTAGCCTTTA TTTGTGTTTA AGGAATCAGG GAATAAGTTC AAAATTCCTT 360
 5 TTCAAGAAAT TTTTGGAAGT CTCTTCTCAC TAAGAACTG TAAAGTCTTA TAAAAGAGAC 420
 ATTATTATT TTCTCCAAGT ATTGCTTGGC AGGTGAATG AAGGTTTTTT TTTTATCAAC 480
 AGTTGTTTTA TAAGATCGTT TGAGGACTAA AAGGGCTGAT TGTAATCACC TGTAACATGT 540
 10 TACCCAGCAA GACATTCCTC ACCAGGTTGA AGTAAAAAA ARAAATGAAG TGAGAATATC 600
 AAGCTTATGC AAGTTTGAAA TTNCAAACAA GA 632

15

(2) INFORMATION FOR SEQ ID NO: 24:

20

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1358 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 24:

GGCACGAGGA TAAATGCAA GTATTAAATCG GTCOCAACTT TAATATGGGA TAAAAATAAC 60
 30 AGTCAGTATG TGACCTCCTA AACAATCCCT CTACTGAGCT GTGGAGGGGA GAAGGAGGT 120
 CCTGGGGCCA GGACAGACAG GGCTATTTTC AGTAGTACAA CTTATATGCT ACTCTAAGAA 180
 AAGTCCAGAA AATGCRATTC TCTTCATACG AAGCTTTARA TACCCTCATK ATTTTGATAA 240
 35 ATACATTTTC ARRTCTAATA TGGAGACAGA AAGCTGCCTA GATTTATACC CACAAGTATT 300
 ATAAATTTAG AGAGTCTGAC CAGCCTCAAT TATTCTCTTT CGAAGTGGGA GAGAGAAATC 360
 40 AAAAGTCAGA AATGGTGGRT AATCTCCAAG TCATATCCAT TTGGSTTTGR TCTACTACTT 420
 GTTTTATGTC TTGTATTTGG RGRCAAGRT GCCTGATGTT AAGGGRATTT CMTACMTGA 480
 ATAATGTGAC CAGACTGCCA TCTAGTCAAA AACCTATAAA ATGTTATTTA CTTTAATTCT 540
 45 GGGCTAATTC AACAGAAGTY YYSGATAAAA RCTCTCCAAA CAATAATTAT GARCCTAGT 600
 TTTTGTGTTT GTTTTGATA CAAAACAAAA CAGCTCTGTA GTTGTCTGT GAGGTTTATA 660
 50 AATAGATTTT TTTAACTACT TAATTTTCYG GTTTCYGCCY CTGKGTTCYC TGTACCTATA 720
 GAGGTAGCTC TTTTCAGTTA AGTAGAGAAA AGCTCTTCCC CTGGGTGAA AATAATGCAG 780
 TCCCGAGAGG CTACTTAACT CTACCTTTCT GGAGGTCATG GTAGCAATTG GAGATCTCCC 840
 55 AGGCATTCTA AGGGAGCTA CTAAAGAGCC CCAGATACTC AATTTACCAC TAGAAATTCG 900
 CTTTATCTAC TCTCTGTCAT CTGGGAGRA AAGTATTATA ACTGACATTC AGTATGCACA 960
 60 CAATAAGTGC ATAATAAGA GCTATTGAGG GGATCCAAGG GAGTAAAATG GGTTCGCCA 1020

5 TAGGACTCCA TCAGGGTCCA CCAACACAGA CTTACAGCAA AAATTGGAAG GCTCTTTTCT 1080
 GCTGGATTCT GGAATCTGT GTTCTCTAGT GTGCCAGGA GAGTTGGAAT CAAAACACGT 1140
 AATATAATGT TTCTATTTCAG AGCCCCATTT TTTTGCCAAA TAAAGTAGCA CTGTCAAATA 1200
 ATAAATCTTG TATTCACTTG GGCATGTATG TTTATTATTG GATCTCTAAA ATATGCTTCA 1260
 10 AATAATGCAC TGAAATAAGT GAGGTGATGA ATTTTGAAAT AATAACAGTT TATGATGGGT 1320
 AGCTCCAAAA TTTTAAAAA AAAAAAAAAA AAACTCGA 1358

15

(2) INFORMATION FOR SEQ ID NO: 25:

20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1376 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 25:

CCCACCTTTA GCGAGCCAAC GAGAGAACAC CGCCTGCAGC TAGAACAGCC TGGTCAGGAG 60
 CGTAACGGAG TGGTGCGCCA ACGTGAGAGG AAACCCGTGC GCGGCTGCGC TTTCTGTCC 120
 30 CCAAGCCGTT CTAGACGCGG GAAAAATGCT TTCTGAAAGC AGCTCCTTTT TGAAGGGTGT 180
 GATGCTTGGA AGCATTTTCT GTGCTTTGAT CACTATGCTA GGACACATTA GGATTGGTCA 240
 35 TGGAAATAGA ATGCACCACC ATGAGCATCA TCACCTACAA GCTCCTAACA AAGAAGATAT 300
 CTTGAAAATT TCAGAGGATG AGCGCATGGA GCTCAGTAAG AGCTTTCGAG TATACTGTAT 360
 TATCCTTGTA AAACCCAAAG ATGTGAGTCT TTGGGCTGCA GTAAAGGAGA CTGGACCAA 420
 40 AACTGTGAC AAAGCAGAGT TCTTCAGTTC TGAAAAATGTT AAAGTGTTTG AGTCAATTAA 480
 TATGGACACA AATGACATGT GGTTAATGAT GAGAAAAGCT TACAAATACG CCTTTGAWAA 540
 45 GTATAGAGAC CAATACAACCT GGTCTTCTCT TGCACGCCCC ACTACGTTTG CTATCATTGA 600
 AAACCTAAAG TATTTTGTGT TAAAAAAGGA TCCATCACAG CCTTCTATC TAGGCCACAC 660
 TATAAAATCT GGAGACCTTG AATATGTGGG TATGGAAGGA GGAATGTCT TAAGTGTAGA 720
 50 ATCAATGAAA AGACTTAACA GCCTTCTCAA TATCCAGAA AAGTGTCTG AACAGGGAGG 780
 GATGATTGG AAGATATCTG AAGATAACA GCTAGCAGTT TGCCTGAAAT ATGCTGGAGT 840
 55 ATTTGCAGAA AATGCAGAAG ATGCTGATGG AAAAGATGTA TTTAATACCA AATCTGTTGG 900
 GCTTCTATT AAAGAGGCAA TGACTTATCA CCCCAACCAG GTAGTAGAAG GCTGTTGTTT 960
 AGATATGGCT GTTACTTTTA ATGGACTGAC TCCAAATCAG ATGCATGTGA TGATGTATGG 1020
 60

180

GGTATACCGC CTTAGGGCAT TTGGGCATAT TTTCAATGAT GCATTGGTTT TCTTACCTCC 1080
 AAATGGTTCT GACAATGACT GAGAAGTGGT AGAAAAGCGT GAATATGATC TTTGTATAGG 1140
 5 ACGTGTGTTG TCATTATTTG TAGTAGTAAC TACATATCCA ATACAGCTGT ATGTTTCTTT 1200
 TTCTTTTCTA ATTTGGTGGC ACTGGTATAA CCACACATTA AAGTCAGTAG TACATTTTTA 1260
 AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA 1320
 10 AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAA 1376

15

(2) INFORMATION FOR SEQ ID NO: 26:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 2923 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 26:

25 CTCCTCTCC GGGCCCCCT CCTCCCCCTT TMACTGGTGC AGATGGCCAG CCTGCTATAC 60
 CACCACCGCT TTCTGATACC ACCAAGCCCA AGTCCTCCTT GCCTGCCGTG AGCGATGCCC 120
 30 GTAGCGACCT GCTTTCAGCC ATCCGTCAAG GTTTTCAGCT GCGCAGGGTT GAKGAGCAGC 180
 GGGACAAGA GAAGCGGGAT GTTGTGGGCA ATGACGTGGC CACCATCTTG TCTCGTCGCA 240
 TTGCTGTTGA GTACAGTGAC TCAGAAGATG ACTCCTCTGA ATTTGATGAG GACGACTGGT 300
 35 CCGATTAACT CTTTCTGCCT GCTGCCACC TTTCTTTCT TTTCTTCTA CCTGCCTTCT 360
 TTGATGCCAA CCCCAACAGA CCCGTAGGGG AGGAAAAGG AGGAAAAAG TAATTTTAAG 420
 40 GGGCCAAAGC TTTCCCTGAA GCAACCAAAG ATATATCCAA GTGCTTCCTC CAAGTCAACA 480
 TGTATTTCCT CTCCTCATTT TCAGGCCCTG TGGGGCTCCT GAGGTTCACT AGCTGGGATG 540
 TTCCCTCTTT CCTTCAAGTG CCTGTGTCAT ATTGAAAGGA AGGAGAAATC CCAAAGCAGA 600
 45 TTCTTTGAT CGGTTTCTG TTGGAGATGG GGCTTCCCTT AGGAGCCATA TTCAACTACA 660
 GCCTTCTAAA ACCTGTGCCC TCAGCCACTT CGAATGCCAG CCACCTTCTG GTTCTAAAAC 720
 50 GGGGAGTGGT CTGAATGAAC ACAGCTGACC CCTTTCCCGC GCACTGAAAG GGCAGAGTAG 780
 GCCGAAGGTC CAAGGGCCAG ACTGCCTCAC CCTCTGCCCT AATCAGCAGG GTGGGCCTGC 840
 CTTTGTCTAA GCGATCTCTA TGCCTGGGAT GCCCTTTATT CCAGGAGGCA TCAAGCCTCT 900
 55 AAAGAATGTC TCACCTCCTC TGCCCAAAAA TGATGCCTTT CTGTAGGCTG GTGTTGTTGC 960
 CTCCTCCCA GGATCCCTTT GGTGAGTATG GTGTTGAGGA TGCACCACCA CCACCTCTAG 1020
 60 ATACCTTCAG GCAACACAGC CCAGTTTAA CCTCTAGTAT CCATGACCAA ACTATCCCTG 1080

	ACACATGAGG ACAGGGGCCT CTTCTGGCTG TCAGGAGCAA AGCCTGAAGA CTTGGAGCTG	1140
5	CAGGACTGGA AGAACAGTGG AGCCCCGTGG GTCTCACCCT TTAAGGATGC TGAGGCCTAG	1200
	AGATGGGAAG TGACTTGCTC AAGGTCACAC AATTGGATAG TGACATAGCT AGAGCGCAGA	1260
	GTTCCTGATT CCAAGTCACC TGTGCTTTCT GGGACCAAAG AATGGGCACC TGCTGGAGTC	1320
10	CGGGCAGAGC TTTCTCAGTT GTATTGCTAC TCCAGACCTC ACCATAGGTT GGGGTCCCAG	1380
	TAGGAAGGCT CAGGGTCTGT GCCAGCCCTG TCGGTGCTGC TCAGACCTTC ATAGCCTCTC	1440
15	TTGTCAATTCT TTGTTGCCCC TTTCTGTCA CCAGCCAACC ACATAGCCTT GGGACCAGCC	1500
	TCTCTGGGG ACCAGAAGTA GTGAGAGAAG GAAGGGGATA GGCAGCTTTG ACAGGTGCTG	1560
	CTTTCAATTC CTCTGCAACT CCTCCCCCTT TTATTTCCCC AATTAAACA AAGATTCTGC	1620
20	CAACTGTGGA AACTTCAGTC CCTCAGGCTG GCAGCCATGC CAGTACCTGC CTGGGGGTGG	1680
	GGGGTGCCTG GCAGCCATGA AGCAGGCTGA AAGGCAGAGG GGCTCCAGGT CCTGTTTCCA	1740
25	GCTCCCTCA CTGCACATGG TGAAGCTCGC TCCCTCCCTC CCTCCCTTCC CGCTTTTCCC	1800
	AGAGCTAATA CACAGGTGCT ATTATTCAGA AAAAACTGG TCAGCTCTAG CCAACAGTGA	1860
	AGGTTCTTCT TCTTCTGCCC TNAACTATTG TGTAGCCTCT TATGCTGAAA TCGGCTTCTG	1920
30	CTGGCTTCTC CGGCTTTCAG AGCCCTGAAA CAAAGAGAAA CAGGATCTGT CCCTACCCAG	1980
	CACAGCAAAT GGTGTAGTA ATTGCCAAG CCCTCATAAA GCCCTCCGGC TTGAGGAGAG	2040
35	AGTGTATAGT CATGGGTTCT GCCTCTGTGC CTTGCTGGC CGCTTCTCCT CTGCCTTCTT	2100
	TCCTGGAAT CAGGGTGTGG GGACTGAGCC TGTAGGGGAC AGCATGCCGT CTTGCTGTGG	2160
	CCACTCCCA GTGTGCCCTC TTCCCTCTTT ACACATCAGG TGTCTCTGGC ACAGGACTTG	2220
40	GCACTAAGCT CCATGCTGAG ACACCAGGCT ATGTGGGCCC CCACCTTGTT TCCAGCCTG	2280
	CACCTTAGAA GCCGAAGTGC TTTCATCAGA ACCCTAAAAT GGTGCTGAA GGCGCCTGGG	2340
45	CCGCAGCCAG CAGTAGTTGG AGAGGCAGGC AGAGGGCAGT GGTTCCTCCA AATAGGAGAC	2400
	CTGGGGCCTG GCCAGGCAGG GTTTGGGCCT AATGGCTTTG ACTAAATTAC CCCCATCCTC	2460
	CTTGCCCGGA AAAGGGAGAG CTAGAGCCAC TCACTGTCAT TCTGCTCTGA CCTTGAAGGG	2520
50	GGCGGTGTTG GCCTGGCTTC TGGAATGAC TGAGTCCATC GTGGAAAGGG CTGGGGGCAG	2580
	GAGGAGGTGG GGAGGGGCAC TGCCTGCGGA AGGTAGGATT AGATCATTAG CTCAGTGACC	2640
55	TCCTAGGGTT TCGATGTGCT ATGTTCTCAT CCTACAGTTG GTTTGGTAAT GATCTGCAAG	2700
	TCCCGAGAG CAACAGCACA GCTCTGCCCTG ACGCTCTCAT TAAATCTAT GCAGCCAAGC	2760
	TGGCACTTT GTAGCAGCCG GCCTTGCGAA GCCTCCTCAG CTCGGGGGGC CGGGGACCCA	2820
60	GTGAGCCGNA GAKCSTCTGG GCTCCACTTA TGCATATGCA CCAAAAAAAAA AAAAAAAAAA	2880

AAAAGGGGG CCGCTCTANA AGGATTCTTC NAAGGGGCCC AAG 2923

5

(2) INFORMATION FOR SEQ ID NO: 27:

- 10 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 775 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 27:

GAAGTAGTGN ATCCCCGGG CTGCAGGAAT TCGGCACGAG CCCRACCCSC ACCACCACCA 60
GAATGCAGTT CCAGCTTAGG AAGCCACAAA CAAGCCACCC AGGAGGAACA AACACCGCC 120
20 AGCGTGGATT TTCCCAAATT TCCCTGGAAA GTAAGTCTCG CTCTTGCCAA AGAAAAGTCT 180
GGCTTGGAGA GTCTCTGGAG CCCAGGATGC CAGCATGTGC CAATGACTGT CACCTTCATC 240
25 TCTTCAAAAG AAAAGCCATA GCCGAGGACT GTCCCGCGAC CCCCGTGGAC TCGTCTAGG 300
TCATGTGATT CTGTTTTCAT TTCTCATCCC ATCCAAITTTG TCCTTTTCTC CTGTCAITTT 360
CTTCCTCTGT GGTCCCTTCA AAGTTGTTAT AATTTGTACT GAAGTTCAAA ATGTGTCCCG 420
30 TTCTCCCCAG ACCACTCTAG CCACAGTATA TTGCAATAAA ATTACTTCTT ATATTTGCAG 480
AAATCTTTT GGTGTAATTT TATTTTTC TCTCAATATA TATAATTGGA CAAACGCTGG 540
35 CAAAAAGAAA AAAATGGTAA GCAAAAAACC CAAGATAAAG TTTCGAGGAC ATCAGGCCTT 600
TTGAAATACA ATGTCAAATG ACACATGTGA CGKTTTCAAA AAATCCGCTA GACATGTCAT 660
AAGTTTAAAC TGTAATGCCC AGGAAAGGAT ATCTTAAAT ATCTTAACT TGTGTAACAA 720
40 AGGAATAAAT AACTGTAATA GTTTTCAAT AAATCGAGTT GGGTGTTC ACCGT 775

45

(2) INFORMATION FOR SEQ ID NO: 28:

- 50 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 534 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 28:

GAATTCGGCA CGAGCAAGGG TGAACCTGA GTCTGCTTGT CTGTTTGCCC CATGACAGCC 60
CAGGGGTGGT GGCTCACCC CACCTCCAGG CAMCCACAAG AATATAAAAT CTTGTACAAR 120
60 GATGTCGATA TTAATATTGS CATTCCTAAG TGCACCTGCA CCTGTAGTAT CAGGTGGTTT 180

GCAGCCTTGG CTGCATAGCT GCATATGAGA ATCACCTGGG AAGCTTTTAA AAATCCCAGT 240
 ATCCCCACCT CTTCGCCAGT TACAGTGGAG TCTTGCGGGT GGTGGGGGAC ATCAATTATT 300
 5 TTTGAAAGCT CCMAAGTAAT TCTGGTGTGC AGTGGGGTGA CCAGCTGTCC CAGGGAMCTC 360
 CTTTAAAAAA TAATATCCCG GGCACATGAC AGGCCAATTG CCCTAATGCA ACCAAGGTTA 420
 10 AGAACTACTG GTTTAATGGG AAAATATTTT TTCCNGTGC TTGAATAATA CTGGTTTAT 480
 TAAACTCCNG AATCCCATTT CTTCCTTGC CAAATTTTTT AAAGGCNAAA AAAA 534

15

(2) INFORMATION FOR SEQ ID NO: 29:

(i) SEQUENCE CHARACTERISTICS:
 20 (A) LENGTH: 1827 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 29:

NNCNGCAGCA GCNCGGTCTT GTCCCGTCAG CGTCCCGCCA GCCAGCTCCT TGCACCCCTTC 60
 GCGGCCGAGG CGCTCCCTGG TGCTCCCCGC GCAGCCATGG CTCAGCACTT CTCCCTGGCC 120
 30 GCCTGCGACG TGGTGGGATT CGACCTGGAC CACACTCTGT GTCGCTACAA CCTGCCCGAG 180
 AGCGCCCCGC TCAATTATATA TAGCTTTGCC CAGTTCTAG TTAAGGAGAA AGGGTACGAT 240
 35 AAGGAATTGC TCAATGTGAC CCCAGAGGAT TGGGATTTCT GTTGCAAAGG TTTGGCATTG 300
 GATCTAGAAG ATGGGAACTT CCTTAACTT GCAAATAATG GCACTGTCTT CAGGGCAAGC 360
 CATGGCACCA AGATGATGAC TCCAGAGGTG CTGGCAGAGG CATATGGCAA GAAAGAGTGG 420
 40 AAGCACTTCT TGTCGGACAC TGAATGGCT TGCCGCTCAG GAAAGTATTA CTTTACGAC 480
 AACTACTTTG ACCTGCCAGG AGCTCTTCTG TGTGCCAGG TGGTGGACTA TTTAACAAAA 540
 45 CTGAACAATG GTCAAAAAAC ATTTGATTTT TGAAGGATA TAGTTGCTGC TATACAACAC 600
 AATTATAAAA TGTCAGCTTT TAAGGAAAAC TGTGGAATAT ATTTCCAGA AATAAAAAGA 660
 GATCCAGGCA GATATTTACA TAGTTGTCCT GAATCTGTGA AAAATGGCT TCGACAGCTA 720
 50 AAGATGCTG GGAAAATTCT TCTGTTAATT ACCAGTCTC ACAGTGATTA CTGTAGACTT 780
 CTCTGCGAAT ATATTCTTGG GAATGATTTT ACAGACCTTT TTGACATTGT GATTACAAAT 840
 55 GCATGAAGC CTGGTTTCTT CTCCCACTTA CCAAGTCAGA GACCTTTCCG GACACTCGAG 900
 AATGATGAGG AGCAGGAGGC ACTGCCATCT CTGGATAAAC CTGGCTGGTA CTCCCAAGGG 960
 AACGCTGTCC ACCTCTATGA ACTTCTGAAG AAAATGACTG GCAAACCTGA ACCCAAGGTT 1020
 60

	GTATTATTTG GTGACAGCAT GCATTCAGAT ATTTTCCCAG CTCGTCAC TA TAGTAATTGG	1080
	GAGACAGTCC TCATCCTGGA AGAACTCAGA GGGGATGAAG GCACGAGGAG TCAGAGGCCT	1140
5	GAGGAGTCAG AGCCTCTAGA GAAGAAAGGA AAATATGAGG GACCAAAAGC AAAACCTTTA	1200
	AATACTTCAT CTAAAAAATG GGGCTCTTTT TTTATTGATT CAGTTTGGG ACTGGAAAAT	1260
10	ACAGAAGACT CCTTGGTTTA TACATGGTCT TGTAAGAGAA TCAGTACTTA CAGCACTATT	1320
	GCAATTCCAA GTATTGAAGC AATCGCAGAA TTACCTCTGG ACTACAAATT TACAAGATTTC	1380
	TCTTCAAGCA ATTCAAAAAC AGCTGGCTAC TATCCAAATC CTCCACTGGT CTTATCAAGT	1440
15	GATGAGACAC TGATATCCAA ATAAGTTGTC TTTACTGAAA AATGAAGTGA AGACCCATAT	1500
	ATGCAGTTAA AAAAAAGTTA ATTTTCAAAA AATACTGTAA AAGACTTTAA GGAACAAGTT	1560
20	TTATTGACCA ATAAGTTGAT ATTTGTCCAT AGGTCTCCTT TCTATAAATC ATCTTGATGT	1620
	TTAACAATC TTATTATATT AAAATCTCAG TATCCTAAAA CTTAGGAACC TTATTGGATA	1680
	TTTTCTATTA CAGTAGTTTT GTGGTTGGGA TTCACCCGGG GGGGCCACAC ACTCACACGG	1740
25	CACAGTTCAC TCTTTACACA TATGGCCNCG GTCCCGTGGG GTTCTCNAAG GTGTGGTTCC	1800
	CTTGGGGCCT NMTGGGCTTG GGCCTTT	1827

30

(2) INFORMATION FOR SEQ ID NO: 30:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1479 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 30:

	GGCAGGAGG CGGGTGGCAT CAGCAGAGGG GCACGAGCCA AAGGGTGTGG CTACCTCACT	60
	GCTGGTCCCC AGGCCCGGGA GTGGGGAGC ACACACAGTG CCTTGGGTAC CCAGNTGGGT	120
45	GTCTCCCCG TGCAGAGGAG ACRGCAGCCT GGGTCCTGCC CTTACCTCT GCGGCTTTC	180
	TCTACATCGC CTGGTGAAC GTGCTCCCTG ACCTCTTGA AGAAGAGGAC CCGTGGCGCT	240
50	CCCTGCAGCA GCTGCTTCTG CTCTGTGCGG GCATCGTGGT AATGGTGTG TTCTCGCTCT	300
	TCGTGGATTA ACTTTCCTG ATGCCGACGC CCCTGCCCC TGCAGCAATA AGATGCTCGG	360
55	ATTCACCTCTG TGACCGCATA TGTGAGAGGC AGAGAGGGCG AGTGGCTGCG AGAGAGAATG	420
	AGCCTCCCCG CAGACAGGAG GGAGGTGCGT GTGGATGTAT GTGGTGTGCA CATGTGGCCA	480
	GAGGTGTGTG CCGAGACCG ACACTGTGAT CCCTGTGCTG GGTCCGGGGC CCAGTGTAGC	540
60	GCCTGTCCCC AGCCATGCTG TGGTTACCTC TCCTTGCCGC CCGTCACTT TCACCTCCTG	600

5 GAGTAAGCAG CGAGGAAGAG CAGCACTGGT CCCAAGCAGA GGCCTTGCCC TGCTGGGACC 660
 CCGGGAGTGA GAGCAGCCCA AGGATCCCAG GGTGCAGGGA ACTCCAGAGC TGCCCACTC 720
 CCACTGCCCC CTCAGCACAC ACACAGTCCC CAGGCGGCCT AGGGGCCAAG GCTGGGGCGG 780
 CTTTGGTCCC TTTTCCTGGC CCTTCCTTCC CCACTTCTAA GCCAAAGAAA GGAGAGGCAG 840
 10 GTGCTCCTGT ACCCCAGCCC CACTCAGCAC TGACAGTCCC CAGCTCCTAG TAGTGAGCTG 900
 GGAGGCGCTT CCTAAGACCC TTTCTCTCAGG GCTGCCCTGG GAGCTCATTC CTGGCCAACA 960
 CGCCCTGGCA GCACCAGCAG CTCCTTGCCAC CTCCAGCTGC CAAACAGCAG CCTGCCGGGC 1020
 15 AGGGAGCAGC CCCAGGCCAG AGAGGCCTCC CGGTCCAGCT CAGGGATGCT CCTGCCAGCA 1080
 CAGGGGCCAG GGACTCCTGG AGCAGGCACA TAGTGAGCCC GGGCAGCCCT GCCCAGCTCA 1140
 20 GGCCCCCTTC CTTCCTCATC GAGGTGGGG TAGGTGGGG CGGTGAGGGC TCCACGTTGT 1200
 CAGCGCTCAG GAATGTGCTC CGGCAGAGTG CTGAAGCCAT AATCCCCAAC CATTTCCCTT 1260
 GGCTGACGCC CAGGTACTCA GCTGGCCAC TCCACAGCCA GGCCTGCCCT GCCCTTCACC 1320
 25 GTGGATGTTT TCAGAAGTGG CCATCGAGAG GTCTGGATGG TTTTATAGCA ACTTTGCTGT 1380
 GATTCCGTTT GTATCTGTAA ATATTGTTC TATAGATAAG ATACAAATAA ATATTATCCA 1440
 30 CATAAAAAAA AAAAAAAAAA AACTTGGGGG GGGGNCCCG 1479

35 (2) INFORMATION FOR SEQ ID NO: 31:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 987 base pairs

(B) TYPE: nucleic acid

40 (C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 31:

45 GGCACGAGCG CAATCGCGTT TCCGAGAGA CCTGGCTGCT GTGTCCCGCG GCTTGGCTC 60
 CGTAGTGGAC TCCGCGGGCC TTCGGCAGAT GCAGGCCTGG GGTAGTCTCC TTTCTGGACT 120
 GAGAAGAGAA GAATGGAGAA GCCCTCTTC CCATTAGTGC CTTTGCAATTG GTTTGGCTTT 180
 50 GGCTACACAG CACTGGTTGT TTCTGGTGGG ATCGTTGGCT ATGTAAAAAC AGGCAGCGTG 240
 CCGTCCCTGG CTGCAGGGCT GCTCTTCGGC AGTCTAGCCG GCCTGGGTGC TTACCAGCTG 300
 55 TATCAGGATC CAAGGAACGT TTGGGGTTTC CTAGCCGCTA CATCTGTTAC TTTGTGTTGT 360
 GTTATGGGAA TGAGATCCTA CTACTATGGA AAATTCATGC CTGTAGGTTT AATTGCAGGT 420
 GCCAGTTTGC TGATGGCCGC CAAAGTTGGA GTTCGTATGT TGATGACATC TGATTAGCAG 480
 60

AAGTCATGTT CCAGCTTGA CTCATGAAGG ATTAAAAATC TGCATCTTCC ACTATTTTCA 540
ATGTATTAAG AGAAATAAGT GCAGCATTMT TGCATCTGAC ATTTTACCTA AAAAAAAAAA 600
5 GACACCAAAT TTGGCGGAGG GGTGGAAAAT CAGTTGTTAC CATTATAACC CTACAGAGGT 660
GGTGAGCATG TAACATGAGC TTATTGAGAC CATCATAGAG ATCGATTCTT GTATATTGAT 720
TTTATCTCTT TCTGTATCTA TAGGTAAATC TCAAGGGTAA AATGTTAGGT GTTGACATTG 780
10 AGAACCCCTGA AACCCCATTC CCTGCTCAGA GGAACAGTGT GAAAAAAAT CTCTTGAGAG 840
ATTTAGAATA TCTTTTCTMT TGCTCATCTT AGACCACAGA CTGACTTTGA AATTATGTTA 900
15 AGTGAAATAT CAATGAAAAT AAAGTTTACT ATAAATAAWA AAAAAAAAAA AAAAAAAAAA 960
AAAAAAAAA AAAAAAAAAA ANANAAA 987

20

(2) INFORMATION FOR SEQ ID NO: 32:

(i) SEQUENCE CHARACTERISTICS:
25 (A) LENGTH: 2933 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 32:

TCTACCTCCG AGTAGTATTA GACTGTAAAC ACAGTAATAT AGNCGCCATC ATTCTGTAAG 60
GGGTTTCTTT TCGGGGACAG AGGATCAGAT GTTGAGAGTT TGGACAAACT CATGAAAACC 120
35 AAAATATAC CTGAAGCTCA CCAAGATGCA TTTAAACTG GTTTTGCGGA AGGTTTCTG 180
AAAGCTCAAG CACTCACACA AAAACCAAT GATTCCCTAA GGCGAACCCG TCTGATTCTC 240
40 TTCGTCTGTC TGCTATTCGG CATTTATGGA CTTCTAAAAA ACCCATTTTT ATCTGTCCGC 300
TTCCGGACAA CAACAGGGCT TGATTCTGCA GTAGATCCTG TCCAGATGAA AAATGTCACC 360
TTTGAACATG TTAAAGGGGT GGAGGAAGCT AAACAAGAAT TACAGGAAGT TGTGAAATC 420
45 TTGAAAAATC CACAAAATT TACTATTCTT GGAGGTAAAC TTCCAAAAGG AATTCTTTTA 480
GTTGGACCCC CAGGGACTGG AAAGACACTT CTTGCCCGAG CTGTGGCGGG AGAAGCTGAT 540
50 GTTCCTTTTT ATTATGCTTC TGGATCCGAA TTTGATGAGA TGTTTGTGGG TGTGGGAGCC 600
AGCCGTATCA GAAATCTTTT TAGGGAAGCA AAGGCGAATG CTCCTTGTGT TATATTTATT 660
GATGAATTAG ATTCTGTGG TGGGAAGAGA ATTGAATCTC CAATGCATCC ATATTCAAGG 720
55 CAGACCATAA ATCAACTTCT TGCTGAAATG GATGGTTTGA AACCAATGA AGGAGTTATC 780
ATAATAGGAG CCACAACTT CCCAGAGGCA TTAGATAATG CCTTAATACG TCCTGGTCGT 840
60 TTTGACATGC AAGTTACAGT TCCAAGGCCA GATGTAAAAG GTCGAACAGA AATTTTGAAA 900

	TGGTATCTCA ATAAAATAAA GTTTGATCAW TCCGTTGATC CAGAAATTAT AGCTCGAGGT	960
5	ACTGTTGGCT TTTCCGGAGC AGAGTTGGAG AATCTTGTGA ACCAGGCTGC ATTAAGCA	1020
	GCTGTTGATG GAAAAGAAAT GGTACCATG AAGGAGCTGG GAGTTTCCA AAGACAAAAT	1080
	TCTAATGGGG CCTGAAAGAA GAAGTGTGGA AATTGATAAC AAAACAAAA CCATCACAGC	1140
10	ATATCATGAA TCTGGTCATG CCATTATTGC ATATTACACA AAAGATGCAA TGCCTATCAA	1200
	CAAAGCTACA ATCATGCCAC GGGGGCCAAC ACTTGGNACA TGTGTCCCTG TTACCTGAGA	1260
15	ATGACAGATG GAATGAACT AGAGCCCAGC TGCTTGACA AATGGATGT AGTATGGAG	1320
	GAAGAGTGGC AGAGGAGCTT ATATTTGGAA CCGACCATAT TACAACAGGT GCTTCCAGTG	1380
	ATTTTGATAA TGCCACTAAA ATAGCAAAGS GGATGGTTAC CAAATTTGGA ATGAGTGAAA	1440
20	AGCTTGGAGT TATGACCTAC AGTGATACAG GGAACTAAG TCCAGAAACC CAATCTGCCA	1500
	TCGAACAAGA AATAAGAATC CTTCTAAGGG ACTCATATGA ACGAGCAAAA CATATCTTGA	1560
25	AAACTCATGC AAAGGAGCAT AAGAATCTCG CAGAAGCTTT ATTGACCTAT GAGACTTTGG	1620
	ATGCCAAAGA GATTCAAATT GTTCTTGAGG GAAAAAGTT GGAAGTGAGA TGATAACTCT	1680
	CTTGATATGG ATGCTTGCTG GTTTTATTGC AAGAATAYAA GTAGCATGC AGTAGTCTAC	1740
30	TTTTACAACG CTTTCCCCTC ATTCTTGATG TGGTGTAAAT GAAGGTGTG AAATGCTTTG	1800
	TCAATCATTT GTCACATTTA TCCAGTTTGG GTTATTCTCA TTATGACACC TATTGCAAAT	1860
35	TAGCATCCCA TGGCAAATAT ATTTTGAAAA AATAAAGAAC TATCAGGATT GAAAACAGCT	1920
	CTTTTGAGGA ATGTCAATTA GTTATTAAGT TGAAAGTAAT TAATGATTTT ATGTTTGGTT	1980
	ACTCTACTAG ATTTGATAAA AATTGTGCCT TTAGCCTTCT ATATACATCA GTGGAACTT	2040
40	AAGATGCAGT AATTATGTTT CAGATTGACC ATGAATAAAA TATTTTTTAA TCTAAATGTA	2100
	GAGAAGTTGG GATTAAAAGC AGTCTCGAA ACACAGAGCC AGGGAATATA GCCTTTTGGC	2160
45	ATGGTGCCAT GGCTCACATC TGTAAATCCA GCACTTTTGG AGGCTGAGGC GGGTGGATTG	2220
	CTTGAGGCCA GGAGTTGAG ACCAGCCTGG CCAACGTGGT GAAACGCTGT YTCTACTAAA	2280
	ATACAAAAAA ATAGGGCTGG GCGCGGTTGC TCACGCCTGT AATCCCAGCA CTTTTCAGAG	2340
50	GCCAAGGCGG GCAATCACC TGAGGTCAAG AGTTTGAGAC CAGCCTGGCC AACATGGTGA	2400
	AACCCCATCT CTAATAACA TGCAAAAATT ACCTGGGCAT GGTGGCAGGT GCTTATAATC	2460
55	CCAGCTACTC TGGGGGCCAA GGCAGGAGAA TTGCTTGAGC CTGGGAGATG GAGGTTGCAG	2520
	TGAGCTGAGA TCATGCCACT GCACTCCAGC CTGGGCAACA GAGCAAGACT CTGCCTCAA	2580
	AAAAAATTAA AATAAATTTA AATACAAAA AAAATAGCCA GGTGTGGGGT GCATGCCCTGG	2640
60	AATCCCAGCT ACTTGAGAGG CTGAGGCACG AGAATTGCTT GAACCCAGGA GGTGGAGGTT	2700

GCAGTGAGCC AAGATCACAG GAGCCACTGC ACTCCAGCCT GGGTGACAGA GTGAGACTCT 2760
GTCTCAAAAM AAAATTAAAT AAATTATTAT AACCTTTCAG AAATGCTGTG TGCATTTTCA 2820
5 TGTTCITTTT TTTAGCATTA CTGTCACTCT CCCTAATGAA ATGTACTTCA GAGAAGCAGT 2880
ATTTTGTTAA ATAAATACAT AACCTCAAAA AAAAAAAAAA AAAAAAACT CGA 2933

10

(2) INFORMATION FOR SEQ ID NO: 33:

15

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1366 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 33:

GGGAATACCT ATTCTCCTTT ACCGTGTGTC TTTTCCCCCT GGAATTGAGC CAGCAAGTTC 60
25 TTGGCATGGC AGGTGTTTCT GAAATATCAG TGTGTTTTTY TTTGCTTTCT TTGTTTTCCT 120
TGTPTTGCTC TTTCTATTTT CCTAAGCAGG CAACTCCAAA AAGAGATTG TTTGTGCAGG 180
AGTCAGGAAA AGGGAAGAGG AATACTGAAA GCTGGGAGTA GGCAGGACA GAAGAGGGG 240
30 AGGAGTCTAT TTTCAATTG TGAGTCTTGA ACTTCCACCA ATGCCAAAGT CACGGACATG 300
TGTGCACTTG GATGTCGAG TTAGAGCAGC CCCAAGGGCC TGTAACTGA ATAGCAGGCA 360
35 CTCACCCAGC TGATAACTCA AGTTCCAAAT GGACCACAGC TGAGTTGTAG GGGATGTGTG 420
TGTGTGTGTA CGCGTCCGTT TGAGATTCTT GGAACAGATT TCCTCTGAGA TCTCAACAGG 480
CTTTTTCATT ATCATTGGGG AGCTATGGTT TCTCTTATTT CACAAGGCCC ATTTCTTCCT 540
40 TTTGAGATGT GCAAGGAGAT GACTCCATCC ATGACTTGGC TTTACACTCT CCTCCTTGG 600
CTTTTATCA TCAGTGCAGR AGARATTCTT GCTCGTCTT CAAACAATCT CATTCGAGCT 660
45 TTATAAAGAT TATTGGARTT TAAATAATAT TCATATCTAT GGCCTAGAAC AATGTTCTTC 720
AAGTATGCGT CAGAATCATG AGTGGTAGAG GGAGGATTAT AATGTAGTTT CCTACATTTT 780
TACCTCCAC CACCTGGAG TCTGCATTTT AACGTACTTC TGTGTGAGGA TCAGATTTG 840
50 GGAAGCGTGG GCTTGAGAT GTTTCTKGA CATTGATTTA TGTGAGACC AGACCAAGAA 900
GCAGATGGAT GGACATGATC AGTTCATAAA CATGTTCTT TCTTAGGGTC AAATTGGAGG 960
55 AGGCTCTAGA GAAGCACTGT CCAATAGAAA TATAATGCCA ACAATATATG TWATTTTAAG 1020
TCTTCTATTG GTGCATTTAA AAAGTAAAAG AAGGCTGAGT GGCTGGGCAT GGCTCCTCGT 1080
GCCTGTAATC CCAGCACTTT GGGAGGCCGG GGTGGGCAGA TCACCTGAGG TCAGGAGTTC 1140
60

GAGACCAGCC TGCCCAACAT GGTGAAACCC CATATNTACT AAAAATACAA AAAATTAAACC 1200
GGGCATAGTG GCAGGTGCCT GTAATCCCAG CTACTCGGGA GGCTGAGGCA GGAGAATCGC 1260
5 TTGAACCTGG GAGGCAGAGA CTGCAGTGAG CTGAGATCGT GCCACTACAC TCCAGCCTGG 1320
GTGATGAGCG AAACCTCCGTC TCAAAAAAAA AAAAAAAA ACTCGA 1366

10

(2) INFORMATION FOR SEQ ID NO: 34:

(i) SEQUENCE CHARACTERISTICS:
15 (A) LENGTH: 667 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 34:

ATTTTCGGCA CAGGCCGGA GCTACCTATC TGGTAGGGAG CTCCCCCAGC ACCGAAGACT 60
GCGATGACTT CTGCRCTGAC CCAGGGGCTG GAGCGAATCC CAGACCAGCT CGGCTACCTG 120
25 GTACTGAGTG AAGGTGCAGT GCTGGCGTCA TCTGGGGACC TGGAGAATGA TGAGCAGGCA 180
GCCAGTGCCA TCTCTGAGCT GGTGAGCACA GCCTGCGGTT TCCGGCTGCA CCGCGGCATG 240
30 AATGTGCCCT TCAAGCGCCT GTCTGTGGTC TTTGGAGAAC ACACACTGCT GGTGACGGTG 300
TCAGGACAGA GGGTGTMTGT GGTGAAGAGG CAGAACCGAG GTCGGGAGCC CATTGATGTC 360
TGAGCCTGCC GGAGGGCGAG GGTGCGAGAA GCGATTGGG TCCTGGGCCT CTGTGATGAG 420
35 GCAGGCACAN CTGTCGGTCT TGGCTTGCTG CTAGAACTAG GGCCTTCTGC TCGCCACCT 480
CCCACCCCTA CCTGGACGGG CCCAGGCTTG GGGACTCTGA GCTGTGTTAA GGAGAACAAG 540
40 GGCAAGGAGA CCTCCCTTTG TGCTCCCTCA CTCCTAATA AACATGAGTC TGATGTTCTC 600
CARMMMAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA 660
AAAAANN 667

45

50 (2) INFORMATION FOR SEQ ID NO: 35:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1710 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
55 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 35:

60 GGCACGAGCC AGAGCAGGCT GCTAGGCCTG GGGCCACCAC TGCCCCCTGG TGCTACACCC 60

	AGTGTGCTGG GTCACCTGGA ACTTCCTGAA GTGGTGTAC CTGAACTGGG CCCCCAAGGA	120
	TGGGGTGCGG GCAGTACCGC AGGAAGAGGA GCAGCCCCTG TGAAGATTGA GAGCTGCCAG	180
5	AGGCTCTGTG ATTGGCTGCG GCACGATGAC CCGCGCACGG ATTGGCTGCT TCGGGCCGGG	240
	GGGCCGGGCC CGGGGACAG AATCCGCCCC CGAACCTTCA AAGAGGGTAC CCCCCGGCAG	300
10	GAGNTGGCAG ACCTTAGGAG GTGCGACAGA CCCGCGGGGC AAACGGACTG GGGCCAAGAG	360
	CCGGGAGCGC GGGCGCAAAG GCACCAGGGC CCGCCCAGGG CGCCGCGCAG CACGGCCTTG	420
	GGGGTTCTGC GGGCCTTCGG GTGCGCGTCT CGCCTCTAGC CATGGGGTCC GCAGCGTTGG	480
15	AGATCCTGGG CTTGGTGTG TGCTTGGTGG GCTGGGGGGG TCTGATCCTG GCGTGCGGGC	540
	TGCCCATGTG GCAGGTGACC GCCTTCCTGG ACCACAACAT CGTGACGGCG CAGACCACCT	600
20	GGAAGGGGCT GTGGATGTCG TGCGTGGTGC AGAGCACNGG GCACATGCAG TGCAAAGTGT	660
	ACGACTCGGT GCTGGCTCTG AGCACCAGAG TGCAGGCGGC GCGGGCGCTC ACCGTGAGCG	720
	CCGTGCTGCT GCGCTTCGTT GCGCTCTTCG TGACCCCTGGC GGGCGCGCAG TGCACCACCT	780
25	GCGTGCCCC GGGCCCGGCC AAGGCGCGTG TGGCCCTCAC GGGAGGCGTG CTCTACCTGT	840
	TTTGCGGGCT GCTGGCGCTC GTGCCACTCT GCTGGTTTCG CAACATGTTC GTCCGCGAGT	900
30	TTTACGACCC GTCTGTGCCC GTGTGCGAGA AGTACGAGCT GGGCGCANGC TGTACATCGG	960
	CTGGGCGGCC ACCGCGCTGC TCATGGTAGG CCGCTGCCTC TTGTGCTGCG GCGCCTGGGT	1020
	CTGCACCGGC CGTCCCGACC TCAGCTTCCC CGTGAAGTAC TCAGCGCCGC GCGGGCCAC	1080
35	GGCCACCGGC GACTACGACA AGAAGAACTA CGTCTGAGGG CGCTGGGCAC GCGCGGGCCC	1140
	CTCCTGCCAG CCACGCTGC GAGGCGTTGG ATAAGCCTGG GGAKCCCCGC ATGGACCGCG	1200
40	GCTTCGCGCG GTTAGCGCGG CGCGCAGGCT CCTCGGAACG TCCGGCTCTG CGCCCCGACG	1260
	CGGCTCTGG ATCCGCTCCT GCCTGCGCCC GCAGCTGACC TTCTCCTGCC ACTAGCCCGG	1320
	CCCTGCCCTT AACAGACGGA ATGAAGTTTC CTTTTCTGTG CGCGCGCTG TTTCCATAGG	1380
45	CAGAGCGGGT GTCAGACTGA GGATTTCGCT TCCCCCCTCA GACGCTGGGG GTCTTGGCTG	1440
	CTGCCCTTACT TCCAGAGGC TCCTGCTGAC TTCGGAGGGG CGGATGCAGA GCCCAGGGCC	1500
50	CCACCGGAA GATGTGTACA GCTGGTCTTT ACTCCATCGG CAGGCCCGAG CCCAGGGACC	1560
	AGTGACTTGG CTTGGACCTC CCGGTCTCAC TCCAGCATCT CCCAGGCAA GGCTTGTGGG	1620
	CACCGAGCT TGAGAGAGGG CGGAGTGGG AAGGCTAAGA ATCTGCTTAG TAAATGGTTT	1680
55	GAACTCTCAA AAAAAAAAAA AAAAAAAAAA	1710

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1096 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 36:

10 GGCCAGTGGG CAGGGTCACA GGGCAAGGTC CCGGGGGCCG CTGGGTGCGG CGACTTCCGT 60
GCTCCCGGCG AGCGGGCGGA GAGCGGGGCG CGCACTGGGG AGTGTGGGCT GGGCCGCAGA 120
15 TGTCATGTGG CCTGKTTTTT GGACCGTGGT TCGTACCTAT GCTCCTTATG TCACATTCCC 180
TGTTGCCTTC GTGGTCGGGG CTGTGGGTTA CCACCTGGAA TGGTTCATCA GGGGAAAGGA 240
CCCCCAGCCC GTGGAGGAGG AAAAGAGCAT CTCAGAGCGC CGGGAGGATC GCAAGCTGGA 300
20 TGAGCTICTA GGCAAGGACC ACACGCAGGT GGTGAGCCTT AAGGACAAGC TAGAATTGTC 360
CCCGAAAGCT GTGCTGAACA GAAACCGCCC AGAGAAGAAT TAATGGAGGA CACAGGGCCC 420
TATGGTCTTA CTGTGGGTGG TGACTTGTCC TGCTACCATG TTGACAGAGC CCCAGAACCC 480
25 ACATCTAATT GGCTTTGTG CTTATTCTGG CCCTTCCCAC ACCACACAGC CACACAAATA 540
CTGGCTGCTC CTGATGGCC AGGCAGACCC AGCAGCAGCC GAGGGGCCAG TGAAGAGGAA 600
30 GGCCGCATCT GTTGTGTGGT GGCCACAAGC ACTCAGGCAT CTGAGTTTAC TGGTGCACTG 660
CTGGGAGGAG AGTTATGAGA TGAACATTGG CTGTCAATCT CTGTGGGCAG GCGGTTTGGC 720
CTCTAGTGGG AATGGCTGGG ATTTGGGCGT TGCCTTTAGG AGGGATACCT GCATGTCTAG 780
35 TTCCAGTCTG CACTGGAAAG AATTCAAATA TGCACCTGGC TCCCTTCACT ATTTTGCCCT 840
ATCCTTTGTG CTCATICTTA CTGAAATCTG TCTGTTCAGC TCAGGAATGG GATTCCCCCA 900
40 GGAAGGAAAG CACTTTTCTG TTCTGGGAAG CCCAGACTGT TCACTTTGGG GCAGGGACGA 960
ACATGTGCCT CGTGAATTTG CTTGAAAACA GTCACCATCT TCTACCCCCA TCACTGTATA 1020
GTGAAAAACC TGATTAAAGT GGTATCTGAG AACCAWAAAA AAAAAAAAAA AAAAAAAAAA 1080
45 AAAAANGGGG GGNCCC 1096

50

(2) INFORMATION FOR SEQ ID NO: 37:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2279 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 37:

60

	GGTGGGCAAG GGGCTCAGCT CGCAGCGCAT GCCCCGCGAC AGGTTGCTGC TGGCCGTGGG	60
	CAGCGCCGTC TTAAATGCCA TGTTCAACGG GGGMATGGCC ACAACATCCA CGGAGATTGA	120
5	GCTGCCCCGAC GTRGAACCCG CCGCCTTCCT CGCACTGCTC AAGTTTCTCT ACTCGGACGA	180
	GGTGAGATT GGCCCGGAGA CGGTGATGAC CACGSTATAC ACCGCCAAGA AGTACGCGGT	240
10	GCCAGCGCTC GAGGCCCATTT GCGTGGAGTT CCTGAAGAAG AACCTGCGAG CCGACAACGC	300
	CTTCATGCTG CTCACGCAGG CGCGACTCTT CGATGAACCG CAGCTGGCCA GCCTGTGCCT	360
	GGAGAACATC GACAAAAACA CTGCAGACGC CATCACCGCG GAGGGCTTCA CCGACATTGA	420
15	CCTGGACACG CTGGTGGCTG TCCTGGAGCG CGACACACTG GGCATCCGTG AGGTGCGGCT	480
	GTTCAATGCC GTTGTCCTGCT GGTCCGAGGC CGAGTGTGAG CCGCAGCAGC TGCAGGTGAC	540
20	GCCAGAGAAC AGGCGGAAGG TTCTGGGCAA GGCCCTGGGC CTCATTGCTT TCCCGTCTAT	600
	GACCATCGAG GAGTTCGCTG CAGGTCCCGC ACAGTCGGGC ATCCTGGTGG ACCGCGAGGT	660
	GGTCAGCCTC TTCTGCACTT CACCGTCAAC CCCAAGCCAC GAGTGGAGTT CATTGACCGG	720
25	CCCCGCTGCT GCCTGCGTGG GAAGGAGTGC AGCATCAACC GCTTCCAGCA GGTGGAGAGT	780
	CGCTGGGGCT ACAGSGGGAC CAGTGACCGC ATCAGGTCTT CAGTCAACAA GCGCATCTTC	840
30	GTGGTGGGAT TTGGGCTGTA TGGATCCATC CACGGGCCCA CCGACTACCA AGTGAACATC	900
	CAGATTATTC ACACCGATAG CAACACCGTC TTGGGCCAGA ACGACACGGG CTTCAGCTGC	960
	GACGGCTCAG CCAGCACCTT CCGCGTCATG TTCAAGGAGC CGGTGGAGGT GCTGCCCAAC	1020
35	GTCAACTACA CGGCTGTGC CACGCTCAAG GGCCAGACT CCCACTACGG CACCAAAGGC	1080
	CTGCGCAAGG TGACACACGA GTCGCCACC ACGGGCGCCA AGACCTGCTT CACCTTTTGC	1140
40	TACGCGCGCG GGAACAACAA TGGCACATCC GTGGAGGACG GCCAGATCCC CGAGGTGATC	1200
	TTCTACACCT AGGCTGCCCG ACACCGACAC CGCCTCCCT CCGTGGGGAT AGCCGCAGCC	1260
	CCAGGCCATC ATCTGCTGCT GGGGYCCCC CACCACGCGG TGCCAGGCC AGTGTCCCC	1320
45	AGGCCGTCTG TCCACTCCAT GCCACCTTTC TCAGCATCAG GACGGGGTTG CCCTGTGTTC	1380
	ACCACGAGTK TGGCTGCTGG ATCAGGGCAG CCGGGGAGGT GGCCAGGCCA GTGGCCAGGC	1440
50	CCTGTGGAGA CAATCCCTCA GGAATAGGGA CAGGGCTGTG CCGGCTGGG CCAGGGCCCA	1500
	CGGACCGCA GCTCAGGGCG CTGCCCCAGC TCGTCTGCCG GCGGTGCGCC GCGGGCGTCC	1560
	CTCGCGTCTC TTCACTGCAC ATTGCAATGC ATTTGCGATT CCCATTTCTC TGCTAGGAGC	1620
55	CAGCCTGGGT GCGCTGCTC CCAGAGCCGT GGGTCCCAGA CCTTGCGTTC CTTTGTGTTCC	1680
	TGTCCGTTTA TCAGGACACG GGGCCACCT GTCACGTGCC CGAGGCCACC CAAGCCCAGC	1740
60	CTCGGGGCG TTCCCACTGC CTGGATGCCG GCTTGAGTTC TGCGCACGCA GGATTGAGT	1800

5 TGGGGACGGC CCCTGCCGGA TAGGCCTAGC CCTGGCCCAG GTGGTGAGCG GTTTGCAGTG 1860
TCCGTCTCTCA TCCACCTGAT GGGCCCAGAT AAAGGCCCCC GCTGTCCAGC CTCCTTGGAC 1920
GGCCCTCGCG GTCCCTGCAG CCCAAGATGG GACTCAGACC CTGTGCCCCA GAGCTCCCTT 1980
GCCGCAGAAAT GGGGCCCCAG CCGGCCCCGA CCGGGTCCAG GAGCACTGCT CGCCTGTACA 2040
TACTGTTGCC CTAGCCCACC TGGTGCCGTG GGAGCCACCC CCAGGTGCTG GGGCACAGCC 2100
CCTCCCCACT CCGGCCACGC CCCACCCAC CCGCGTGTTC TCTGCCCTGT GACTCCTGGA 2160
ACCTGCGTCC TCCCCAAAGC CATGGGAGGG GTGTCTCTCT CAGACCATGC CCCAGATGA 2220
15 TTTTTTTTAAA TAAAGAAACA AATGCACCTG CAAAACAAAA AAAAAAAAAA AAAACTCGA 2279

20 (2) INFORMATION FOR SEQ ID NO: 38:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 745 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 38:

30 GTACAGGACT GAGAAGCAGA TAACAAGAGT GACGCTCACA GGGCTGGGCT GACGCTAACA 60
GGAGGCAGTG TGTGGCTCGA AGATTCTTGA ACCCACAGCA GCAGCTGCGG CCACCCCATC 120
CTGCCCACAG CTCCAGCCCT GAGACGACGA GGAGGAGAGT CGACTTTGCC TCTTGCCCAA 180
35 GGGACCATGC CCAGGTGCGG GTGGCTCTCC CTGATCCTCC TCACCATTC CTTGGCCCTG 240
GTGGCCAGGA AAGACCCAAA AAAGAATGAG ACGGGGGTGC TGAGGAAATT AAAACCCGTC 300
40 AATGCCPTCA ANTGCCAACG TGGAAGCAGT GTYYGTGGTT TTGCCATGCA AGAATACAAC 360
AAAGAGAGCG AGGACAAGTA TGTCTTCCTG GTGGTCAAGA CACTGCAAGC CCAGCTTCAG 420
GTCAAAATC TTCTGGAATA CCTTATTGAT GTAGAAATG CCCGACGCGA TTGCAGAAAG 480
45 CCTTTAAGCA CTAATGAAAT CGCGCCATTC AAGARAATC CAAGCTGAAA AGGAAATTAA 540
GCTGCAGCTT TTTGGTAGGA GCACCTCCCT GGAATGGTGA ATTCACTGTG ATGGAGAAAA 600
50 AGTGTGAAGA TGCTTAATGG TGTTTTGAGG CATCCCTCCA ACCTCTGTGA CTACTTTATC 660
CATGAAAAATG AAGCAATGGT CAGGTGGGAG GCTCTTCCCA ATGTGCTTTC TTCAAAAAAA 720
AAAAAAAAAA AAAAAAAAAA CTCGA 745
55

60 (2) INFORMATION FOR SEQ ID NO: 39:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1718 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 39:

10	CCCCATAGGC AGGAGGCCCC CGGGCAGCAC ATCCTGTCTG CTGTGTCTG CTGCAGAGTT	60
	CTGTCTCTGC ATTGGTGGCG CTCAGGCCAG GCTGCACTGC TGGGACCTGG GCCATGTCTC	120
	CCCACCCAC CGCCCTCCTG GGCCTAGTGC TCTGCCTGGC CCAGACCATC CACACGCAGG	180
15	AGGAAGATCT GCCCAGACCC TCCATCTCGG CTGAGCCAGG CACCGTGATC CCCCTGGGGA	240
	GCCATGTGAC TTTCGTGTGC CGGGGCCCGG TTGGGGTTCA AACATTCCGC CTGGAGAGGG	300
20	AGAGTAGATC CACATACAAT GATACTGAAG ATGTGTCTCA AGCTAGTCCA TCTGAGTCAG	360
	AGGCCAGATT CCGCATTGAC TCAGTAAGTG AAGGAAATGC CGGGCCTTAT CGCTGCATCT	420
	ATTATAAGCC CCCTAAATGG TCTGAGCAGA GTGACTACTG GAGCTGCTGG TGAAAGAAAC	480
25	CTCTGGAGGC CSGGACTCCC CGGACACAGA GCCCGGCTCC TCAGCTGGAC CCACGCAGAG	540
	GCCGTCGGAC AACAGTCACA ATGAGCATGC ACCTGCTTCC CAAGGCCTGA AAGCTGAGCA	600
30	TCTGTATATT CTCATCGGGG TCTCAGTGGT CTTCCTCTTC TGTCTCCTCC TCCTGGTCTT	660
	CTTCTGCCTC CATCGCCAGA ATCAGATAAA GCAGGGGCCC CCCAGAAGCA AGGACGAGGA	720
	GCAGAAGCCA CAGCAGAGGC CTGACCTGGC TGTGTATGTT CTAGAGAGGA CAGCAGACAA	780
35	GGCCACAGTC AATGGACTTC CTGAGAAGGA CAGAGAGACG GACACCTCGG CCCTGGCTGC	840
	AGGGAGTTCC CAGGAGGTGA CGTATGCTCA GCTGGACCAC TGGGCCCTCA CACAGAGGAC	900
40	AGCCCGGGT GTGTCCCCAC AGTCCACAAA GCCCATGGCC GAGTCCATCA CGTATGCAGC	960
	CGTTGCCAGA CACTGACCCC ATACCCACCT GGCCTCTGCA CCTGAGGGTA GAAAGTCACT	1020
	CTAGGAAAAG CCTGAAGCAG CCATTGGAA GGCTTCTGT TGGATTCTC TTCATCTAGA	1080
45	AAGCCAGCCA GGCAGCTGTC CTGGAGACAA GAGCTGGAGA CTGGAGGTTT CTAACCAGCA	1140
	TCCAGAAGGT TCGTTAGCCA GGTGGTCCCT TCTACAATCG AGCAGCTCCT TGGACAGACT	1200
50	GTTTCTCAGT TATTTCAGA GACCCAGCTA CAGTTCCCTG GCTGTTCTA GAGACCCAGC	1260
	TTTATTACCC TGA CTGTTTC CAGAGACCCA GCTAAAGTCA CTGCCTGTT CTAAGGCC	1320
	AGCTACAGCC AATCAGCCGA TTCTCTGAGC AGTGATGCCA CCTCCAAGCT TGTCTAGGT	1380
55	GTCTGCTGTG AACCTCCAGT GACCCAGAG ACTTTGCTGT AATTATCTGC CTGCTGACC	1440
	CTAAAGACCT TCCTAGAAGT CAAGAGCTAG CCTTGAGACT GTGCTATACA CACACAGCTG	1500
60	AGAGCCAAGC CCAGTTCTCT GGGTGTGCT TTA CTCCACG CATCAATAAA TAATTTTGAA	1560

GGCCTCACAT CTGGCAGCCC CAGGCCTGGT CCTGGGTGCA TAGGTCTCTC GGACCCACTC 1620
TCTGCCTTCA CAGTTGTTC AAGCTGAGTG AGGGAAACAG GACCTACGAA AAAAAAAAAA 1680
5 AAAAAAATCG AGGGGGGGCC CGTACCCAAT CGCCTGTA 1718

10 (2) INFORMATION FOR SEQ ID NO: 40:

(i) SEQUENCE CHARACTERISTICS:

15 (A) LENGTH: 1966 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 40:

20 GTCGCGCCTG CAGGTCGACA CTAGTGGATC CAAAGAATTC GGCACGAGCT GGGGAGCGGG 60
ACTSGAGAAT ACTGCCAGT TACTCTAGCG CGCCAGGCCG AACCGCAGCT TCTTGGCTTA 120
GGTACTTCTA CTCACAGCG CCGATTCCGA GGCCAACTCC AGCAATGGCT TTTGCAAATC 180
25 TGCGGAAAGT GCTCATCAGT GACAGCCTGG ACCCTTGCTG CCGGAAGATC TTGCAAGATG 240
GAGGGCTGCA GGTGGTGGAA AAGCAGAACC TTAGCAAAGA GGAGCTGATA GCGGACTGCA 300
30 GGACTGTGAA GGCCTTATTG TTCGCTCTGC CACCAAGGTG ACCGCTGATG TCATCAACGC 360
AGCTGAGAAA CTCAGGTGG TGGCAGGGC TGGCACAGGT GTGGACAATG TGGATCTGGA 420
GGCCGCAACA AGGAAGGGCA TCTTGGTTAT GAACACCCCC AATGGGAACA GCCTCAGTGC 480
35 CGCAGAACTC ACTTGTGGAA TGATCATGTG CCTGGCCAGG CAGATTCCCC AGGCGACGGC 540
TTCGATGAAG GACGGCAAAT GGGAGCGGAA GAAGTTCATG GGAACAGAGC TGAATGGAAA 600
40 GACCTGGGA ATTCTTGGCC TGGCAGGAT TGGGAGAGAG GTAGCTACCC GGATGCAGTC 660
CTTTGGGATG AAGACTATAG GGTATGACCC CATCATTTCC CCAGAGGTCT CGGCCTCCTT 720
TGGTGTTCAG CAGCTGCCCC TGGAGGAGAT CTGGCCTCTC TGTGATTTC TCACTGTGCA 780
45 CACTCCTCTC CTGCCCTCCA CGACAGGCTT GCTGAATGAC AACACCTTTG CCCAGTGCAA 840
GAAGGGGGTG CGTGTGGTGA ACTGTGCCCC TGGAGGGATC GTGGACGAAG GCGCCCTGCT 900
50 CCGGGCCCTG CAGTCTGGCC AGTGTGCCG GGTGCACTG GACGTGTTTA CGGAAGAGCC 960
GCCACGGGAC CGGGCCTTGG TGGACCATGA GAATGTCATC AGCTGTCCCC ACCTGGGTGC 1020
CAGCACCAAG GAGGCTCAGA GCCGCTGTGG GGAGGAAATT GCTGTTTCA TCGTGGACAT 1080
55 GGTGAAGGGG AAATCTCTCA CGGGGGTGTG GAATGCCAG GCCCTTACCA GTGCCCTCTC 1140
TCCACACACC AAGCCTTGGA TTGGTCTGGC AGAAGCTCTG GGGACACTGA TGCAGCCTG 1200
60 GGTGGGTTC CCCAAAGGGA CCATCCAGGT GATAACACAG GGAACATCCC TGAAGAATGC 1260

5 TGGGAAGTGC CTAAGCCCCG CAGTCATTGT CGGCCTCCTG AAAGAGGCTT CCAAGCAGGC 1320
 GGATGTGAAC TTGGTGAACG CTAAGCTGCT GGTGAAAGAG GCTGGCCTCA ATGTCACCAC 1380
 CTCCCACAGC CCTGCTGCAC CAGGGGAGCA AGGCTTCGGG GAATGCCTCC TGGCCGTGGC 1440
 CCTGGCAGGC GCCCCTTACC AGGCTGTGGG CTTGGTCCAA GGCACCTACRC CTGTACTGCA 1500
 10 GGGGCTCAAT GGAGCTGTCT TCAGGCCAGA AGTGCCTCTC CGCAGGGACC TGCCCTTGCT 1560
 CCTATTCGG ACTCAGACCT CTGACCCTGC AATGCTGCCT ACCATGATTG GCCTCCTGGC 1620
 AGAGGCAGGC GTGCGGCTGC TGTCCTACCA GACTTCACTG GTGTCAGATG GGGAGACCTG 1680
 15 GCACGTCATG GGCATCTCCT CCTTGCTGCC CAGCCTGGAA GCGTGAAGC AGCATGTGAC 1740
 TGAAGCCTTC CAGTTCCTACT TCTAACCTTG GAGCTCACTG GTCCCTGCCT CTGGGGCTTT 1800
 20 TCTGAAGAAA CCCACCCACT GTGATCAATA GGGAGAGAAA ATCCACATTC TTGGGCTGAA 1860
 CGCGGGCCTC TGACACTGCT TACACTGCAC TCTGACCCTG TAGTACAGCA ATAACCGTCT 1920
 AATAAGAGC CTACCCCCAA AAAAAAAAAA AAAAAAAAAA ACTCGA 1966
 25

30 (2) INFORMATION FOR SEQ ID NO: 41:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 972 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 35 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 41:

40 GGCACGAGCC AAGTGGTCCC CCAGACAAGG CTCAGGATGT CCACATCCAC TGCATCCTGG 60
 ACCCTGTGCA GGTGAAGATG TCCCGACCCA CGCATACTCC TCTTTCGCCT GCCACCATTT 120
 CTCCAACCAT CACAGTAGCA GTCTTCTTCG CTGTGTTCGT CGCCGCGGCC GCCGCCACCG 180
 45 CCGTGTTCGC CGTCGCTGCT GCAACCACCA GCAGCGGSCG CAGAACTASA GACAAATCCC 240
 CCATAGCCAC TCAGTCTTCC GTAACCCACA TGCAGCCAA AAGATGTCAC AACTACACCG 300
 AGTGCCTTTC TTTGATCAGG ARGACCCGGA TTCTTACCTG GARGARGARG ACAACCTGCC 360
 50 CTTCCCGTAT CCCAAGTACC CACGTCGCGG CTGGGCGGGG TTTTATCAGA GAGCGGCGCT 420
 GCCTCCAATG TGGGCGTGTG GGGCCACCAG GGTGTATCCT GGCCAGTCTG CCACCACCTT 480
 55 CTCTCTACCT GTCACCTGAG CTGCGCTGCA TGCCCAAGCG TGTAGAGGCC AGGTCTGAGC 540
 TGAGGCTCTG CCCGCTGGC GTCTCTGAC TACCTCTGCC TCCCTCACGG TGTGGACGA 600
 60 GGCTCCCAT CAACGGACCC CAGCTCCAAG CTCAGTGTG GTCCCCCATT CTTCCAGCC 660

CTGGCCCAAA GTCCAGGCTG CGGACCCCTGC CCCTCCCCCG ACCATGTTTG TCCCCTCAG 720
 CCGGAATCCA GGGGGCAATG CCAACTACCA GGTGTACGAC AGCCTGGAGC TGAAGCGGCA 780
 5 GGTGCAGAAG AGCAGAGCCA GGTCCAGCTC ACTGCCACCG GCTTCCACCT CCACCTTGAG 840
 GCCCTYTCTG CACAGGAGCC AGACCGAGAA ACTCAACTGA CCAGCAGGCG GATGTGGGGT 900
 10 GTGGGGCAGG GCATGGAGGG AGAGGAATAA AGAGAAACAG AGTCCAGGAA AAAAAAAAAA 960
 AAAAAAATC GA 972

15

(2) INFORMATION FOR SEQ ID NO: 42:

(i) SEQUENCE CHARACTERISTICS:

20

- (A) LENGTH: 1536 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 42:

GGCACAGGCC AACTTAGTATT GAGTTCTTCT TCTGGACTCT GTATGTCTTT GTGTGTACCC 60
 TATGCCGTTT ACAGTCCGTA CTCTCTCTGT GARATTGGCT GTCTAATCCA GGTGGATCAG 120
 30 GAGGTGCTTT GTGGTTTTTT TGCAAAGAAA TGAAGTCTGG CAAGCAAACA ATGATTAAAC 180
 ATGTTTCGAT TCGTGACTTG TCTTTTGGCG AAATGCAAAG GTGGGTGTGC ATTCTTGAAT 240
 TCAAAGAAAA TCTCTTTCAA ATCCCCCTCAT CCCTTGTGTC TCTTCTAAAT ACTCTCTTTC 300
 35 TAGATATCTT GCACCCCCAA AACTCCCTCA GCCCCCATGG CAGCTTTTCT CTCTCCTCTC 360
 TCTCTTTCCC GCCTCTCCCT GTCTCCTCAC TTCAGCCTTT CCTCTTTCTT AGATCTTTAT 420
 40 TATGTAGATA AAAACCCCTC CAACCTCCTT AGCCTTCTCT CCATTGCATC CCCTACCCGA 480
 ATTATCTTCA AGAAGAGGCG CAGGATCCGA CACAGCGATC AGAAATCTTC CTCCCTTASA 540
 AGCSCAGGGG TGAGGGAGTT CAGGAATATT CATACACTGG TAATCCTTGT CCCTGTTACA 600
 45 GTCACTTCCT TGTATCAGGA CCCTTGTAC TATTTACAGA CTATTTTCCA TCTCTCTTAA 660
 TGCAATTGCT CAAAGGGCAC TTTAAGNATA ATCATTATCC ATTGATGTTT TTTGGAGGCT 720
 50 TTTATTCCTT CCAATAAGTT CTGCCGAATA CTGGCCGCTG GCTCTATTTG TTAAACAATG 780
 GAGGCTTTG TTCCGCTTTT TTTTTTTTTT TTWTCWTAA CCTGAGCTTT CTGCCACCC 840
 TTAGTATGGG GCCAAAGGGA AGATTTTTAT GCCACCCCTT TTGGTGAGAA GAGTCACTTC 900
 55 CTGATTAGTG TTTGGGCTGA AAATGGGTCC CCCTTTGGGA AGAAACATGG GTGCAGTGTA 960
 CTTCTGTGT CACAGGATTA ACAGCTCCTG CCCCCTCCC AAGGAGGCAG CTCYTCGGGG 1020
 60 CAGTTCYTCT TTGAGAATTT CATGGTCATT AAGAAGCAGG YTCCAGGGA CCCCAGAGTG 1080

5 GGAACCTTTG ACTGAAGTCA CCACAGTGGG TGTAAGATAA ACATAAGAGA CTTTCTCAG 1140
 GGAAGATTTG GAACGAAGAA AAAGAGTAAA AAGTTCACAT GGAACCATGA GTGTINTGGA 1200
 AAAGGGCCCA GAAAGGAAG CTGTGGCTAA GAAGATAAAC TGCCTGATTG CAGAGACCCA 1260
 GGAGAGGGGA TGAATCTCTT TTGTCTGGTC ACATTTCTCW WTAATGATKY TCCACATGTA 1320
 10 CAAAGCTAGC CAGTTTACCA AGTGCTTCCA CACACATTGC TTCATTCTGT GTCTCTTAAG 1380
 CAGATIGACT CCTTGGAAAA GCCTCACGTC TGGCATCTG CACCTGCCCC TCACCAGTTT 1440
 GGCTTGGTC TGCTGGGCTG GTTGGGTCTC CCCATGGTGA GCTCCCATGG TATCTCCTCT 1500
 15 TCACCTTTAT ATCACTCATT AGACACCGGT GACAAC 1536

20

(2) INFORMATION FOR SEQ ID NO: 43:

25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 2541 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 43:

AATTGGGCAC GAGGTTCCTG GCCAACCTGC TGCTGGAGGA GGATAACAAG TTTTGTGCAG 60
 ATTGCCAGTC TAAAGGGCCG CGATGGGCCT CTGGAACAT TGGTGTGTTT ATCTGCATTTC 120
 35 GATGTGCTSG AATCCACAGG AATCTGGGGG TGCACATATC CAGGGTAAAG TCAGTTAACC 180
 TCGACCAGTG GACTCAAGTA CAGATTCACT GCATGCAAGW GATGGGAAAT GGAAAGGCAA 240
 ACCGACTTTA TGAAGCCTAT CTTCCTGAGA CCTTTCGGCG ACCTCAGATA GACCCAGCTG 300
 40 TTGAAGGATT TATTCGAGAC AAWTATGAGA AGAAGAAATA CATGGACCGA AGTCTGGGAC 360
 ATCAATGCCT TTAGGAAAGA AAAAGATGAC AAGTGAAAA GAGGGAGCGA ACCAGTTCCA 420
 45 GAAAAAAAT TGGAACCTGT TGTTTTGTAG AAGGTGAAAA TGCCACAGAA AAAAGAAGAC 480
 CCACAGCTAC CTCGGAAAAG CTCCTCGAAA TCCACAGCGC CTGTCTATGA TTTGTTGGGC 540
 CTTGATGCTC CTGTGGCCTG CTCCATTGCA AATAGTAAGA CCAGCAATAC CCTAGAGAAG 600
 50 GATTTAGATC TGTGGCCTC TGTTCATCC CCTTCTTCTT CGGGTTCCAG AAAGGTTGTA 660
 GGTTCATGC CAACTGCAGG GAGTGCCGGC TCTGTCTCTG AAAATCTGAA CCTGTTTCCG 720
 55 GAGCCAGGGA GCAATCAGA AGAAATAGGC AAGAAACAGC TCTCTAAAGA CTCCATCTTT 780
 TCACTGTATG GATCCAGAC GCYTCAAATG CCTACTCAAG CAATGTTTAT GGCTCCCGCT 840
 60 CAGATGGCAT ATCCACAGC CTACCCAGC TTCCCGGGG TTACACCTCC TAACAGCATA 900

	ATGGGGAGCA TGATGCTCC ACCAGTAGGC ATGGTTGCTC AGCCAGGAGC TTCTGGGATG	960
	GTTGCCCCCA TGGCCATGCC TGCAGGCTAT ATGGGTGGCA TGCAGGCATC AATGATGGGT	1020
5	GTGCCGAATG GAATGATGAC CACCCAGCAG GCTGGCTACA TGGCAGGCAT GGCAGCTATG	1080
	CCCCAGACTG TGTATGGGGT CCAGCCAGCT CAGCAGCTGC AATGGAACCT TACTCAGATG	1140
10	ACCCAGCAGA TGGCTGGGAT GAACTTCTAT GGAGCCAATG GCATGATGAA CTATGGACAG	1200
	TCAATGAGTG GCGGAAATGG ACAGGCAGCA AATCAGACTC TCAGTCCTCA GATGTGGAAG	1260
	TAAAAACAAA ACACCTGTAT GGCTGCCATT CTCCTCAGCC CTCGCTCTCC CCTTTCCACA	1320
15	GOCTCCACCC CTGACCCCCA TCCTCTTTTC CTACCTCTCT GTTTGGTTTA GAAATTGCTC	1380
	AATAAGTCAT TTGGGGTTTG GCATCCTGCC CAGCCACTTC CCAAACATGA AGACCTCTCT	1440
20	GTGTCTTTAT GTGTACATG CCCCATAGCC ATCCCAACGT CCTCCCCAGT CCTCTCCTGG	1500
	CACCAGCACC TTAGAAGTTG TTGGCAGAAG GCACTTAAAC TGTGGGAGAA GTGTGCACAC	1560
	CTTTGAGTCC CTTCCTCAA GGTAAAGCT CCTGTCAGAC TCTCAGAAGG GTCTGTGGGT	1620
25	GTGTATATAT AGGCAAACAG GGGAAAGCTT AGAGGTCTTT CTATATGTGT TAATAAGCTG	1680
	TTTCTAAGTG TTAAATTTG AAAAGCATCA TGTCTCATG ATTTATGGGA ATGAAGCAAG	1740
30	TACTGAAATC AAATTAATA CTCCTGGGT CCTGGGTCAG TTTGACCCTA GCCCTGGGGT	1800
	GAGGCAAGCC CCTCCTATG AGGATGAGCA AAAATACTAC TCTCTTCCCC CTGAGTTGCT	1860
	TTCTGGATCT GGGGCTTCAG GACTTGCTGC TTCAGTCAGC CTTTATTAGC ACCAAAGACT	1920
35	TTATGAAGAT CCCACACACA GACACACATC CCTTCCCGCC TCCCCCTGCG CTTCAGTAGG	1980
	ATCTGGCTCC GTGGCTGGAG GACCAACCCC TATAGTGGGA ATGCAGAGCT TAACGTGTAC	2040
40	TGCTTGTTGT TGTGCTGAG TGTGTGTGTG TGTATGAGTG TGTGTTCCGC CTCCCACCT	2100
	CTCCCCATCT GCTCTGGGTA TTTTGTGTTT TGTTAGTTT TAGGTTTACA ACAGAGAGGA	2160
	ATTAATTTAT CAGCAGCCTA AAAGTGTGT GTTTTCTTA TGGTTTAAAA AACGCCATGT	2220
45	CATTGATAAC TCCCTTCTC CCTTCCCTTC TCCCGGTCTG CTGATCACTC TTTTCATGCT	2280
	GTGTATCCAG GGTGCTCTGT TTCCCCACCG TTCCCAGGTG TACGAGGCAG AGGGCCGGGA	2340
50	CAGCTTTCCT CTCAGTCATT GTTCACCCCA CTGAAAATT CAGACAAGAA AACITTGCTT	2400
	AAAAGATTTT ATGTGTGGGA ACCACAGTTC CTGGCTGCCT TTCTCCTGTG TATGTGTAAG	2460
	TTCTTAAATA AATATTGCAG GGAAGGACAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA	2520
55	AAAAAAAAAA AAAAAACTCG A	2541

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2418 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 44:

10 CCCACGGTC CGCCACGCG TCCGCCACG CGTCCGCCA CGCGTCCGGG ACTCAGCGAA 60
 GGGTGGGCGC CGCCGAGGCC TCCTGCCGCT GGCGGGTTTC CGCGGAGTGC CGCCCGGCTC 120
 CGCTCTGCCG CGGCGCGGC TCATGGGCAG AGTCGGCCGG GCGGGCCGGC ATTAACTGA 180
 15 AGAAAAGATG TCCCTGTACG ATGACCTAGG AGTGGAGACC AGTGACTCAA AAACAGAAGG 240
 CTGGTCCAAA AACTTCAAAC TTCTGCAGTC TCAGCTTCAG GTGAAGAAGG CAGCTCTCAC 300
 20 TCAGGCAAAG AGCCAAAGGA CGAAACAAAG TACAGTCCTC GCCCCAGTCA TTGACCTGAA 360
 GCGAGGTGGC TCCTCAGATG ACCGGCAAAT TGTGGACACT CCACCGCATG TAGCAGCTGG 420
 GCTGAAGGAT CCTGTTCCTA GTGGGTTCCT TGCAGGGGAA GTTCTGATTC CCTTAGCTGA 480
 25 CGAATATGAC CCTATGTTTC CTAATGATTA TGAGAAAGTA GTGAAGCGCG CAAAGAGAGG 540
 AACGACAGAG ACACGGGGAG TGGANAAGAC AAAAGGAAAT AGAAGAAAGG GAAAAAAGGC 600
 30 GTAAAGACAG ACATGAAGCA AGTGGGTTTG CAAGGAGACC AGATCCAGAT TCTGATGAAG 660
 ATGAAGATTA TGAGCGAGAG AGGAGGAAAA GAAGTATGGG CGGACTGCCA TTGCCCCACC 720
 CACTTCTCTG GTAGAGAAAG ACAAAGAGTT ACCCCGAGAT TTTCCTTATG AAGAGGACTC 780
 35 AAGACCTCGA TCACAGTCTT CCAAAGCAGC CATTCCTCCC CCAGTGTACG AGGAACAAGA 840
 CAGACCGAGA TCTCCAACCG GACCTAGCAA CTCCTTCCTC GCTAACATGG GGGGCACGGT 900
 40 GGCGCACAAG ATCATGCAGA AGTACGGCTT CCGGGAGGGC CAGGGTCTGG GGAAGCATGA 960
 GCAGGGCCTG AGCACTGCCT TGTCACTGGA GAAGACCAGC AAGCGTGGCG GCAAGATCAT 1020
 CGTGGGCGAC GCCACAGAGA AAGGTGTGTC CCCAGGGAAG CGTGTGACTA GAGGGAAAGG 1080
 45 ACTGGCCCCA TCCATATCAG ACATGGCCAG TCTTGATCCT CATGTGTCAG CAGGGGGACA 1140
 ATGAGGCGTG TGGCCAGAGG GAGAGGGCTG GCCCTGCCAT CACTAGAACA CAGGCCGTCC 1200
 50 TGTTCATATG ATGCACTGCC ACTTCCGTTT TGTGAAACCA GGAATCCTGA GGCTCATCTT 1260
 TATTTTTTCA GAACAGACGT AGAGAGATGA AGGCTTGTGG AGGAAAAGAT GGTGAGAGAC 1320
 TTGGGCAGAA AATGAGTAGT CCTCAGGAAG AAATCTTGGT TATGTGTTTA GAGCATGAAG 1380
 55 GACAGAGCCA TATAGTGTGG CAGTGAATAT ACCTGCTATC TCCATCTCAG AGGTGCTCTC 1440
 TACTTTTCCC TTTTGCCCTT TCAGTATAGA TGTGATTCTT GATTCTCTTA CAGATTGTTT 1500
 60 GCTTTGCGAG ATCTGATGTT ATGTTGCAGT CTCTTGSTAA ATGATGCCTA GTTGGTGTTC 1560

5 TATTTTCATT TAATTTTAC AGTCTGTTCT GTGTTGAGGG AATTCAGGAA AGAGACAAAC 1620
 ATATGTTAGC ATTTTAATCA GGGAAITTAAG TTTGAGTCAG CCTAGCTGAA CTTCTTTTGC 1680
 TAAAGAAAGA AGAAAACTTT TCTGGCAGCC CCGTTCATGC ACAGCTTAGG GATACATCAC 1740
 GAGCCTGACA GATGCATCCA AGAAGTCAGA TTCAAATCCG CTGACTGAAA TACTTAAGTG 1800
 10 TCCTACTAAA GTGGTCTTAC TAAGGAACAT GGTGTTGCGG GGAGAGGTGG ATGAAGACTT 1860
 GGNAAGTTGA AACCAAGGAA GAATGTGAAA AATATGGCAA AGTTGGAAAA TGTGTGATAT 1920
 TTGAAATTCC TGGTGCCCOCT GATGATGAAG CAGTACGGAT ATTTTITAGAA TTTGAGAGAG 1980
 15 TTGAATCAGC AATTAAAGCG GTTGTGACT TGAATGGGAG GTATTTTGGT GGACGGGTGG 2040
 TAAAAGCATG TTTCTACAAT TTGGACAAAT TCAGGGTCTT GGATTTGGCA GAACAAGTTT 2100
 20 GATTTTAAAGA ACTAGAGCAC GAGTCATCTC CCGTGATCCT TAAATGAACT GCAGGCTGAG 2160
 AAAAGAAGGA AAAAGGTCAC AGCCTCCATG GCTGTTGCAT ACCAAGACTC TTGGAAGGAC 2220
 TTCTAAGATA TATGTTGATT GATCCCTTTT TTATTTTGTG GTTTTTTAAT ATAGTATAAA 2280
 25 AATCCTTTTA AAAAAACAAC AATCTGTGTG CCTCTCTGGT TGTTCCTCTT TTTTATTATT 2340
 ACTCCTGAGT TGATGACATT TTTTGTTAGA TTTCATGGTA ATTCTCAAGT GCTTCAATGA 2400
 30 TGCAGCATT TTTGCACT 2418

35 (2) INFORMATION FOR SEQ ID NO: 45:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1337 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 45:

45 TCGACCCACG CGTCCGGAGC GACCTCTCTG CTCCGCTCGT CTCGTGGTT CCGGAGGTCG 60
 CTGCGGCGGT GGGAAATGCT GGC GCGCGCG GCGCGGGCA CTGGGGCCCT TTTGCTGAGG 120
 GGCTCTCTAC TGGCTTCTGG CCGCGCTCCG GCGCGGCCT CCTCTGGATT GCCCGAAAC 180
 50 ACCGTGGTAC TGTTCGTGCC GCAGCAGGAG GCCTGGGTGG TGGAGCGAAT GGGCCGATTC 240
 CACCGGATCC TGGAGCCTGG TTTGAACATC CTCATCCCTG TGTAGACCG GATCCGATAT 300
 55 GTGCAGAGTC TCAAGGAAAT TGTATCAAC GTGCTGAGC AGTCGGCTGT GACTCTCGAC 360
 AATGTAATC TGCAAATCGA TGGAGTCCTT TACCTGCGCA TCATGGACCC TTACAAGSCA 420
 AGCTACGGTG TGGAGGACCC TGAGTATGCC GTCACCCAGC TAGCTCAAAC AACCATGAGA 480
 60

	TCAGAGCTCG GCAAACCTCTC TCTGGACAAA GTCTTCCGGG AACGGGAGTC CCTGAATGCC	540
	AGCATTTGTGG ATGCCATCAA CCAAGCTGCT GACTGCTGGG GTATCCGCTG CCTCCGTTAT	600
5	GAGATCAAGG ATATCCATGT GCCACCCCGG GTGAAAGAGT CTATGCAGAT GCAGGTGGAG	660
	GCAGAGCGGC GGAAACGGGC CACAGTTCTA GAGTCTGAGG GGACCCGAGA GTCGGCCATC	720
	AATGTGGCAG AAGGGAAGAA ACAGGCCAG ATCCTGGCCT CCGAAGCAGA AAAGGCTGAA	780
10	CAGATAAATC AGGCAGCAGG AGAGGCCAGT GCAGTTCTGG CGAAGGCCAA GGCTAAAGCT	840
	GAAGCTATTC GAATCCTGGC TGCAGCTCTG ACACAACATA ATGGAGATGC AGCAGCTTCA	900
15	CTGACTGTGG CCGAGCAGTA TGTCAGCGCG TTCTCCAAAC TGGCCAAGGA CTCCAACACT	960
	ATCCTACTGC CCTCCAACCC TGGCGATGTC ACCAGCATGG TGGCTCAGGC CATGGGTGTA	1020
	TATGGAGCCC TCACCAAAGC CCCAGTGCCA GGGACTCCAG ACTCACTCTC CAGTGGGAGC	1080
20	AGCAGAGATG TCCAGGTAC AGATGCAAGT CTTGATGAGG AACTTGATCG AGTCAAGATG	1140
	AGTTAGTGA GCTGGGCTTG GCCAGGGAGT CTGGGGACAA GGAAGCAGAT TTTCTGATT	1200
25	CTGGCTCTAG CTTCCCTGCC AAGATTTTGG TTTTATTTTT TTTATTTGAA CTTTAGTCGT	1260
	GTAATAAACT CACCACTGGC AAACCAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA	1320
30	AAAAAAAAAA AAAANN	1337

(2) INFORMATION FOR SEQ ID NO: 46:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1276 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 46:

45	CTCACGCGTC CGGGACGGCN GGACGCGTGG GTGCATTTGC TGAGTGTITT ACTTCCAATT	60
	ATGTGATTCN ATATTACAGG NGCTGCCATG TGGTAATGAG AAGAATGTAT ATTCTGTTGT	120
	TTTGGGGTGG ARTGTTCAT AGATGTCTAT CARGTCTGTT TGATCCAGAR CTGARITCAR	180
50	GTCTGGTAT CTCARTCTTT ACTGTGARTC TTCAAATGAC ATAAGAATGA CAGAAMTTGT	240
	AGTTAAGGAC AACAGRGCAW TSCAAGGCAG CAGCATAGTC CAAATAGAC GTGTCTTCTT	300
55	CCCGAAGTCA CTGTAGTGGG GGACATAAAA TTTAAGGAAC CTCTGGGTCT TACTACCTGA	360
	TGTGGCCAAT TGGACTAAAA CCAATAACCA TTAAGGAAWA AATSSACTWA ACCACAAGCA	420
	ACTCAATTAA MAAATAGGCA AAGAACTTGA AGAGGCATTT TCCCAAAGAA GCCAACAAGC	480
60	ATGTGAAAAG ATGCTCAACA TCATTAGACA TCAGGGAAAT ACAGATCAAA ATCAAAATGA	540

5 GATACCAGTT TATACTAAGG TGGCTATAAT AAACATCATA ATAATGAAGG ACATTAACAT 600
 GTATTAGTGA GGATGTGGAG AAATGGAACC CATTCTGGT AGGAATGTAA AATAGTGCAG 660
 CCACTGTGGA AAACAGTTTG GTGGTTCCCC AGAAAGCTAA GCATAGAGTT ACCAGAGAAC 720
 CTAGCAATTT AACTTATAGG TACATACTTC AAAGGAATTG AAAACATAGA TYCTAACAGA 780
 10 TACTKGTACA GCAATATYCA TKGTTGCWTT ATTACAGATA GCCAAAAGGT AAAACAACTC 840
 AAGTGTCAT CAAATATATA ATGTGTAAAC AATGTGGTAT ATTCCTAGAG GGAATATTA 900
 15 TTCAGCTTTA AAAAGGAATG AAGTACTGGT ACATGCTACA AAGGTGGATG AGCCTCAGAA 960
 ACATGCTGAG TGAAAGAAGC CAATGATAAA AGACCATATA TTGTATGATT CCATTATATG 1020
 AAATKTCAG RACATTCAAG TCTATAGAGA CAGAAAGTAG ATTAGTGAYT GCTTAGGGCT 1080
 20 GGCAGGATA AGGGGKTCAT GGCTAAAGGG TATGGGTTT TGTGTGGA GGTGAAAAAT 1140
 TTTAAACTT GKSTGATGG TTGCACAAGC CTGTGAAGAT ACTGAAAACC ATTGAATTGT 1200
 25 GTGCTTTAAA TGGATGAATT GTATGGTGT TGAACATATAT CCAATAAAG CTGTTTTTTA 1260
 AAAAAGAAAA AAAAAA 1276

30

(2) INFORMATION FOR SEQ ID NO: 47:

35 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1282 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 47:

GGCACGAGAG AAAGGCCAGT TTGTGGGCA AATTAGACTA AACTCTGTGC TGGTAGAACT 60
 GCTTTCCAAG AATGCTGTCA CTGCTATAGT TTTAATGCT TCAAATCTCA ACTCNCCTCCC 120
 45 TCCATTGCGC ATAGCTCAAC CATGTTCCAG GAGTGTATTC CAATCAGCTT GTTTTCTCTT 180
 AACTGGTCAA AGGAATGTTG CTCATTCAAC TGCCCCAACT CACATATTAA CAATTGTTTA 240
 ACTGGGATTA GATAAAAGGA AAGCTGACTT ACAGATGAAC CAAGAGGGAG CTATTTATGC 300
 50 CACAGCCCCC AGCCAGTAA CTTTATGTTT CTGATCTCCT GCAAAATTTT TTTATAAAAA 360
 AAGCTTAGCC AGGAAGTAGT AGAAAGAATA AAGTAAAGAT GGTGTAAGAA ATATATGGAT 420
 55 AGGCAAGTTC CWNVGTGAG ACCTTAYGAA GAATGGTGAG GTGTGGTTAA ATGGAGGAGA 480
 TAATCAGCAG ATAAGAGCTC AGATGGTCMS AAACATWTAG AACTATAATG CCATCTCCAA 540
 AGTATTGCAT GCATACAAAT GACGTCAAT CCGTTGAATA TAATGGAGAC AACTATTTTC 600

60

5 AAAAAATTAAAG TTCTTCTWTC TTGAGCTTTA AAAGTATACA CATTTACCCM AATGAATTWA 660
 AAACATGCMC ACMAATATTT ATATCAAAAG TGTACATGAT TTCCAAAAC TGAAGTWAC 720
 10 CAAGATTTAC TTCCWTGGGT TAGTGCATAA ATTAACGTGT ATACATATAT ACTATGGAAT 780
 WITAYTCAGC AACAGAAATA AATGAGHTAT CAAACCACAG AAAGACATGG AGGAAACTTA 840
 AATCCAGGTG GMITAAGTGAW AGAAGCCAAT ATGAAAAGGC TACATTSTAT ATGATTTCAA 900
 15 ATATATGACA TTCAGGAAAA GGCAAGGCTG CAGAGACAGT AAARAGATCA GCTAGGTGCA 960
 TGKGGSTCAC GCCACTTTGG GAGGCTTGAG GCAGGKGGAT TATMTGAAG TCAGGAGTTC 1020
 NAGACCAGCN TGGGCAACAT GNTGANACCC CATATNTCCT AAAAGNACNA AAATTTAACT 1080
 GGGCGTGGTG GCACGTGCCT GTANTCCAN CNACTCTGGT GGCTNAGACN GGNGAATTGC 1140
 20 TTGAACCCAG GAGGCAGAGG TTGCGGTGAG CCAATGATTG CACCCTGCA NTCCAGCCTG 1200
 GGTGGTAGAG CGAGACTCAG TCTCAACNTT NATCAAGATA GGANNGAAAT AGAANGGAAG 1260
 AAAGAGAAAA AATAAAAAATA NA 1282

25

(2) INFORMATION FOR SEQ ID NO: 48:

30

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 645 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 48:

AAGGTAGAAA AGTACAGAAA AACTAAATT TTCATTGTGC TGTTTCAATG TGGCAGATTC 60
 40 TTTAAATAC TTCGACACGC TACAATAATT AAAGGTTTGA AGAACATTAA GATACTTAAA 120
 AAATAAAAGC CCACAATTGA ATAACAAAAA TGAACCTTGT TTTATTTTAT ATTGGCATTA 180
 45 ATGTAGGTG CCGTGGTGAA AATAGTTTGA AATACTTCAC AGTAACAGTT TTGTGCAGCC 240
 CTAGAGATTA AAAACAGCAA AGTAAATAAG CAGGACTCTC AACGACTCAT ACTCACAGAC 300
 ATGTTTAATG TAATCCTAGC ACTTCGGGAG GCTGAGGCGG GAGGATTACT TGAGCCTAGG 360
 50 AGTTTGAGAC CAGCCTGGGC AACATAGCAA GATCCCATCT CTACAAAAAA GTGAAAAAGT 420
 TAGCTGAACA AGGCGGCATG CACATGCTAC TCCAGACGCT GAAGTGGGAA GATCACTTAA 480
 55 GTCCGAGAGA TCGAGGCTTC AGTGAGATAT GGCTGAGACA CTGCTCTCAG CCTGGATGAC 540
 AGAGTGAGAA CCTGTCTCAA ACAAGAGAAA AAAATAAATC AAATGCTATT CAAAATCTA 600
 AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAA 645

60

(2) INFORMATION FOR SEQ ID NO: 49:

5

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1495 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 49:

TGTGGAAAAC AGTAGGAAAG CAATGAAAGA AGCTGGTAAG GGAGGCGTCG CTGATTCCAG 60
 15 AGAGCTAAAG CCGATGGTAG GTGGAGATGA GGAGGTGGCC GCCCTCCAAG AATTTCACCT 120
 TCACTTCCTC TCTCTCTCTG TCTTCACTGA CTGCACTTCT TCAGGAGAAG CTTTGTGTAT 180
 CTGTATCACG CAGACATGCT GCTCTTTCTG TTTGTGTGCT TACCCATCAC TTGGATGGCA 240
 20 GAATTCCTGT CACAACCTGAG ACCACCTTCT ATAAAAGTAA GCTGAAAGGA ACAGCATCCT 300
 CGTCAGTGCT CGGCAGGGG GGTAGGGGA TGATGGTTTT TTCCTAAGG TAAAACTGCT 360
 25 GTTGCTCTTG TTTCCTTTTT AACTGTCAGT GTTTGGCTTT CATCAGAMTG AACATTTTGG 420
 TGTTCACCTT GAACTGACGG TTTGATTTTT ATCATTTTGG AAAGGTGATC ATAGCAATTC 480
 CTTTCCAAC TGTCTAAAAT CCATACTCCC CCTTTTAAA ARWATKGTTS TGCCTTCATT 540
 30 GCTKIMCWIT TSCCTTGKCT SMCITTTTCY TCCTGTGKSC TGAARTTKTW CYTTCYTKT 600
 TTCTTAAGST WTTTTTCAGT AGCAAACAAG GCTGTTTTCA TCAATACCCA CATTCCCAYT 660
 35 CRGKRRGRMM ATYTAGTYTT YTCCCAGKTT AAKTGKGRGR KGRKGAAAA TRATKTCKGG 720
 KANGKGGAWA TKAWAWAKG KWWATGKAAA CACAAATATA TYTYTYTAMA TTCCACITTA 780
 ATTKGGGAAA AAAGGCAGCT KAAGTGGAGT GTWAAGRARR ACCTKGRST GCTTTTCAAC 840
 40 ATGGGATATG GTCACATATG CATRGGAAC ANGATGCCTT CTATCAWAKA TGGGTCTAAT 900
 TACTYCCTAA TTAAACACAC GTATTTTTTT AAATAGCATG TTTATTTTCA AATATDATAT 960
 45 AATGGTCGSG CRTCTTAAA TAATTTTAAA CAANGTGTCC CCGRGACNGC ATATAATGTT 1020
 CAAAWGKAG AGGTAAGGAC TTYCCTTTCT GTCTYCTTAA CACTTWAGTA AATRATTNGA 1080
 WITAWAGCAA GTTGTGCCAA CTGKCNNCCT GNGGNCCGCA NANGGMWGRG GAAGGGCTTT 1140
 50 TCMAACACAA ATTCGTAAAC TTTATTAAAA CATGAGATTT TTGCTTTT TTTTTTAAG 1200
 CCCATCAGCT ATCCTTAATG TATTTTANAT GTGGCCCAAG ACAATTCCTC TTCCAGGATG 1260
 55 GCCTGGGGAA GCCAAAAGAT TGGANACCCC TGATTTGTAG GTTTTCAACT TTAAATATA 1320
 TGCTATAAAA TAAGTTCATT TAAGTAGGCT AGGCATGGTG GCTCATGTNT GTAATCCTAG 1380
 CACTTAGGGG GCCCGAGGCA GAAAGATTRM CTGAGCTCAG CAGTTTGAGA CCAGCCTGGG 1440
 60

CCAAACGGTG NAACCTGTT TTTACTNAAA TACCCAAAAA AAAAAAAAAA AAAAA 1495

5

(2) INFORMATION FOR SEQ ID NO: 50:

(i) SEQUENCE CHARACTERISTICS:

10

(A) LENGTH: 1630 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 50:

15

GAATTCGGCA CGAGATTATC TGTCTTCTTC TTACCAATTT ATAGAACTTT TTAGTATTGC 60

AGATAAAGTT CCTCATCGGA TATCTTCTCT CCTTCTATTG GGTACCTTTT TATTGTCTTA 120

20

ATGGGGGTCT TTTAATGACC AGAAGTTCTT AGTTTAAAA TAGTCCAGTT TATCCATTTT 180

TAAATGTGTA GTGCTATTTG TGTCTGCTT GAGAGATTTT TGCCTACTGC AAGGTCACAA 240

25

AGATGTTTTT CTCTAAAAGC CTTTTGGTTT TGCCCTTTTG TTTTAGATCT GCAGCTCATC 300

TGGAATTGAG TGTGTGGTGT GTGTGTGGTG TGAGGTAGGG GTCCTTTTTT TCATATGGAT 360

ATCCAATTGA CCCAGAACAG TGTATTGAAA AAAAAAATCT GTCTTAGTCA ATTTGGACTG 420

30

CCGTAACAAA ATACCATAAC CTGGGTGGCT TAGACTACAG AAATGTAGCG CTCACAGYTC 480

TGGAGGCTGG AAGGCCAGGA TCAAGACACC AGCAGATTCG GTGTCTNGTG AGGACCCACT 540

TTGTGMITCA TAGATGTCAC CTTCTTGCTG TGTCCAGTG GTGRAAGGGG CAAACTAGCT 600

35

CCCTTAAACC TCTTTTATA AGATCCCTAA AACCTTTAAT GAGGGCTCCA CCCTAATGAT 660

CTAATCACCT CTCATACCT TATCTTGGGG GTTAAGATTT GAACAGAGGA ATTTGGGGGA 720

40

GACATAGACA TTTGGAGCAT AGCATCTTCT TTTCTCAGT GCACAGCAGT GCTGCCITCA 780

TCATCAGTCA GGTGTCGTA GGTGTGTGGC TATTTCTGGA CTTGGCACTC TGTCTACTTT 840

GTGATTTTCT CTGCCITATA CCAATGCCAC ACCATCTTAA TTATTGTAAC CATCTTAATT 900

45

ATTTATAAAA AGTCTTTTTT TTTTTTTTGA TACAGTCTCA CTCTGTCCCC CAGGCTGGAG 960

TGCAGAGGTA CAGTATTGGC TCACTGCAAC CTCTGTCCCC AGGCTTAAGC AATTCTCATG 1020

50

CCTCAGCCTC CTGAGTAGCT GGGATTACAT GTGCACCACC ACACCTGGCC TTCTTTCTTT 1080

TCTTTCCAAY CCATTGTTTT TTTATTTCTT TCCCTKGCTT TATKGCACGT GCTAAGATTT 1140

CCAGTGCTGA ATAGGAGTGA TGACAGTGGG CACCTTGTC TTTCTCCCAA CCTCAGAGGG 1200

55

AAAAGTATCC AATGCATTTG TAGATAITCT TTATCAGATT AGCTTCCTTT CTAGCGGCTT 1260

GTGTCTTTC ATTGTTTTTC ATGAGCAAGT GTTGAACTTT TTCCTGAGT TTCCAAATA 1320

60

CTTTTCCAT TGAGTTTTTT TACTTTAACC GTCATATTGC CAAAAGTCTG CATTTGTTAT 1380

TTCTCCCAA ATGCTGGGA TTATAGGCAT TAGCCACTGC ACCCAGCCAG ACTTTATAGA 1440
 AAATCTTGAT ATCTGGTCAT GGAAGTCCCC TAGCTTGGTT ATTTTTTTTTT GGTACCGCTT 1500
 5 TGTCTATTTT CGGCCCTTTC CATTTCATG TAACTTTTAG GATCAGCTTG TCAGTTCCTA 1560
 CCAAAAAAAA AAAAAAAA ACTCGAGGGG GGCCCGGTAC CCAAATCGCC GGGTAGTGAT 1620
 10 CGTAACAATC 1630

15 (2) INFORMATION FOR SEQ ID NO: 51:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2420 base pairs

(B) TYPE: nucleic acid

20 (C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 51:

25 GCCAACAGTG CTCCCTCATA GATGGACGAA GTGTGACCCC CCTTCAGGCT TCAGGGGGAC 60
 TGGTCTCCT GGAGGGAGAT GCTCGCCTTG GGAATAATC ACTTTATTGG TTTTGTGAAT 120
 GATTCTGTGA CTAAGTCTAT TGTGGCTTTC CGCTTAATC TGGTGGTGAA GGTCAGCACG 180
 30 WGGCCGGGGG AGAGTCACGC AAATGACTTG GAGTGITCAG GAAAAGGAAA ATGCACCACG 240
 AAGCCGTCAG AGGCAACTTT TTCCTGTACC TGTGAGGAGC AGTACGTGGG TACTTTCTGT 300
 35 GAAGAATACG ATGCTTGCCA GAGGAAACCT TGCCAAAACA ACGCGAGCTG TATTGATGCA 360
 AATGAAAAGC AAGATGGGAG CAATTTCACC TGTGTTTGCC TTCCTGGTTA TACTGGAGAG 420
 CTTTGCCAGT CCAAGATTGA TTAGTGCATC CTAGACCCAT GCAGAAATGG AGCAACATGC 480
 40 ATTTCCAGTC TCAGTGGATT CACCTGCCAG TGTCCAGAAG GATACTTCGG ATCTGCTTGT 540
 GAAGAAAAGG TGGACCCCTG CGCCTCGTCT CCGTGCCAGA ACAACGGCAC CTGCTATGTG 600
 45 GACGGGGTAC ACTTTACCTG CAACTGCAGC CCGGGCTTCA CAGGGCCGAC CTGTGCCCAG 660
 CTTATTGACT TCTGTGCCCT CAGCCCCTGT GCTCATGGCA CGTGCCGAG CGTGGGCACC 720
 AGCTACAAAT GCCTCTGTGA TCCAGGTTAC CATGGCCTCT ACTGTGAGGA GGAATATAAT 780
 50 GAGTGCCTCT CGCTCCATG CCTGAATGCA GCCACCTGCA GGGACCTCGT TAATGGCTAT 840
 GAGTGTGTGT GCCTGGCAGA ATACAAAGGA ACACACTGTG AATTGTACAA GGATCCCTGC 900
 55 GCTAACGTCA GCTGTCTGAA CGGAGCCACC TGTGACAGCG ACGGCCTGAA TGGCACGTGC 960
 ATCTGTGCAC CCGGGTTTAC AGGTGAAGAG TGCGACATTG ACATAAATGA ATGTGACAGT 1020
 AACCCTGCC ACCATGGTGG GAGCTGCCTG GACCAGCCCA ATGGTTATAA CTSCCACTGC 1080
 60

	CCGCATGGTT GGGTGGGAGC AAACGTGTGAG ATCCACCTCC AATGGAAGTC CGGGCACATG	1140
	GCGGAGAGCC TCACCAACAT GCCACGGCAC TCCCTCTACA TCATCATTTG AGCCCTCTGC	1200
5	GTGGCCTTCA TCCTTATGCT GATCATCCTG ATCGTGGGGA TTTGCCGCAT CAGCCGCATT	1260
	GAATACCAGG GTTCTTCCAG GCCAGCCTAT RAGGAGTTCT ACAACTGCCG CAGCATCGAC	1320
	AGCGAGTTCA GCAATGCCAT TGCATCCATC CGGCATGCCA GGTTTGGAAA GAAATCCCGG	1380
10	CCTGCAATGT ATGATGTGAG CCCCATCGCC TATGAAGATT ACAGTCTTGA TGACAAACCC	1440
	TTGGTCACAC TGATTAAAC TAAAGATTTG TAATCTTTTT TTGGATTATT TTTCAAAAAG	1500
15	ATGAGATACT ACACCTCATTT AAATATTTTT AAGAAAWTAA AAAGCTTAAG AAATTTAAAA	1560
	TGCTAGCTGC TCAAGAGTTT TCAGTAGAAT ATTTAAGAAC TAATTTTCTG CAGCTTTTAG	1620
	TTTGGAAGAAA ATATTTTAAA AACAAAATTT GTGNAACCTA TAGACGATGT TTTAATGTAC	1680
20	CTTCAGCTCT CTAACACTGTG TGCTTCTACT AGTGTGTGCT CTTTTCACTG TAGACACTAT	1740
	CACGAGACCC AGATTAATTT CTGTGGTTGT TACAGAATAA GTCTAATCAA GGAGAAGTTT	1800
25	CTGTTTGACG TTTGAGTGCC GGCTTTCTGA GTAGAGTTAG GAAAACCACG TAACGTAGCA	1860
	TATGATGTAT AATAGAGTAT ACCCGTTACT TAAAAAGAAG TCTGAAATGT TCGTTTTGTG	1920
	GAAAAGAAAC TAGTTAAATT TACTATTCTT AACCCGAATG AAATTAGCCT TTGCCTTATT	1980
30	CTGTGCATGG GTAAGTAACT TATTTCTGCA CTGTTTGTGT GAACTTTGTG GAAACATTCT	2040
	TTTCAGTTTG TTTTGTGTCAT TTTCGTAAAC GTCTGCGAAC TAGGCCTCAA AAACATACGT	2100
35	AACGAAAAGG CCTAGCGAGG CAAATTCTGA TTGATTTGAA TCTATATTTT TCTTTAAAAA	2160
	GTCAAGGGTT CTATATGTGR AGTAAATTAA ATTTACATTT GAGTTGTTTG TTGCTAAGAG	2220
	GTAGTAAATG TAAGAGAGTA CTGGTTCCTT CAGTAGTGAG TATTTCTCAT AGTGCAGCTT	2280
40	TATTTATCTC CAGGATGTTT TTGTGGCTGT ATTTGATTGA TATGTGCTTC TTCTGATTCT	2340
	TGCTAATTTT CAACCATATT GAATAAATGT GATCAAGTCA AAAAAAAAAA AAAAAAAAAA	2400
45	AACTCGAGGG GGGTCCCGT	2420

50 (2) INFORMATION FOR SEQ ID NO: 52:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1172 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 52:

60 AAAATTATTC TGTACCATCA CAGCTTTTCA CAACGATGGC AAGCCTTATG TCTTGGGAGC 60

	CTGTTTTCCT AGGCAAAGTT ACAAGTGACC TAATGGGAGC TCAAATGTGT GTGTGTCTCT	120
	CTGTGTGTTT GTGTGTGTGT GTGCACTCAA GACCTCTAAC AGCCTCGAAG CCTGGGGTGG	180
5	CATCCCGGCC TTGCCATTAG CATGCCTCAT GCATCATCAG ATGACAAGGA CAACCCTCAT	240
	GACGAAGCAA CATGAATTAG GGGGCCTCTT GGCTTGGTC CAAAATTGTC AATCAGAAAT	300
10	GAACATAAAG GACTCCAGAG CAGTGGGACT GTCTGTCAA AGACTCTGTA TATCTTTTGT	360
	GGATGAGTTT TGTGAGAGAA CAGAGAGACC ATTGTACCTG GCACAAGGGC TSTTCATGAA	420
	AAGGGAGACT TACTGGGAGG TGCAAGACAG TGGCATTTCT CCTCTCCTCT TGCTGCTCAG	480
15	CACAGCCCTG GATTGCAGCC CCGAGGCTGA GACCAGACAA AGCCCGGGAG GCAGAAAGAT	540
	GCTCCAAGAA CCAACACTAT CAATGTCTTT GCAAATCCTC ACAGGATTCC TGTGGGTCCA	600
20	GCTTTGGAAC TGGGAAACCT TTCTTCGGAT CGCACTCAT TCCACTGATG CCAGCTGCCC	660
	CTGAAGGATG CCAGTACTGT GGTGTGTGAG TCTCAGCAGC CGCCACACG CTCCTAACTC	720
	TGCTGCATGG CAGATGCCTA GGTGGAAATA GCAAAAACAA GGCCAGGCT GGGGCCAGGG	780
25	CCAGAGGGGA AGGCCCTGGA TTCTCACTCA TGTGAGATCT TGAATCTCTT TCTTTGTCT	840
	GTTTGTTCAG TTAGTATCAT CTGGTAAAT AGTTAAAAA CAACAAAAA CTCGTATCT	900
30	GTTTCTAGCA TGTGCTGCAT TGAATCTATT AATCACATTT CAAATTCACC CTACATTCCT	960
	CTCCTCTTCA CTAGCCTCTC TGAAGGTGTC CTGGCCAGCC CTGGAGAAGC ACTGGTGTCT	1020
	GCAGACCCCC TCAGTTCTTG TGCCTCAGCC CACAGGCCAC TGTGATAATG GTCTGTTTAG	1080
35	CACCTCTGTA TTTATTGTAA GAATGATTAT AATGAAGATA CACACTRTAA CTACAAGAAA	1140
	TTATAAATGT TTTTCACATC AAAAAAAAAA AA	1172
40		

(2) INFORMATION FOR SEQ ID NO: 53:

- 45 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1589 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear
- 50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 53:

	CCCACGCGTC CGCCACGCG TCCGCCACG CGTCCGTTTC AAAGGGAGCG CACTTCCGCT	60
55	GCCCTTCTTT TCGCCAGCCT TACGGGCCCG AACCTCGTG TGAAGGTGC AGTACCTAAG	120
	CCGAGCGGG GTAGAGCGG GCCGGCACCC CCTTCTGACC TCCAGTGCCG CCGGCCTCAA	180
60	GATCAGACAT GGCCAGAAC TTGAAGGACT TGGCGGGACG GCTGCCCGCC GGGCCCCGGG	240

	GCATGGGCAC GGCCCTGAAG CTGTTGCTGG GGGCCGGCGC CGTGGCCTAC GGTGTGCGCG	300
	AATCTGTGTT CACCGTGGAA GGCGGGCACA GAGCCATCTT CTTCAATCGG ATCGGTGGAG	360
5	TGCAGCAGGA CACTATCCTG GCCGAGGGCC TTCACCTCAG GATCCCTTGG TTCCAGTACC	420
	CCATTATCTA TGACATTGGG GCCAGACCTC GAAAAATCTC CTCCCCTACA GGCTCCAAAG	480
	ACCTACAGAT GGTGAATATC TCCCTGCGAG TGTGTCTCG ACCCAATGCT CAGGAGCTTC	540
10	CTAGCATGTA CCAGCGCTA GGGCTGGACT ACGAGGAACG AGTGTGCGG TCCATTGTCA	600
	ACGAGGTGCT CAAGAGTGTG GTGGCCAAGT TCAATGCCTC ACAGCTGATC ACCCAGCGGG	660
15	CCCAGGTATC CTTGTTGATC CGCCGGGAGC TGACAGAGAG GGCCAAGGAC TTCAGCCTCA	720
	TCCTGGATGA TGTGGCCATC ACAGAGCTGA GCTTTAGCCG AGAGTACACA GCTGCTGTAG	780
	AAGCCAAACA AGTGGCCAG CAGGAGGCC AGCGGGCCMA ATTCTTGSTA GAAAAAGCAA	840
20	AGCAGGAACA GCGGCAGAAA ATTGTGCAGG CCGAGGGTGA GGCCGAGGCT GCCAAGATGC	900
	TTGGAGAAGC ACTGAGCAAG AACCTGGCT ACATCAAAC TCGCAAGATT CGAGCAGCCC	960
25	AGAATATCTC CAAGACGATC GCCACATCAC AGAATCGTAT CTATCTCACA GCTGACAACC	1020
	TTGTGCTGAA CCTACAGGAT GAAAGTTTCA CCAGGGGAAG TGACAGCCTC ATCAAGGGTA	1080
	AGAAATGAGC CTAGTCACCA AGAACTCCAC CCCCAGAGGA AGTGGATCTG CTTCTCCAGT	1140
30	TTTTGAGGAG CCAGCCAGGG GTCCAGCACA GCCCTACCCC GCCCCAGTAT CATGCGATGG	1200
	TCCCCACAC CGGTTCCCTG AACCCCTCTT GGATTAAGGA AGACTGAAGA CTAGCCCCTT	1260
35	TTCTGGGGAA TTACTTTCTT CCTCCCTGTG TTAAC TGGGG CTGTTGGGGA CAGTGCCTGA	1320
	TTTCTCAGTG ATTTCCTACA GTGTTGTTC CTCCCTCAAG GCTGGGAGGA GATAAACACC	1380
	AACCCAGGAA TTCTCAATAA ATTTTATTA CTTAACCTGA AGTCAAGGCT TCACGTGTTT	1440
40	ATGAACTGGG TAACTGGCAG CAAGCATGCG CACGTTTACA TGTGCGCTCC TGGGTCTGTC	1500
	TTTGTGTGTG CCAGCAGGGG GCGCAAAGA ATCTGGCTGG GGCGGCTAAN GGAAGCAAG	1560
45	GCCTGGGCTC CGAAACANGA CCCAACTGG	1589

50 (2) INFORMATION FOR SEQ ID NO: 54:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2074 base pairs

(B) TYPE: nucleic acid

55 (C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 54:

60 CCGCCTGACC GCCCCGGGCT TAAGGGAGCC TGGCTAGGCC GGCAGCCGGA TGGTCCCGCA 60

	GCTCGGGGCC GGCCATGCTT CGCGGTCCGT GCGGCCAGCT TTGGCTCTTT YTCCTGCTGC	120
5	TGCTCCCGGG CGCGCCTGAG CCCCGCGGCG CCTCCAGGCC GTGGGAGGGA ACCGACGAGC	180
	CGGGCTCGGC CTGGGCCTGG CCGGGCTTCC AGCGCCTGCA GGAGCAGCTC AGGGCGGCGG	240
	GTGCCCTCTC CAAGCGGTAC TGGACGCTCT TCAGCTGCCA GGTGTGGCCC GACGACTGTG	300
10	ACGAGGACGA GGARGCAGCC ACGGGGCCCC TGGGCTGGCG CCTTCCTCTG TTGGGCCAGC	360
	GGTACCTGGA CCTCCTGACC ACGTGGTACT GCAGCTTCAA AGACTGCTGC CCTAGAGGGG	420
15	ATTGCAGAAT CTCCAACAAC TTTACAGGCT TAGAGTGGGA CCTGAATGTG CGGCTGCATG	480
	GCCAGCATTT GTCCAGCAG CTGGTCTTAA GAACAGTGAG GGGCTACTTA GAGACGCCCC	540
	AGCCAGAAAA GGCCCTTGCT CTGTCTTCC ACGGCTGGTC TGGCACAGGC AAGAACTTCG	600
20	TGGCACGGAT GCTGGTGGAG AACCTGTATC GGGACGGGCT GATGAGTGAC TGTGTCAAGG	660
	TGTTTCATCGC CACGTCCAC TTTCCTCACC CCAAATATGT GGACCTGTAC AAGGAGCAGC	720
25	TGATGAGCCA GATCCGGGAG ACGCAGCAGC TCTGCCACCA GACCCTGTTC ATCTTCGATG	780
	AAGCGGAGAA GCTGCACCCA GGGCTGCTGG AGGTCTTGG GCCACACTTA GAACGCCGGG	840
	CCCCTGANGG CCACAGGGCT GAGTCTCCAT GGACTATCTT TCTGTTTCTC AGTAATCTCA	900
30	GGGGGATAT AATCAATGAG GTGGTCTTAA AGTTGCTCAA GGCTGGATGG TCCCGGAAG	960
	AAATTACGAT GGAACACCTG GAGCCCCACC TCCAGGCGGA GATTGTGGAG ACCATAGACA	1020
35	ATGGCTTTGG CCACAGCCGT CTGTGAAGG AAAACCTGAT TGACTACTTC ATCCCCCTCC	1080
	TGCCTTTGGA GTACCGTCAC GTGAGGCTGT GTGCACGGA TGCCTTCTG AGCCAGGAGC	1140
	TCCTGTATAA AGAAGAGACA CTGGATGAAA TAGCCAGAT GATGGTGAT GTCCCCAAGG	1200
40	AGGAACAACT CTTTTCTTCC CAGGGCTGCA AGTCTATTTT CCAGAGGATT AACTACTTCC	1260
	TGTCATGAAG GCTAGAGGAA GACTTCCTGG AACTGCCITT CTTCCTACTA CAGGACCCTG	1320
45	GGACCTGTAG GAGCACCCCG TTTGGGACTG TGAGGTGTTT GAGGGTGTGG ACTGGCATCC	1380
	AGCAGCCACT AACAAACACA CAACTGGTGT GTAAAAGGCA GGCCTTACAT TAGAAGCCAA	1440
	GCCAATCCTT TTTCTTTTTT TTGGAGGTCC CACCGAGATA GATAGGAACT TGGATTGCTG	1500
50	AATTCAAAAA CAGAGCCCAT TCTTAAGATC ACTTGGTGCC TTAAAGACAC GCATTCCAAA	1560
	GTGGAATGTG GTTGAAGAAA GTGGGCCAGG TGGTGAAGA AAGCCATGTG GGAGCTCAGC	1620
55	AAATCCCAAG GGCTTATTAT GACACTCCAG ATGGTCTCCT TAGCATCTCA GCTCTTCTGC	1680
	AAGGAAGAGC TTGGGTGTTA GGCCTCAGAG GCTGTAGGGT CCTTGGGTTA CAGAGCCGGG	1740
	GAGAACGAAG TTCTGTGACC CAGGGGTGGA GAATACACTC TAGGTTTGGC GGCTGGTGGG	1800
60	CTTTCAAATT GGTACTTCCA GAGGAAAGCC AAGCTGCTTC TGTGTGAGC GAATCAGCCA	1860

5 AGAGCCTGAG GCTGAAGGGA AAAGTACACA GAGGAAGATA TTTTACAAAC CAGGTCAGTG 1920
 TAGGCCAAGA CTTATGGTCT ACAGATTTTG GCGGGGGAGG GGGGACCTTT TCAAAGACAA 1980
 TAGGGGGTCT TGACATGTTT GTTGATGTGA AAGATGATAA GATTAAAAAT TTTGATTTTC 2040
 CTAAAAA AAAA AAAA AAAA TTNC 2074

10

(2) INFORMATION FOR SEQ ID NO: 55:

15

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1483 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 55:

GAATTCGSCA CGMCGGTGGA GCGCCACGT CCCTTGCGGC GCGGGGAGAG AAATCGCTTG 60
 25 GACTTCGGGG CGGCCTCGGA CGGCCATGGC CTTTACCCTG TACTCACTGC TGCAGGCASC 120
 CCGTCTCTGC GTCAACGCCA TCGCAGTGCT GCACGAGGAG CGATTCTCTCA AGAACATTGG 180
 30 CTGGGGAACA GACCAGGGAA TTGGTGGATT TGGAGAAGAG CCGGGAATTA AATCACAGCT 240
 AATGAACCTT ATTGATCTG TAAGAACCGT GATGAGAGTG CCATTGATAA TAGTAAACTC 300
 AATTGCAATT GTGTTACTTT TATTATTTGG ATGAATATCA GTGGAGAAAA TGGAGACTCA 360
 35 GAAGAGGACA TGCCAGTAGA AGTTATTACT TTGGTCATTA TTGGAATATT TATATCTTAG 420
 CTGGCTGACC TTGCACTTGT CAAAAATGTA AAGCTGAAAA TAAAACCAGG GTTCTATTT 480
 ATCTGTTTTT TTTTTTAATG TTGCACTTGT AGTTTCATTA CAAAAGATCA GATCATGAAA 540
 40 GGCAGTAACT CTCAGGACT GGAATATCTG ATTGCTCAGT GTTAATAGTA GTTCATGCTG 600
 TGGTGAGATT GTTAAAAGGG TGCAAGACTG TTGCTTCTCT TTTTTTAGAT ATTTTTCTAT 660
 45 CTCTCACTTC TCAGGGATGA AATTCTTTTT CAAAGTTTTG AAGTTCCTTG CAACTTAGCC 720
 ATGATGTGAG TGGTTATCCC TAGATAAAAT TAAAAGGATT TTAAAAAGT AATTACTGCA 780
 CATAAAATGA TAAATAGGTA ATTGAATAA TTTTATTTTA AGCTCCTTGG TTAATTATTT 840
 50 TGTCTATTGT CTCAGCTATA AATTCAAATT TATACATACT ATTGAGTATT AATATTCTCT 900
 GATTTCAAGG AGAATTCTGT CAGTCACATG ATGATTATGT TTTTNTTTAA CATTCTTTCC 960
 55 ATGCACTTGT TATTTTATTA ATTGCTCTGA ATGATGAGAC CAGACCAGTG TCTACAGATT 1020
 TTCATTGTCA GAAAAATCTA TAAGTCTGCC CTTTTTACAA TGATGGATTT AAAAAAACA 1080
 ACAGCGTAAA TATTAGCCCA CAAGAGCAGT CCTAAACAAT CACAATTACA CTGTACTACC 1140
 60

CAAGAAGACT GTTTATTGTG AAGCATTTAC CTTTCAAAAA ATCATTACAT TTCTATTTCT 1200
 TGGTGGAGCA GCACATTGTG GAGTGTGATT CTTAATTCTT CATTGAGTTT GTCAATAGGA 1260
 5 CATTGATGCT GGATAGGTTG TCTTTTGTIT TTATGTCTCA GACCATCTTG TGAGATTGTT 1320
 TGCCTATCTC ATAATACAGT TTTATGCAGA AAGGTTGAAA CTATGTAAAT GGTMTTATG 1380
 GAAATTATCA GTTACAATAT TTTAAAGGTG TAGAATGGCA TCTTTGTTTA TAGGAGAACA 1440
 10 TTGTAAATA AAGTTAAATT TCTAAGTCAA AAAAAAAAAA AAA 1483

15

(2) INFORMATION FOR SEQ ID NO: 56:

(i) SEQUENCE CHARACTERISTICS:

20

- (A) LENGTH: 1123 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 56:

25

CAAAAATAAT AATAGTCATC ACATTTGTAT AGCACTGGGT CATTTTCC C AAGACCATTT 60
 AGTTACTTGA CCTCAGCTGT TGTCCAGCTT CCAGTCTTGG GGTAAATGGCA GCTTAATAAT 120
 30 CTGAAAATTG CCAAGAGAAA GATGTGGAAG GATGAAATGG AGGCAACATG AATTTCTGTC 180
 ACCTTGTCAT ATGTTCTCAT TTCCAKGCCT TGNGAGCAAG AGAGTTAGGT ATATCTTCTG 240
 TAACTCAGAC AATTTTCTTC CTCTTTGCAG AATGGCCCTT AGGAATCAAG GTAGCTTTTC 300
 35 TTTTGGAAAC TTCATGCTGT TTTTAGTGTG GATAGAAAGG AGGTATCTGC CATTTCTGTC 360
 ACCTATTTTA TTTTGTGTGA GCACCCATAA TAGATCAGCT GTCACAGCCA CAAATCTCTG 420
 40 AGGAGACTGG AATCATTTCC AGATAAATCA GAAAGTCAGA ATCACTTTAT GGTATATAGTC 480
 CTGGCTTCTT GAGAGCTTGT CTGGAGGTTG TAGCAGGGGA GCACAGCTAG TCATATACCC 540
 TWGACTARSG ACCGGTCTWC CTCTATTGGG GATGGTTGTC CTCTTCTACT GAGCTTGCAG 600
 45 CTTTGGGAGG GACGCACATG GAGTGGTGAG GGAGGAAGGG GACACCCGCC TAGCCAGCCA 660
 GATCAGCTGA ATCAACCTG GCAATCAATG GGGTGACAGA TGTTCAGCC AGATCGCCCT 720
 50 CACATCCAGT CCTACCTTCT TGGTAACAAA ACAATTGGTT TTGCTGGTCT AGAAACTGTA 780
 GGGCTAGACA TGTATTATAG GACTGGCTTA GGGAGAGTTA CTTTATATTA GCACTCATGT 840
 55 TTCTACTCAT TTATTCTTG TAGCTCATTA AAAGAAAAAC CATAATTGAG CATCTACTAT 900
 ATGCCATGCA TTGTGCTGAG TATCCATGAT GCTCAGGTGA ACGGACATG GTCCTGTAAA 960
 AAGTGTAAG TCTGCTGGGA AAGTTAGTGC TCAAAAGTGT AACTAAATAC TTGAGGCAAG 1020
 60 TGCTTTACTA GGAATAAAC TAAATATCAA GAGAACAAG ATAAGCAATT CCTTCACGAT 1080

GTTTTACATG GTAAATCCAT ACAATTTTAA AAAAAAAAAA AAA 1123

5

(2) INFORMATION FOR SEQ ID NO: 57:

10 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1239 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 57:

GTATTGATAC GAATTTTGAC TACATTTCTG ATGGTGTGTT TTGCTGGTTT TAACTTAAAA 60
GAAAAGATAT TTATTTCTTT TGCATGGCTT CCAAAGGCCA CAGTTCAGGC TGCAATAGGA 120
TCTGTGGCTT TGGACACAGC AAGGTSACAT GGAGAGAAAC AATTAGAAGA CTATGGAATG 180
GATGTGTTGA CAGTGGCATT TTTGTCCATC CTCATCACAG CCCCATTGG AAGTCTGCTT 240
25 ATTGGTTTAC TGGCCCCCAG GCTTCTGCAG AAAGTTGAAC ATCAAAATAA AGATGAAGAA 300
GTTCAAGGAG AGACTTCTGT GCAAGTTTAG AGGTGAAAAG AGAGAGTGCT GAACATAATG 360
TTTAGAAAGC TGCTACTTTT TTCAAGATGC ATATTGAAAT ATGTINAWGT TAAGCTTAAA 420
30 ATGTAATAGA ACCAAAAGTG TAGCTGTTTC TTAAACAGC ATTTTTAGCC CTNGCTCTTT 480
CCATGTGGGT GGTAAATGATC TATATCACCA ACCTKAATCT CTCTGCCTTT TTTTTCAAAC 540
35 ACCCCTTCAT CATCCATCTT AATTTCGATA AGGACATATC TACTTTAATG TACTACCACA 600
GTTTACAGTT AATGTGGGAA AGACCAGCTT CAGTATCCTC TTCAGCTAGG ATGCCCCTAA 660
CTTTTAACCT TCACAGTTTC CTGATTCATA TTTGCCCAGG CTCTGATGCC TTGAATTGGT 720
40 TTTGGCTCTC TTTTGTGGAT CTGTTTTTGT TGTAAACAT CATAATGCAG TCTCTCATTA 780
ATTTTACCA TCATTTACCC TGATAATCTG CCTCTCTCC ATTTCTCCTT CCCTTACTAC 840
45 CTTTCTTTGA ATTACTGTAA CTGATTGGTC CCACCAAAT TTTAAAGTAC ATGAAGTATC 900
TTCATTGGTT CATCCTCTTG CCCCCTCCAG ATGTCAAAAA ACTTTATCCT GCCCCCTAGC 960
TGACCACCCA GGTTCCTTTA TTTCAGTGGC CCATGTGAGT CTACCTTCCC CTAAGGAGTG 1020
50 CCCTAATCCA GCCCTTTTTT TGTTCCTTAT GACCCATATC TTTAGGCTCT TCCCATTTCT 1080
AGGTGGGAGA TAGGTAAGTT TCAAATCTAT GCCAGTCTTA TGAATATTAC ATTAGGGTAA 1140
55 TGTGCTATAA TGAAGAAATA AAAAATACAG TGCTTAAAG AAAATAAAAT TCTATTCTG 1200
TCTAAAAAAA AAAAAAAAAA CCNNGGGGGG GCGCCCGGT 1239

60

(2) INFORMATION FOR SEQ ID NO: 58:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 803 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 58:

GGCAGAGGTC AATCCAGGAC TACAAACACC TGTGCCAAGA CCTGAGCTTC TGCCAGGACC 60
TGTTCATCCTC CCTCCATTTC GACAGCTCCT ACCCACCGGA TGCGGGCCTG TYTGACGACG 120
15 AGGAGCCTCC CGATGCCAGC CTGCCTCCTG ACCCGCCACC CCTTACTGTG CCCCAGACGC 180
ACAATGCCCG TGACCACTGG CTGCAGGATG CCTTCCACAT CAGCCTCTGA AGGGCTGGGG 240
20 GGCAGGGGGC ATGCACCCAT GCAAAAGGCT CAGAACTCC CCCTCCGCA AGCCCTCAGA 300
CTTCGGAGCC TGCGCCTTCC CCCCTACCGC CTCACCTCAC AGGAGGGCCA GGCATGTATT 360
CCTCAGAGGC GAAACTGCCA AACTCTTTCT CCTGTCTTGG GTTGGCTGGC ACTGGGGCGG 420
25 GCATCTAGGG TACAGCCTCT GCTCATGGCA CTGGGCCTCC AGTTCCTCCA CATGTGTGCA 480
CCCCCAGCTT GGCCAACCTT CAGCCTTGGC GTGGGGCCCG AAGCATCTTC CCTTCCGCTT 540
30 GCGCTCTCTG GGATTGGGAT GAGTGCCTGG CTCCCATCTC CTCCTCACCT TTTGTTGCTA 600
TCGGCAGCTG CTGGCTCAGG GGCATCCAM CTCCGGGCTC TGGGTTCTC TGCCCTGGAA 660
GGGCTCCAGG ACCCGTCCA ATAACCAACC ACGGCCAGKA RGCCAAGGCC CCGTGTGGA 720
35 TATTTAAATT TAGGGGCCG TCTCCAGGGC GCGTAGATAA ATAAATACAC TCAGCGTCAA 780
AAAAAAAAA ARAAAAAAAAA ATT 803
40

(2) INFORMATION FOR SEQ ID NO: 59:

(i) SEQUENCE CHARACTERISTICS:

- 45 (A) LENGTH: 995 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
50 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 59:

GATTTGNGCA CGAGGNAACA GCTTTATTCT TGGTTATTCC TAATGTCCAC CTAGTCCTCT 60
55 TTWACTTTTC TTGGTAGGGT TAGGGTGGCA TGGGGAAATG GGACGGTATC ATTTTGTCTT 120
TTTAACTTTT TTTTTTTCCA CCTACAGCAG CTGTTTTTAC CCTGTGGTCA GTCAGGTACT 180
ATATTTAGTT TGCAGTTGCA CTGCTGATCG ACCCTTGATG GCCCCAGTTG GAAGTTGTTT 240
60

	GGGGGAAGG AAYTAGGAGA GGCCAGGSCC TCCATTTAAA CCATGTCTGT AATGTCTCCT	300
	TGGAAGAAA AAAAGATACT GTTCCAGTCA TGGTTTCCTG GTAGTTGACG TTTAAAATGG	360
5	GCCTCATTTA AAAATTTCAA TAATTCAGGC TAATTTTTC CCTTTATATG GTAACCCAC	420
	CAAGTTTGTC TAAATGTATG ATTTTATCA TGATTAAGTT TTTAYTTCCA CATCATGTGA	480
	CAACTGGCCT GGGATGGGAT ATAAGCTCAG AACACAAAGT CATTACCTC TTAATAAAT	540
10	AATCTATCT GTGGGGGT ATGTTATTT TGTCAAAGA GGACACAATA TGATGCAGAA	600
	TACACCATG AAGGATTTT TGGTTGGCA AGTTCTTATT TTTTAAATG GCTGTAAAC	660
15	CTAGCAGTGT TTCTGAAAT GCATACCTTA CCTGATGTC AGAGATCCGA TTTACTTCTT	720
	GATTCCCAG CAAGTGATT TGAAAACATT TAATCTAATC ATTCCCCCA CCGTCTGTC	780
	AAATCAAAG AAGTGGCATC CAGCACTAAT TTTCATGCAT TTATGAAAG ATGCCTGAGG	840
20	ACCTTAAGT ATAATCAAA ATTTGTGTTA ATGTGTGTC CTGTGAAG TTCTTTAGGA	900
	GTCGTAGAAC GAACTGATG CCCACTGATC ATCAAATGCA AGTTATGAAC ATTTAATAAA	960
25	AATTAAAC CAAAAAAAAA AAAAAAAAAA CTCGA	995

30 (2) INFORMATION FOR SEQ ID NO: 60:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 966 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 60:

40	GACAGTACGG TCCGAATTCC CGGGTCGACC CACGCGTCCG GGAGAGGACA TGCAGTGGGC	60
	ACAGAAAGTT CAATGGAACA GATGCCACTG TGGGCACCAA GACTGTAATG ACTCTGTGTG	120
	GTAGGTAGTT TTAAGGACT GCATGCTTG GAAATGATC TTCACTTGGA GAACATACTT	180
45	GCCTCTAGAT ATGTTGTCA CTCTAAGCAT CCTGAATATA ACAATAGAGA AAGATAAGTC	240
	AACCAACAGA TTAGGGATG TGTTCCTCA GCACATTTG GTCATTTGA TGCCAAGTTT	300
50	GACATACTGT TTAATTGGGC AGCACCTTG CTCCTTACC AGGTATGTAT CACTTTGTTA	360
	CTCCAGGTGC CATTCTTGGT GATGACAGAA TGTATCAC TATCGTTGTT AGCAAGAGGA	420
	AGCTTCAAT ATAGGAAGTT AACATCTTC CATGAGTATA AATGAATTTA AGACATTTGA	480
55	ATCAAACTT CAGTAGAGG AGGTTTAGA ATTCATAAAA CTGGTTAAG GAAATCTTT	540
	TTACTTTCC CAAGTTAAT CTTTTAAAT ATCTCTAGAC ATCAAATACT TTCTGTATGT	600
60	ATTAGCTGTG TCTGTCTATG ATGCAAGTAA CTCCTCCTT ATTTGGGGGA TAGTTCAGAG	660

5 AGGTAGGAGC ATTATCTCCC ATTTTCTCGG TGACTTCTTG GAGTATAGAA TTCACCATTT 720
 TATCOGTAAG TCTTCAAAGG ATTATGGTGG ACTAGAACTT ACATAGTGCA AAATAGTCTT 780
 CTATTTTTTAA TAGGAACCTA GAAAAAACTT AGAATTATAT ATAGAGTTGT TTCCTTTAGA 840
 AACCAGAGCT ATTTATTTGT ATTTAAAGCA CTGTTTATTA TTGTACTGA TTCTTATCCC 900
 10 TCTGTGTGAA TAAATGTAAG ACGGTGAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA 960
 ACTCGA 966

15

(2) INFORMATION FOR SEQ ID NO: 61:

20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 262 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 61:

TTGCAGGTAT ACATCCAGAT GCACAGAATG TCCATTTGTC CCTTATTGGT GATGCTAATT 60
 TTGATCACTT GGGTAAGATG TCCAGTTTCT CCAGTGTATC GTTATTGTTT TTCCTTTTGC 120
 30 AATTAGTGGG TAATTGTGA GGAGAACTT TGAGACCTTG TTTGACAATT CTGTTCTTCC 180
 ATCAAATCTA CCCCTCCCTA GGTTTAGCAT CCTTTGACAA TCCTGTCTCT GAATAAATTT 240
 35 TTAACATAAGA TGTTTNCCCA AN 262

40 (2) INFORMATION FOR SEQ ID NO: 62:

45 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 753 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 62:

50 GGCACAGGTT CTTTGTCCAG TCATGACAGA ACCATGCAAG ATATTGTTTA CAAATTGGTA 60
 CCAGGCCTCC AAGAAGGTGA GTGTCTGACT GTCTTGCTGA TCCCTGAGGT CCCAGCCTGG 120
 CCTCTGCAGC CCCTGCTCTC CTGGAAGTTT GGTCTCGGA TGGGAGGCC CTTTCTTTT 180
 55 GGCCGAATCA CGTCTTCTC ATCCCTGCTC TCAGCCCAAC TTCATCTCCT TGGCTGGTCT 240
 CTTCTTTTCT CTAAGATGCG TAKACATCTT TTTACCCCTT ATGTGTATTC ATTCAGCAAG 300
 60 TATGGATCGC ATGTTTACGA CATGGGAMCC CCAGGGNICA ACGCAGCTCC TGCCCTTCCC 360

5 AGGACCTGCTG CTTSTTCTG GGGCCACCT CCTGTCCAG GCCTGCCTCC CCTCATCCCA 420
 CAGCGCCAGC TTCCCCACAA CAGAGGAGCA GCACGTTGGC ATAGCGGTA GCTGGTGTTT 480
 CTAGAAAAAC TTCACCATAA AGTCAAATTT CATTTAGAAT TAAAAGAAAT ACCAAGTAGT 540
 ACAAATACCC TGAAAGTGA AATCGGTTGC TTGGGGATCG CTCAGCTGAA AGCTCCCCCA 600
 10 GCTCCCGACA CTCTACGGT GGTGGCCCT CCGCTGGCGA ACCGGCAANG AAGCCCAAGG 660
 AAGGGGGCCA GGTTCAGCGC CCAGGTGGG CTGTCCCTG GTTATTCCTG CTCCATCCAN 720
 AACCTTTCCA AAAGGCAGAA TAGAAAAACN TGA 753
 15

(2) INFORMATION FOR SEQ ID NO: 63:

20

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 739 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

25

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 63:

30 ACAATACATG CATCATATCT TTTGACTTTG AAGGATATCT CATGTCAAAG GAATCAAGTT 60
 ATGATTTATA GAGGATTCAG CTGGAATACC TTGTGGGTGC TGGCTGAGGG TGGCAAAACG 120
 CCTACCGAGA CATGAAGGTT TTAGCCACTA GTTTGTCTCT TGGGAGCCTG GGGTTGGCCT 180
 35 TCTACCTGCC TTTGGTGGTG ACTACACCTA AAACACTGGC CATCCCTGAN GAAGCTGCAA 240
 GAAGCTGTGG GGAAAGTTAT CATCAATGCC ACAACCTGTA CTGTCACTG TGGCCTTGGC 300
 TATAAGGAGG AGACCGTCTG TGAGGTGGGC CCTGATGGAG TGAGAAGGAA ATGTCAGACT 360
 40 CGGCGCTTAG AATGTCTGAC CAACTGGATC TGTGGGATGC TCCATTTTAC CATTTCTATT 420
 GGCAAGGAAT TTGAGCTTAG CTGTCTGAGT TCAGACATCT TGGAGTTTGG ACAGGAAGCT 480
 45 TTCCGGTTCA CTGKAKACT TGCTCGAGGT GTCATCTCCA CTGACGATGA GGTCTTCAAA 540
 CCCTTTCAAG CCAACTCCCA CTTTGTGAAG TTTAAATATG CTCAGGAGTA TGACTCTGGG 600
 ACATATCGCT GTGATGTGCA GCTGGTAAAA AACTTGAGAC TCGTCAAGAG GCTCTATTTT 660
 50 GGGTTGAGGG TCCTTCCTCC TAACTTGGTG AATCTGAATT TCCATCAGTC ACTTACTGAG 720
 GATCAGGACT AATAGAGAA 739
 55

(2) INFORMATION FOR SEQ ID NO: 64:

60

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 476 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 64:

GAATTCGGCA CGAGAGGACA TGGATTATGG GTACTACTCA GCAGGCCAGT TTTTACTCCA 60
10 CCTCTTTCTA GCTGACTTGA CACAAGCAAC AACCCAACAG AAAACCAATA CTCTGAGAA 120
TGGCTGCAAG TTTGTTTGTG CTGTCTTTTG AGGTAAGAAA TCAAGGCTGA GCTCTTCTTT 180
CTCCTAATTC TCAGGAAGGA GGAAGGCAGA TGTGAGAACA CTGATTGGGT CTGAGTGATC 240
15 TGGGCAGCAT CACTGTATAA AGGTCAGCAC ACAGATGCAA GCTCACTTGT CTGCTTNCMT 300
TCATGTGACT GAAGTGGTTA AGAARGTTGT NCAACTCCCC CCTGCACCCC CCTCACCACC 360
20 GCAGTAAGGG AGAGACAGGG CCAAACCTGC AGCTTCGGTA GAAGAGGCCA AGGCAGGTGT 420
CCAAGGCCAG ATCAGCAGTC AGCCAGGGCA AATGGGCTCA CTCTGGTTAC ATGACC 476

25

(2) INFORMATION FOR SEQ ID NO: 65:

(i) SEQUENCE CHARACTERISTICS:

30

- (A) LENGTH: 754 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 65:

AATTCGGCAC GAGACCAATT GTACTTTTAT TATATCAGGC TGATTCAC TG TTTCTAATGC 60
AATGAAGTAC ACACAGATTT TAAATTTTTY CTCAATCTGT CCCATTGTGT AGACAAATTA 120
40 ATTCAAAGTT CTTTTTCTTC CTCTCTTTT TCATCTAAGC CTGTGCTTAT GAGTAGAAAA 180
AGAGAAGAGG CTACCTTGAA ATGCCTCGGG CCCAAACTCA GAAGGCTCTG CACTCAACTG 240
45 AGCCTCCCTT CCTACTAAGA ATGGAATAGT GTTGCTTATA GGGGTGTTGG TCCAAGTATC 300
AGCTGTGGAT GATTAATTCC CAGGGCTGCT ATCACCTAAG GTAACCTCAG TAATCTTATG 360
TGTTTGGAAG GGAGGATGAG GATTATTTTT CAAATACATA ATTTTGTTTT ATTTTGAAAC 420
50 AATCTCACAC CTACAGAAAA GTTGCAATTA TAATACAAAG AGCTTCCCCC TCGCCTGAAC 480
TGTTTGATAG TAAGTTTGCC AACTGATAT ACCCAGGATC CCCAAATGCT TCAGTGTTAT 540
55 TTCTCCCGC CCAAGGACAT TCTCCCTGCA TAACCCACAA TACAACCCAT AAAAGTCAGG 600
AAAATTTAAC ACCCAGTTCC ATTTTGAAC CCATCCTGAA ATTCCAGGTG TTCATTCCAT 660
GTTTTGGCC AGTTGGTNC TTTGGTATGT TCCCTCCNT AGCCCAAAAA AAAAAAAAAA 720
60

AAACNCCAAG GGGGGGGGCC CCGGTCCCCA ATCC

754

5

(2) INFORMATION FOR SEQ ID NO: 66:

(i) SEQUENCE CHARACTERISTICS:

10

- (A) LENGTH: 1890 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 66:

15

GGCAGAGRAA AAACAAAATG GGTAAATGCAT TCGAGGTGAC AGGGTTAATG TTGGCATTAC 60

TTTGTATATGT TGTGTATGGG CAGAAACCCA AGGKGGGGTT TTKTTGAGCA TAAACACAAG 120

20

AAGCAATTAT TTGTGGCACT AGACTTAACC CAAAGGACAG ACCCCTACAT GTATATAGTA 180

GAGAAATCCT GTCTTTTAGC ACTATCTCAC AGGGGAAGCT GAGGAATCAC ATTATCTTTA 240

25

ATATAAATAA ATGAAATGCN AGCACTGTAT AATTTATATC CTTAAGCAAC TGGATTTCAMC 300

GTACCACTAA TGGCCTGGTC ATGTTTAA CATTACCCCA AAACAGCCTA ACTGTTCTGT 360

GACTCAGTGT CTCTGTGGAA TCCTATTTAG TAGCACCATG GTCTCTAAAT GTTTTGATTA 420

30

CACATCAGTA TTAGGAAAAC ATGTTTGAAG CATTGTCTAA GTCTGTTTGT GCTGATGTAA 480

CAGAATACCA TAGACTGGGK AGTTTATAAA GAGAGAAAT ATTGGCTTAC AGTTGTGGAG 540

35

GCTGGAAAGT CTAGTATCAG CGTACTGGGA TTTGGCAAGG GCCTTCTTGG TGCATGATAG 600

TATGGTGGAA GGTATCACAC GGCAGGCAGA AAGGCAGAGA GAGAACAAA GGGGGCGAAC 660

CCACTCCCTT GATGAGAACC TAAATACCTC TTAAGTCC TAACTCTCAA TGCTGTTTAC 720

40

AATGGCAACC AAATTTAAAC AAGAGTTTGT TAGGGAACAA AACTCAATC AAAACCATAG 780

CAAGTATGTA CCATGACTGT ATGTGTATTT ATAAAATACA TTCATATATT TCTACAGCAA 840

45

TATATATGAG GTACATTTAA GCATGTAAAA ATAGGAATTT TTAAAAATAG GACAGTTGTA 900

ATAATTTCTT TGTACATTCC ACTTTGGAGA CTGTTTATAT ATGGRGCTTG TTTTATCACC 960

AAAAGGCATT TTAATTTTGC ACACTTTAGA WTTCTTACAA TGTGTAATTG ACTGCTAGTT 1020

50

GCTGAACAAA GGACAGATAA AGTGTTCCTT GCACCTGAGC AGCCTAAAGG TGAGTGTAAAT 1080

ACAGATGCAC AAGTGACTGG TTGATAATGG AATGAGACCC CTTATAAGAA AGACATACAG 1140

55

AGCACGGCAG AGGAGCAAGA ACMACACAGA GGCAATGACA TTTGAGCTAG GCCTCTTATA 1200

TCTGTAGATG AACATTTGAT GGTAGGTAGT AGGGAAGATG GAACTAAGAA TATTTGAGCT 1260

ACTTAATATA TGCCAGGCAG CATGCTGAGT GCTTGTGTTT ATTTAATTCT CAAGACAGCC 1320

60

ATAAGCGGCA ATACAGGTAT TGGGCCTATT ATTCTAAATC CCATTTTATA AGAGAGTTAG 1380

5 GATTAGATTC AGTTCATCT TTCTACAAA CCTGGCACTG TCATTCCAGG CAAAGGGAGT 1440
 ACAATCCATT TTTCTCTTAA GAGGTGATT TTGCCAATGA GACAGAATGA ATCTCTACAG 1500
 CTGTGTAAGT TTCWACCGT CTTTGGGTGA CTGAAAAATT CAAATGTAAA GATGTGGCAA 1560
 AATTGGTCT CTAAGGATTT TAAGTACAGC CAAATGATAT GTCACAAGTT TTTCTCTAAA 1620
 10 TATCCAACCA TTTAGTCTTT CATAAGCTTT TAATTCCTACT AGCCTCACTT TCTGAGATTG 1680
 TTGATGTTTT CTGTCTCTAA CCTGAAATTT TCTTTGTTTG ATGTTAACAG GAGTATAATG 1740
 AAGGAGTAAC CATTTTTATT TTATGATAGT CTATCAATAG ACTTTTTTTA ACCTTCTTTA 1800
 15 AGCTAGGTGT GTTGTCTCTT TATTAAAGTC AGTTTGACCC AGCCTGTACA ACATTGCAAG 1860
 ACCTTAACTT TAATAAAAAA AAAAAAAAAA 1890
 20

(2) INFORMATION FOR SEQ ID NO: 67:

25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1614 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 30 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 67:

AAATAAGACN TCCTTGAGCA GCGATTGCTG GATCATTGAT CTGTTTGAGG AATGTCTGAC 60
 35 CTGGGCCTRA RAGCTGGAGA AGGTGCAGAT TCAAAGTRAG CGGCTCCTRA GGAGAGCCCC 120
 AAGSTGCTCG CCTTCTCCGT GGCTTCCGCA GCTACCGTCT GCACGGTGAG AGGGCACGGG 180
 CACACGGTTC GGGCTGGCGT GCAGTCTCCC AGCCAGCCAC GCTCTGCTCA GGCCTGGAAG 240
 40 TGAAAGCCGC CTCCTTCCCG TTATGCCCCC CATACAGGAG CCTCGGTTTT TCAGCAAAAC 300
 GCGGCCAGTC CCCTTCTCCA CTGCTGCCTC CCAGCAGAGG GCCCCAGGAT CTCCAAGGTC 360
 45 CCAGCTATGG CTTTGGACAA CGTGGCTTCG GCCCCTGGGG TTGCAGAGCT TGCATTGGGT 420
 TTACCTCGGT CTCATTCACT CATGGAGCCA AGGGTGGGGT TTCACCTGCG AACATCAGAC 480
 TGACTTGCTG GCGTCAAGAG CAGTIGACTC ACTGATGAAG GCCCTGGTGA GGAGAAAGCA 540
 50 CTCTGTCTTT CGCCTACTCT GTAATCGTTT TGTGATAATG AGCCATGAAA AAAGTAATGA 600
 ACTTGTGCTG TTAATCGTCA CTGTAATGAG AAGTCTTACG TACAACATAG CTGTGGTGGC 660
 55 TCGGTGGTTT AATGGCTGCA TTAGATAGGA TCCTCACATC CCATTGAGAA CCAAACTGA 720
 TACAGTGAAG CAATTAAGGT GAGCAATAG TTTTAACTTT TCTTTTTTTT TTTAAGTTTC 780
 ATTCTTCCTA GAATATTTTT CTAACAATTT TTATTTTCAGC TTAAAGATG GGTGATATAG 840
 60

CCAAACGGGC CATATAATCC AACATTGTTG AGATGTCTTA GGACATCTAA GGCAAAACTG 900
 GCACATTTGT TCTGCAGACT ATTGCAGGAA TGTTTTTTCC TAGCATTTCT ATATTATCTG 960
 5 TCCATTTCTGA GGAACCAGTG AATGTCCTAT AAATGCACCT CCTGTCAAAA CCATGCCTGA 1020
 GAGGTCCCGG CTGGGAGTGA CAGGGTGCTT NCTTAGATTG TATTGGTCTT TCTCTCATTC 1080
 TCCGAACCTA CTCCTTTTGA TGGGTAAGTC AACTAGGTYV ACAGTCCCTT ATTTTAAATG 1140
 10 CCTAAGTTTTT GACAGCAGGN AAGAAAACAA TTTTAAATAA ATTCTCATTA CATAGACGCA 1200
 CAAGAATATG TCACATAAAG AAAATGTGTT TAGAATACTG GTTTTCTATT TACGCATGAT 1260
 15 ATTTTCCTAA GTAAAATTGC CAAGTGGACT TGGAAGTCCA GAAAGGAAAA TAATTTAAAT 1320
 TAATGCTGGT GATCTTAACA ATATTTTGTA AAATGATGCT TCCCCCTTCT CCATGGTGTA 1380
 GTCAATTTTG TACAATTAGG TATCTGACTT TACAAGTTTG TTATCCTTTC TAATTTTAC 1440
 20 TGAAGTAAA GCACAAAGAA GACTACACAG AAAATCTGGA AACAGTTGCA GGTGTTGGGA 1500
 GGAAGATGAA ATCGAGCTGT CTTTAACTT TCGTATGTT TTTATCAGAA TTTGCTGGAC 1560
 25 TATGCTAGCA AGGACTTTGT TTACNATCA ATTGTACTAG TGTCTGCAGG GTTT 1614

30 (2) INFORMATION FOR SEQ ID NO: 68:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 596 base pairs
 (B) TYPE: nucleic acid
 35 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 68:

40 CTFTTCACCC TTAGAGACAG GGTTCACCTT TTTTGCCTTC TTAATGGAGA TATTCAGTTT 60
 TCTTTTTTTC ATTTAAACAA AGAAAAAATA TGTATCTACT CTACCTTCCC TCTGCTCTCC 120
 TCCCTCCCTA TCCTACTTGC CCATATGAGC ACGGCTCCCC ATGGCCACAT ACTCCTGCAA 180
 45 AGCTTTTATG CTGCTTCGCT TTTCTCTAAA CAGATCTGAT ATTGCTGCTC CTGTGGTTTT 240
 CTCAAAATTA ACTTTGCCGT GGTTTTAAAT AAGGAATCAA AATGCATTGT TGCATTAAGC 300
 50 TTTTCAATA AAGGAAATTT ACGGAAGGAA AATAGGCAAC ACCAGCAAAT TATATGTGGA 360
 CAGGTTCTAA ACTCTATATA TACATATATA TATATATATC TATATATCTA TATACGTAAT 420
 CATCTAGTTC TGTCTCTTGA CTGAAAGGAA TAACACTTCT AAAGATCACC ATTTCTGAGA 480
 55 AGTTCTTGGA AATCTTTATG TCTAAGTATG TGTATTAGAT CAGCAATAAT GACTATGTAA 540
 TCTCAAAAAA CAAATAAAAT ATTCTTAACA TGGAAAAAAA AAAAAAAAAT ACTCGA 596

(2) INFORMATION FOR SEQ ID NO: 69:

5

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1524 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 69:

ATCCGGAATT CCCGGGTGTG TTCGACCCGT CCGGGACTTT GCACAGCACC TTCCAGCCCA 60
 15 ACATTTCCCA GGGAAACTT CAGATGTGGG TGGATGTTTT CCCCAAGAGT TTGGGGCCAC 120
 CAGGCCCTCC TTTCAACATC ACACCCCGGA AAGCCAAGAA ATACTACCTG CGTGTGATCA 180
 20 TCTGGAACAC CAAGGACGTT ATCTTGACG AGAAAAGCAT CACAGGAGAG GAAATGAGTG 240
 ACATCTACGT CAAAGGCTGG ATTCTTGGCA ATGAAGAAAA CAAACAGAAA ACAGATGTCC 300
 ATTACAGATC TTTGGATGGT GAAGGGAATT TTAAGTGGCG ATTTGTTTTT CCGTTTGACT 360
 25 ACCTTCCAGC CGAACAAC TCATATCGTTG CGAAAAAGA GCATTTCTGG AGTATTGACC 420
 AAACGGAATT TCGAATCCCA CCCAGGCTGA TCATTTCAGAT ATGGGACAAT GACAAGTTTT 480
 CTCTGGATGA CTACTTGGGT TTCCTAGAAC TTGACTTGCG TCACACGATC ATTCTTGCAA 540
 30 AATCACCAGA GAAATGCAGG TTGGACATGA TTCCGGACCT CAAAGCCATG AACCCCTTA 600
 AAGCCAAGAC AGCCTCCCTC TTTGAGCAGA AGTCCATGAA AGGATGGTGG CCATGCTACG 660
 35 CAGAGAAAGA TGGCGCCCGC GTAATGGCTG GGAAAGTGGA GATGACATTG GAAATCCTCA 720
 ACGAGAAGGA GGCCGACGAG AGGCCAGCCG GGAAGGGGCG GGACGAACCC AACATGAACC 780
 CCAAGCTGGA CTTACCAAAT CGACCAGAAA CCTCCTTCCT CTGGTTCACC AACCCATGCA 840
 40 AGACCATGAA GTTCATCGTG TGGCGCCGCT TTAAGTGGGT CATCATCGGC TTGCTGTTC 900
 TGCTTATCCT GCTGCTCTTC GTGGCCGTGC TCCTCTACTC TTTGCCGAAC TATTTGTCAA 960
 45 TGAAGATTGT AAAGCCAAAT GTGTAACAAA GGCAAAGGCT TCATTTCAG AGTCATCCAG 1020
 CAATGAGAGA ATCCTGCCTC TGTAGACCAA CATCCAGTGT GATTTTGTGT CTGAGACCAC 1080
 ACCCCAGTAG CAGGTACGC CATGTCACCG AGCCCCATTG ATTCCAGAG GGTCTTAGTC 1140
 50 CTGGAAAGTC AGGCCAACAA GCAACGTTTG CATCATGTTA TCTCTTAAGT ATTAAAAGTT 1200
 TTATTTTCTA AAGTTTAAAT CATGTTTTTC AAAATATTTT TCAAGGTGGC TGGTTCCATT 1260
 55 TAAAAATCAT CTTTTTATAT GTGTCTTCGG TTCTAGACTT CAGCTTTTGG AAATTGCTAA 1320
 ATAGAATTCA AAAATCTCTG CATCCTGAGG TGATATACTT CATATTTGTA ATCAACTGAA 1380
 60 AGAGCTGTGC ATTATAAAAT CAGTTAGAAT AGTTAGAACA ATTCTTATTT ATGCCACAA 1440

CCATTGCTAT ATTTTGTATG GATGTCATAA AAGTCTATTT AACCTCTGTA ATGAAACTAA 1500
ATAAAAATGT TTCACCTTTA AAAN 1524

5

(2) INFORMATION FOR SEQ ID NO: 70:

10

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 819 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 70:

GGCACGAGGG AGAGGGACGG GGAGGGGGCG AGGGGCGGAG GCCGAGGGGG CAGGGGNTGG 60
20 GCGGCGGCCA GTGTTTACAG ATGAGCTTTA ACTGCCGCTT CAGGCGTGGA GACGGAGACC 120
CCGAGCCCG GCGGCGCTC AGCCCTTCAA CGACAGTATT GAGTGGTCAG GTTACAATAA 180
ACCGGAGAGA AAAGGTCCGC TTGCACTTTT TTTAGTTTTT TTATTTTTAG ACACCCCTCC 240
25 CCTCCAGGGT GATCTTTAAA AAAGCAAAAC AAAAAACAG ACTTTTCCAG CGCTCAGCGT 300
TTTTTCCTTT CGTCCGAAGC CGTTTTCTGA TTGACTTTT CTGCGCGGCC GGTCTCAGGC 360
30 CCACAGACGT TCCAGAGGAG GAGGGTGACA TTTTACTTCC CTTTTGGGG CTAACCAATT 420
ATGCTTTTGT ACATCAACCG TCGCGGCGCG GAGGGGCGAG GGGGCGGGG GCGAGGGGCG 480
TTCCAATCAA ATTTCTAATT TCTGTAAAT ATTAATCCCC KTTTTACTGC GGTTCCTGTT 540
35 GTCATTTTTA AAATTTTTTT AATTTTTTTT TTTTTTTTAC TTTTACTTTT TACCTCTTGT 600
GTATATGTAG GGAATTTATA GGGAAATATG TACTTTATGG AATAAATTTT AAGAACTAAA 660
40 ATATATTTTA TTTTAAATAA AGTAATGGAC CTTTAATCTT ACACAGCTAA ATTACTGATT 720
ATATATTTSC TGAGCTGATT TAAGGGTTAA AAAAATTGTA TCAAGAGTTT TATTTTTTGA 780
CTTCAAAGCC TTCTTAATAA AGCCTCTTTT CTACATGTG 819

45

(2) INFORMATION FOR SEQ ID NO: 71:

50

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1442 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 71:

AAATGCTTGG CATGAGTTTA CTTTAATGGC TGTTTCTGAG TTTGATCCCT CTCCGGAACC 60

60

AACCSTCTG ATGTGTCCTG TTCCAGCAGG AAGAGACAGA CCTGGAGGTT CTGTACTTGT 120
 GATTTCGGT TGTGGATCCT GAGAACAAGA AGTACTGGGA TCCTAAAGTT CTGACATTTG 180
 5 CAAAGCAGAT TAATGACCTA CCACATTCCA GATCATTGG TGAYYWIGTG TTGTGCGTGT 240
 GGGTGTGTGT GTGTGTGTGC CAAATTCAAG GTGGTCCCAG CCTTTCTAGT CTTCTCTAAC 300
 CTTTCTTCTC ARAARTCGCA CCTGTTCTGT CTTTCTAGGA TATAATTTTT TTTCTATTAG 360
 10 CCTGGGTAAC ACCCCAACCA ATAAAGTTTG CAATATCCAA GCCTCCTAAT TTCTCTACTT 420
 ATTAGCTTAT ATTAAGCTTC AGCATGAGCA AGCCTAAAAA CTCGCCATTA TCTGGAAAAG 480
 15 TTCTATTCA CAGGCTTTAA TCTCTCCTAG AGTAGTTAGC ACTCTTTTGT GGCTTTGTGT 540
 TCCTGTACTA GCTTGAATTC CACAGTCTGA CGTTAATAAT TAGCTCCTTA ACACGTCCAT 600
 CCTCTCTGA TGTCTGCTC TCTATTTTTT CTTCTTTCTT CCAAGTTGGG ATAAATTCAG 660
 20 CTTCTTATTT TCCTGCTCCA GAMCTTGGTT GTGGAGAAAG ATAGAAAAAG TTCCATACAG 720
 GGGACTCTGT GATCCTGCTA ACATCATTAT TTACCTAAGC TCTTTAGACT CCAGTGAAAG 780
 25 CTTCTGATTT AATGTCATGT CCTACTTTA TGCCACATGT CCCATACCAT TTTCTTTGTT 840
 TTATGCAATT TATTCCACT ATCTGATCCC ATTCCACCCA CATGACTTTG AGTGGAAAAC 900
 TTCATCTCTT CATTGCTGAG TAAACAAACT TCAGGATGAA CAAGCCCTGT CCACTATTTT 960
 30 CCCTTTTACT KTAAARKYCT GGAATTTTWWA TGATCTACGT TTTTTCCTC TGTTTTATT 1020
 CTTCACTCCA TATCAACTTA CTTGGGGATC TACACCTTCA TTCATYCTTT TCATTCTGTC 1080
 35 GGCACCTGGC TATGGAGTTT ACATTTCTCA TCATATTTAC TCCTCATAAT AATCCTGTGA 1140
 GGTATATACC ACTCTGAGTC TTGTATAAGA GAAAAAGAAA CTGAGATAGG GATAACTCAA 1200
 AGGGATAATT CATTTGCTGG AGCTACCAAC TAGCTACTAA CCATGCTAGA ATGGACAGAG 1260
 40 ATGACATTCA TGCCAAAGAC CATGTTGACT TGCTATCTCT ACATTTGCTC TAAGTTTGA 1320
 AAAAAAAAT CCCTTCAATT TATCCTCCAA CAGTCTCTT AGAACCTTAC CATGGATGCC 1380
 45 TTGTWTAACA CATTTACCT TTCTGGTAAA AAAAAAAAAA AAAAAAAAAA AAAAAACTC 1440
 GA 1442

50

(2) INFORMATION FOR SEQ ID NO: 72:

55 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1223 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

60 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 72:

AACCTGAGGA GGCTGTCATG ATAGGAGATG ATTGCAGGGA TGATGTTGGT GGGGCTCAAG 60
 5 ATGTCGGCAT GCTGGGCATC TTAGTAAAGA CTGGGAAATA TCGAGCATCA GATGAAGAAA 120
 AAATTAATCC ACCTCCTTAC TTAACCTGTG AGAGTTTCCC TCATGCTGTG GACCACATTC 180
 TGCAGCACCT ATTGTGAAGC AATGTGTGCA TCTGAAGCAA CTGAAATGC AGCTTCTTAT 240
 10 TGTCTGGAAT GAATCCCTTA CCAACTCAGT GCCAGCATCG GTAGACACCA GTCAGTGCTG 300
 ATCGCTTTT AACCCTCTTT TGTGTGCAT TAAATTAGAAA GAAAGGTATT GAATTGCGGC 360
 TAGCCAGTAA GCCTTGCTAA TCTCTTTTAT TTTGTAAC TGAGATGAGAC CCAAAGAAAG 420
 15 GGAAAGCTGA GATTTGTGTC CATTCCTTTT AAAATATTCA TCAGGTTAGG TGGGGCTGTG 480
 GGGGAAAAGC TACTACAGGG AAGAGTGTTC TCTGCTGTCT CTTCACTGGA AAACAGGGAG 540
 20 GGGGGATTTC AGACTGTGAA GAAAGTTGAA TGGTGGTTTT TAAATTATAA AGTAATGTAT 600
 TAAAAGGTGC ATTAGGCTGT AGTTCTAATA TTGAGTTCAA CTGTGAAATC CATCAGATGT 660
 GCCAAATGGA GAAGACAGAA AGCAACAAAG TGAATTGTTC TTTAGCCCAA GTGGTACAGT 720
 25 GAATTGCTT TAACAGATGT TGAAAATAA ATTTTCTACT GTATTCCCAG CACGGGTGAC 780
 TTCTTTTCT CTTCAATTAGC CAGAGATGAC TAAATTAAAT TTAGAACCAG ATTTTAATTT 840
 30 AAATTAATAT TTCCATTAAAT AACCTATTCA TTGCAGATAC CTATTATACT GTGTAACAGT 900
 TGTMTTGGA ATTTTATGTA AAATTAATAA TATCAGTATT TTACAGATGT TTTAATTAGA 960
 CATGTTATTA ACAGGAACAG TGCAGAACT AGAATCAAGC CTTATAATAT CTTATAGACC 1020
 35 ATGCATTTTG AAGTTAGTGT CCACTARGGT CCTATTAACT GTACATTGCA AGATTCATTA 1080
 TTTTGCTCT GACACTAWGG GAAAATTTT AGAAGCCAAT GGGACAGATT CCAGCCTTTA 1140
 40 AGCACTGGGT ACTACAGCCG TAAAAGGAAA TCCCGCCTGG TAGCCAGGGA TATNCCTCCC 1200
 CAGGTTAAAN CCCCCAAAT NAA 1223

45

(2) INFORMATION FOR SEQ ID NO: 73:

50 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1814 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 73:

CAAGCTTTGT ACTTAGATCT TTTACTTAGA TCTGCTTTTT GTCTTATTCT TTTTAGTGA 60
 60 TGTTCCTAAG GATTGTCTTC AGTCATGGCC TTGGGATTAA AGTGCTTCCG CATGGTCCAC 120

	CCTACCTTTC GCAATTATCT TGCAGCCTCT ATCAGACCCG TTTCAGAAGT TACACTGAAG	180
	ACAGTGCATG AAAGACAACA TGGCCATAGG CAATACATGG CCTATTCAGC TGTACCAGTC	240
5	CGCCATTTTG CTACCAAGAA AGCCAAAGCC AAAGGGAAAG GACAGTCCCA AACCAGAGTG	300
	AATATTAAATG CTGCCTTGGT TGAGGATATA ATCAACTTGG AAGAGGTGAA TGAAGAAATG	360
	AAGTCTGTGA TAGAAGCTCT CAAGGATAAT TTCAATAAGA CTCTCAATAT AAGGACCTCA	420
10	CCAGGATCCC TTGACAAGAT TGCTGTGGTA ACTGCTGACG GGAAGCTTGC TTAAACCAG	480
	ATTAGCCAGA TCTCCATGAA GTCGCCACAG CTGATTTTGG TGAATATGGC CAGCTTCCCA	540
15	GAGTGTACAG CTGCAGCTAT CAAGGCTATA AGAGAAAGTG GAATGAATCT GAACCCAGAA	600
	GTGGAAGGGA CGCTAATTCG GGTACCCATT CCCCAAGTAA CCAGAGAGCA CAGAGAAATG	660
	CTGGTGAAAC TGGCCAAACA GAACACCAAC AAGGCCAAAG ACTCTTTACG GAAGGTTCGC	720
20	ACCAACTCAA TGAACAAGCT GAAGAAATCC AAGGATACAG TCTCAGAGGA CACCATTAGG	780
	CTAATAGAGA AACAGATCAG CCAATGGCC GATGACACAG TGGCAGAACT GGACAGGCAT	840
25	CTGGCAGTGA AGACCAAAGA ACTCCTTGGA TGAAAGTCCA CTGGGGCCAG CAATACTCCA	900
	GAGCCCAGTT TCTGCTGGAT CCCATGGGTG GCACATTGGG ACTTCTCTCC CTCCCCATC	960
	TACACAGAAG ACTGTCACCA TGCTGACAGA AGCCTGTCTT TGTAAGGCC AGCCTTCCAG	1020
30	GGGAACACTC AGACATGTTT ATTCTCTTCC TGCTTCTGCT CTGGGCCGGT GGGTGGCTCT	1080
	CAGAAAWTAC TTGCTGCTGG CAAAAGCCT GTACTCAGGC ATTTGCTTTG ACTTGATGTT	1140
35	GCCAAGGGAC TGAGGCCATT GGCAGGCTTA GTACCACCTG CTCCTCATCT TAGGAGTCTC	1200
	CTTTTCAAAT AATTAGGCTC TGTTCCTATT TTAAACTCT GATATTGGCC TTCACCTGTG	1260
	ACTGGACACT TTAAGTAGAG CCCATTTTCA CTAAACAATA AAATCTAAAT AAATTGGAAG	1320
40	GAATAACAAC CACAAAGGAA AGAATAGAGT TGGTCTGGAT TGATGATCAC TGAGGATCTG	1380
	TATGTGAGGC ACCCATAACA GTAGTTTTCG CTGTGAGTCG TCTTCACACA TGCTGTTTTC	1440
45	TCTGCCTGGC TCTCTCTCC CCTCCTTACC TGGCCAGTCC TGTATTATCAT CAGGCCCTGT	1500
	CTTGATATC ACGTCCTCTG GGAAGTCTTC TTTTCCCCTC TAACCTAGGA CCTCATTAC	1560
	CGGCTCTCAT AGCAGAGTCT ACTGCTTTGT ACGAATTCTA AGTATTCTTG TTGCACTTAA	1620
50	TTAGCCTGTA TATCCTCAGA ACTTTGTGTA ATGCCTGGAG CATAGTAGGC AGTCATATGT	1680
	TGTATCGTGA ATAAATTGCA CATAGTAGCT ACCCAGCAAA TGCTGACTTC TTTTCTTTCT	1740
55	AGTCTTAACA CTCCCCTTCT AATNCATTTT CACTNTTGTA NTGTTCTCAA CATTACTTGG	1800
	TAGTGACAAA CTTT	1814

(2) INFORMATION FOR SEQ ID NO: 74:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 4712 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 74:

CATGGTACGC CTGCAGGTAC CGGTCCGGAA TTCCCGGGTC GACCCACGCG TCCGCCCAYG 60
CGTCCGGGCG CTCCGAGCCA GGGGCTATTG CAAAGCCAGG GTGCGCTACC GGACGGAGAG 120
15 GGGAGAGCCC TGAGCAGAGT GAGCAACATC GCAGCCAAGG CGGAGGCCGA AGAGGGGCGC 180
CAGGCACCAA TCTCCGCGTT GCCTCAGCCC CGGAGGCGCC CCAGAGCGCT TCTTGTCCCA 240
20 GCAGAGCCAC TCTGCMTCG CCTGCCTCTC AGTGTMTCCA ACTTTCGCT GGAAGAAAAA 300
CTTCCCGCGC GCCGGCAGAA CTGCAGCGCC TCCTCTTAGT GACTCCGGGA GCTTCGGCTG 360
TAGCKKGCTM TGCGCGCCCT TCCAACGAAT AATAGAAATT GTTAATTTTA ACAATCCAGA 420
25 GCAGGCCAAC GAGGCTKTGC TCTCCGACC CGAACTAAAG CTCCCTCGCT CCGTGCCTG 480
CTACGAGCGG TGTCTCCTGG GGCTCCAATG CAGCGAGCTG TGCCCGAGGG GTTCGGAAGG 540
30 CGCAAGCTGG GCAGCGACAT GGGGAACGCG GAGCGGGCTC CGGGTCTCG GAGCTTTGGG 600
CCCGTACCCA CGTGTCTGCT GCTCSCGCG GCGCTACTGS CCGTGTGGA CGCACTCGGG 660
CGCCCCCTCC AGGAGGACGA GGAGCTAGTG GTGCCGAGC TGGAGCGCGC CCCGGGACAC 720
35 GGGACCACGC GCCTCCGCTT GCACGCTTTT GACCAGCAGC TGGATCTGGA GCTGCGGCCC 780
GACAGCAGCT TTTTGGCGCC CGGCTTCACG CTCCAGAACG TGGGGCGCAA ATCCGGGTCC 840
40 GAGACGCGC TTCCGGAAAC CGACCTGGCG CACTGCTTCT ACTCCGGCAC CGTGAATGGC 900
GATCCAGCT CGGCTGCCG CCTCAGCCTC TCGAGGGCG TCGCGGGCG CTTCTACCTG 960
CTGGGGGAGG CGTATTTTCAT CCAGCCGCTG CCCGCCGCA GCGAGCGCT CKCCACCGCC 1020
45 GCCCCAGGGG AGAAGCCGCC GGCACCACTA CAGTTCCACC TCCTGCGGCG GAATCGGCAG 1080
GGCGACGTAG GCGGCACGTG CGGGTCTGT GACGACGAGC CCCGGCCGAC TGGGAAAGCG 1140
50 GAGACCGAAG ACGAGGACGA AGGGACTGAG GCGGAGGAC AAGGGCCTCA GTGGTCGCCG 1200
CAGGACCCGG CACTGCAAGG CGTAGGACAG CCCACAGGAA CTGGAAGCAT AAGAAAGAAG 1260
CGATTTGTGT CCAGTCACCG CTATGTGGAA ACCATGCTTG TGGCAGACCA GTCGATGGCA 1320
55 GAATTCCACG GCAGTGGTCT AAAGCATTAC CTCTCAGCT TGTTTTCGGT GGCAGCCAGA 1380
TTGTWCAAAC ACCCCAGSAT TCGTAATTCA GTTAGCCTGG TGGTGGTGAA GATCTTGGTC 1440
60 ATCCACGATG AACAGAAGGG GCCGGAAGTG ACCTCCAATG CTGCCCTCAC TCTGCGGAAC 1500

	TTTTGCAACT GGCAGAAGCA GCACAACCCA CCCAGTGACC GGGATGCAGA GCACTATGAC	1560
5	ACAGCAATTC TTTTCACCAG ACAGGACTTG TGTGGGTCCC AGACATGTGA TACTCTTGGG	1620
	ATGGCTGATG TTGGAAGTGT GTGTGATCCG AGCAGAAGCT GCTCCGTCAT AGAAGATGAT	1680
	GGTTTACAAG CTGCCTTCAC CACAGCCCAT GAATTAGGCC ACGTGTITAA CATGCCACAT	1740
10	GATGATGCAA AGCAGTGTGC CAGCCTTAAT GGTGTGAACC AGGATTCCCA CATGATGGCG	1800
	TCAATGCTTT CCAACCTGGA CCACAGCCAG CCTTGGTCTC CTTGCAGTGC CTACATGATT	1860
15	ACATCATTTT TGGATAATGG TCATGGGGAA TGTTTGATGG ACAAGCCTCA GAATCCATA	1920
	CAGCTCCCAG GCGATCTCCC TGGCACCTCG TACGATGCCA ACCGGCAGTG CCAGTTTACA	1980
	TTTGGGGAGG ACTCCAAACA CTGCCCTGAT GCAGCCAGCA CATGTAGCAC CTTGTGGTGT	2040
20	ACCGGCACCT CTGGTGGGGT GCTGGTGTGT CAAACCAAAC ACTTCCCGTG GCGGGATGGC	2100
	ACCAGCTGTG GAGAAGGGAA ATGGTGTATC AACGGCAAGT GTGTGMACAA AACCGACAGA	2160
25	AAGCATTTTG ATACGCCTTT TCATGGAAGC TGGGGAATGT GGGGGCCTTG GGGAGACTGT	2220
	TCGAGAACGT GCGGTGGAGG AGTCCAGTAC ACGATGAGGG AATGTGACAA CCCAGTCCCA	2280
	AAGAATGGAG GGAAGTACTG TGAAGGCAA CGAGTGCCT ACAGATCCTG TAACCTTGAG	2340
30	GA CTGTCCAG ACAATAATGG AAAAACCTTT AGAGAGGAAC AATGTGAAGC ACACAACGAG	2400
	TTTTCAAAG CTTCCTTTGG GAGTGGGCCT GCGGTGGAAT GGATTCCCA GTACGCTGGC	2460
35	GTCTACCAA AGGACAGTG CAAGCTCATC TGCCAAGCCA AAGGCATTGG CTACTTCTTC	2520
	GTTTTCAGC CCAAGTTGT AGATGGTACT CCATGTAGCC CAGATTCCAC CTCTGTCTGT	2580
	GTGCAAGGAC AGTGTGTAAA AGCTGGTTGT GATCGCATCA TAGACTCCAA AAAGAAGTTT	2640
40	GATAAATGTG GTGTTTGGCG GGGAAATGGA TCTACTTGTA AAAAAATATC AGGATCAGTT	2700
	ACTAGTGCAA AACCTGGATA TCATGATATC ATCACAATTC CAACTGGAGC CACCAACATC	2760
45	GAAGTGAAAC AGCGGAACCA GAGGGGATCC AGGAACAATG GCAGCTTTCT TGCCATCAA	2820
	GCTGCTGATG GCACATATAT TCTTAATGGT GACTACACTT TGTCCACCTT AGAGCAAGAC	2880
	ATTATGTACA AAGGTGTGT CTTGAGGTAC AGCGGCTCCT CTGCGCATTT GGAAAGAATT	2940
50	CGCAGCTTTA GCCCTCTCAA AGAGCCCTTG ACCATCCAGG TTCTTACTGT GGGCAATGCC	3000
	CTTCGACCTA AAATTAAATA CACCTACTTC GTAAAGAAGA AGAAGGAATC TTTCAATGCT	3060
55	ATCCCCACTT TTTTCAGCATG GGTCAITGAA GAGTGGGGCG AATGTTCTAA GTCATGTGAA	3120
	TTGGGTTGGC AGAGAAGACT GGTAGAATGC CGAGACATTA ATGGACAGCC TGCTTCCGAG	3180
	TGTGCAAAGG AAGTGAAGCC AGCCAGCACC AGACCTTGTG CAGACCATCC CTGCCCCCAG	3240
60	TGGCAGCTGG GGGAGTGGTC ATCATGTTCT AAGACCTGTG GGAAGGGTTA CAAAAAAGA	3300

AGCTTGAAGT GTCTGTCCCA TGATGGAGGG GTGTTATCTC ATGAGAGCTG TGATCCTTTA 3360
 5 AAGAAACCTA AACATTTTCAT AGACTTTTGC ACAATGGCAG AATGCAGTTA AGTGGTTTAA 3420
 GTGGTGTTAG CTTTGAGGGC AAGGCAAAGT GAGGAAGGGC TGGTGCAGGG AAAGCAAGAA 3480
 GGCTGGAGGG ATCCAGCGTA TCTTGCCAGT AACCAGTGAG GTGTATCAGT AAGGTGGGAT 3540
 10 TATGGGGGTA GATAGAAAAG GAGTTGAATC ATCAGAGTAA ACTGCCAGTT GCAAATTTGA 3600
 TAGGATAGTT AGTGAGGATT ATTAACCTCT GAGCAGTGAT ATAGCATAAT AAAGCCCCGG 3660
 GCATTATTAT TATTATTTCT TTTGTTACAT CTATTACAAG TTTAGAAAAA ACAAAGCAAT 3720
 15 TGTCAAAAAA AGTTAGAACT ATTACAACCC CTGTTTCCTG GTAATTATCA AATACTTAGT 3780
 ATCATGGGGG TTGGGAAATG AAAAGTAGGA GAAAAGTGAG ATTTTACTAA GACCTGTTTT 3840
 20 ACTTTACCTC ACTAACATG GGGGGAGAAA GGAGTACAAA TAGGATCTTT GACCAGCACT 3900
 GTTTATGGCT GCTATGGTTT CAGAGAATGT TTATACATTA TTTCTACCGA GAATTAAAC 3960
 TTCAGATTGT TCAACATGAG AGAAAGGCTC AGCAACGTGA AATAACGCAA ATGGCTTCCT 4020
 25 CTTTCCTTTT TTGGACCATC TCAGTCTTTA TTTGTGTAAT TCATTTTGAG GAAAAACAA 4080
 CTCCATGTAT TTATTCAAGT GCATTAAAGT CTACAATGGA AAAAAAGCAG TGAAGCATT 4140
 30 GATGCTGGTA AAAGCTAGAG GAGACACAAT GAGCTTAGTA CCTCCAACCT CTTTCTTTT 4200
 CTACCATGTA ACCCTGCTTT GGGAAATATG ATGTAAAGAA GTAACCTGTG TCTCATGAAA 4260
 ATCAGTACAA TCACACAAGG AGGATGAAAC GCCGGAACAA AAATGAGGTG TGTAGAACAG 4320
 35 GGTCCACAG GTTTGGGGAC ATTGAGATCA CTTGTCTTGT GGTGGGGAGG CTGCTGAGGG 4380
 GTAGCAGGTC CATCTCCAGC AGCTGGTCCA ACAGTCGTAT CCTGGTGAAT GTCTGTTTCA 4440
 40 CTCTTCTGTG AGAATATGAT TTTTCCATA TGTATATAGT AAAATATGTT ACTATAAAT 4500
 ACATGTACTT TATAAGTATT GGTTTGGGTG TTCCTTCCAA GAAGGACTAT AGTTAGTAAT 4560
 45 AAATGCCTAT AATAACATAT TTATTTTAT AATTATTAT CTAATGAAAA AAACTTTAA 4620
 ATTATATCGC TTTTGTGGAA GTGCATATAA AATAGAGTAT TTATACAATA TATGTTACTA 4680
 GAAATAAAAG AACACTTTTG GAAAAAAA AA 4712

50

(2) INFORMATION FOR SEQ ID NO: 75:

55

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1885 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 75:

	ATGCCARGAA GACTGATGGA GCAGGCTTGC AATATTAAAG TNCCAACCAA GAAGCTGAAG	60
5	AAATWTGAGA AAGAATATCC AGACAATGCG AGAGAGTCAG CTGCAACAGG AAGACCCAAT	120
	GGATAGATAC AAGTTTGTAT ATTGTAGGT AACTCCAGCT GTTGCAITTA TACTGGGAAT	180
10	CTTCATAAGA AGCTGAGAGA AAGAGAGGGG AAAAGAAAG TGGCTTTCTA CTTTCAAAAA	240
	TGAAACAAAA AGGAAAAATG GCAAAGTACT GTTTTAGCTG TGCATGTCAT ATCCACAAAG	300
	ACTTTTAGCA GGTGAAGTGT TCCAAGACTG ACACAAGGAT GTTTCAAACT TGCCTCTGTC	360
15	TGTAGAAAAT GTTAAAAATA CCAACTCACT TGAAGGAAA AATAAAAAATC ACAAAGGTAT	420
	ATTGAGCACA GTAGTGGTGT TTGTTGCAAC ATTTATTTCC ACAAATGAAT TTATGAACAA	480
20	CAGTGATATT TGAAGTAAAG TATGAAGTTT CAGAATCAA ATAATTTCAT TTTAATACGT	540
	TCNGTTAATT GTGAATCTCT TCMATGGTAA TTAGCAACAC TGTTCACAGG ATGCAAAGTT	600
	GGGAAACACT TATTTCCAAC TTATTTTTF TCCAAGTAAA TATTATCTCT CTTCAACATG	660
25	CTTTAACTTT TCAGACTCAC ACAGATACGT WACAGCTCCC TTCTCCCTCC ATATCAATAC	720
	ACTAAGATAA AAGAATACTG TATTTTCAGC ACTGAGCAGC AGTGCCAAAA TCTCTGCCA	780
30	AGAAATGGAC TGTGTGGCAT TATTAATTAA ATCACCACA TTGGGATGAC TTCCACTTTT	840
	GTAAGTAGAG TTATCTTTAT GTGGTCAGAG CTGGACATAG GCAGCATAGT CACACAGAAC	900
	ATCTTATCTC TGTGCKGAA TKGAATAGCA TGGGATGTGT GCAGAGGAAC ATGGKGGGAG	960
35	TATGTAGGTT TGTAGTCAG ACAGACCKGA ACTCAAATCT TGYTCATTTT TTAGAGCACA	1020
	GGATTTGGAY TCCAAATTGA GGGTTTAAAT CCCCATGCCA CCATTCAGCA TCTTCGACTA	1080
40	GTTATTGAAC CTYTTCTCA TSKATAAAG ATATAGTGTT TCTGATTCCT TGATGGATTG	1140
	TTACAAGGAT GAGGGATGCT GTATGTTAAG GACTCAGCTC ATAGTTGIGT TCAATAAATG	1200
	GCTGTTATTT TATGAAGCCT ACTACTACAG ATTATGCAAT TATTACTAGA ATAATGCCAC	1260
45	CTTATGTGGG TCTTCCCTC TAGTCCCTTA TTGATTGTTT TTATTTCTCT CAAGTATTGC	1320
	CAACCAATAA TCTCCCTTG CTTATAGAAG TGGTTCAAGA TCTGATTATA AAATCCACA	1380
50	TACTTCTATA GCAGATAACT ATTAACAGAT AATGTTTGRA CTAATTCAC CACCAACATT	1440
	CCCCCTCAAT AAAACCAGCT TTAAATGTAA ATCACATAGC ATACTGCTTT AGAAAGGCTT	1500
	GAAGGTAGTA ATTATAAAT ATTATTAAGC ATCCAAAATG AAGGTCTCCT TTTGCTAATA	1560
55	TCATTCAGAT TTCTTTATTA CTACAATTAT TATGAATAAA TTCTGTGAAG AGTGCTTTAA	1620
	AATAAGAGAG AAATGGRAGA CCAAAGTGT ACATTTAAAA TCAGGCTGGA ATTGAAGTTG	1680
60	TTATGTGTC TTAAATCCTT TTTTGTGCCA AAGCAGGTAT GTATACATTA ATAGTAAGAT	1740

232

GTACATTATT TTTAAAGTAC TTATMACATG TAAGATTATC AATATGTATA GTTTTATTG 1800
AGAGATCAAA GTAGGATTAA ACTTCTTGTT TTGAAAGCAG GCATTACTTT TTAAAAAAA 1860
5 AAAAAAAAAA AAAAAAAAAA AAAAA 1885

10 (2) INFORMATION FOR SEQ ID NO: 76:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 890 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 76:

20 TTCAAACCTAG CAAAAAATGT ATGAAACTAT GAAGCTCGAT GCGTGTRATC ATCAGCAGAG 60
GCCGACGCTG CAGGCAGGGC CAAAGCTTCT GACCTGGCC CCCAGGGAGG AACCCAGAGG 120
CCAGTCAGGG AGGGGCAGCG AGCTCACGGC CAGGCAGCGC CACAGCACTG GCGACCTCA 180
25 GGGAGAACAG GCACTACCCA GGGCTGGATG CGTAACGGGC CCCCCGGCCA CCCCCACCG 240
CCCATCAGAG CCGCAGCTCC TGAGAACGCA TCCGATGCN AGGCCAAAGT CAGCCATGGC 300
30 ACAACATTT GTGCATCAAG GTCCTGTGTC TCTGCAACAA CTCACCACAA ACAGAAGGGT 360
GGAAACCTCC ATGTATCGG ACGGCCACGG SCAGAATCCA ACGCCATCTC CCTGGGCTGA 420
TGTCGTGCA AGCAGGGCTG ATGCCGTAGC TTTTCCGGCT TCTGGAARCT GCCACAGCCC 480
35 CTGGCTCATG GSACCATCCT CACATCCTCT GAATCCACAT TCTCCTCTGA ATCTCCCGCC 540
TCCCTCTTTC CACTGTAAGG ACCCTGTGAT GACACTGCAC CCTCAGACCC TGGAACCCA 600
40 GGGTCATCTT TCCACCTCAG GCGCTCTGAC TTAAGCCTGC CTGGAGGGTC CCTGTGGTCA 660
CATTCATGGG TTCCAGGCTT CAGACACGGC CACTTTGTGG GATCATTACT CTGCCTACCA 720
CACCATGTGG CCCTGTGTGT GTTTTCAGGG GGCATTTGCG CYTATATGCA AATAATACAT 780
45 ATATGAATAA ACGTGTGAAT GGTGGTCACG TAGGAGARGG CATCTGTATG GGGCCACACC 840
TGTAATAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA 890

50

(2) INFORMATION FOR SEQ ID NO: 77:

55 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1657 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
60 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 77:

	AGAACGGCCT TCCCCACATC TTCCAGCACC TGC GCGCCTG AATCCGTCCC ACCCAGGCCC	60
5	AGACGCAGGC TTCTTCTCGG GTCTTGGTCC TGCATCCTCT CTCTCCAGA GCCTCCGTTA	120
	GGGGTGGGAA AGGACTTTGC CATAGGTCGC TGAGGCCACC ATCTGCTCTC TTAGTGCCA	180
10	AGGCGGTAAA AAGATAGTCY TCCCATTAGC TAGAGAGCAA ACCCCAGAAA GCCTATTGGC	240
	TGCGCCGTCC GCGGGCCTTG GTCCGNTTTG AAGGCGGGCT GCGGCTGCGA GAGGAGGGCG	300
	GGCGGGAGGC TAGCTGTTGT CGTGGTTGCT CGGAGGCACG TGTGCAGTCC CGGAAGCGGC	360
15	GAGGGGAAAC TGCTCCGCGC GCGCCGCGGG AGGAGGAACC GCCCGTCTT TTAGGGTCCG	420
	GGCCCGGCGG GGCATGGATT CAATGCCTGA GCCCGCGTCC CGCTGTCTTC TGCTTCTTCC	480
20	CTTGCTGCTG CTGCTGCTGC TGCTGCTGCC GGCCCCGAG CTGGGCCCGA GCCAGGCCGG	540
	AGCTGAGGAG AACGACTGGG TTCGCCTGCC CAGCAAATGC GAAGGGACTT GCGGTTAATC	600
	GAAGTCACTG AGAACCATT T GCAAGAGGCT CCTGGATTAT AGCCTGCACA AGGAGAGGAC	660
25	CGGCAGCAAT CGATTTGCCA AGGGCATGTC AGAGACCTTT GAGACATTAC ACAACCTGGT	720
	ACACAAAGGG GTCAAGGTGG TGATGGACAT CCCCTATGAG CTGTGGAACG AGACTTCTGC	780
30	AGAGGTGGCT GACCTCAAGA AGCAGTGTGA TGTGCTGGTG GAAGAGTTTG AGGAGGTGAT	840
	CGAGGACTGG TACAGRAACC ACCAGGAGGA AGACCTGACT GAATTCTCTT GCGCCAACCA	900
	CGTGCTGAAG GGAAAGACA CCAGTTGCCT GGCAGAGCAG TGGTCCGGCA AGAAGGGAGA	960
35	CACAGCTGCC CTGGGAGGGA AGAAGTCCAA GAAGAAGAGC AKCAGGGCCA AGGCAGCAGG	1020
	CGGCAGGAGT AGCAGCAGCA AACAAAGGAA GGAGCTGGGT GGCCTTGAGG GAGACCCAG	1080
40	CCCCGAGGAG GATGAGGGCA TCCAGAAGGC ATCCCCCTCT ACACACAGCC CCCCTGATGA	1140
	GCTCTGAGCC CACCCAGCAT CCTCTGTCTT GAGACCCCTG ATTTTGAAGC TGAGGAGTCA	1200
	GGGGCATGGC TCTGGCAGGC CGGGATGGCC CCGCAGCCTT CAGCCCCCTC TTGCCTTGCC	1260
45	TGTGCCCTCT TCTGCCAAGG AAAGACACAA GCCCCAGGAA GAACTCAGAG CCGTCATGGG	1320
	TAGCCACGC CGTCCTTTCC CCTCCCCAAG TGTTCCTCTC CTGACCCAGG GTTCAGGCAG	1380
50	GCCTTGTTGGT TTCAGGACTG CAAGGACTCC AGTGTAAGT CAGGAGGGGC AGGTGTCAGA	1440
	ACTGGGCACC AGGACTGGAG CCCCTCCGG AGACCAAAC CACCATCCCT CAGTCTCC	1500
	CAACAGGGTA CTAGGACTGC AGCCCCCTGT AGCTCCTCTC TGCTTACCCC TCCTGTGGAC	1560
55	ACCTTGCACT CTGCTGGCC CTTCCAGAG CCCAAAGAGT AAAAATGTTT TGGTTCTGAW	1620
	RAAAAAAAAA AAAAAAAAAA CCCCCGGGGG GGCCCGT	1657

(2) INFORMATION FOR SEQ ID NO: 78:

(i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 2015 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 78:

	GGCCGGGCTG AGAGAAGAGC TTGCGGGGTT TGCGGTTGAT GGCCCGGACT GAAGGGCTGG	60
	AGGCGGTGTA TGCCGCTGTT CTTGCTGTCG CTCCCGACAC CTCCGTCCGC TTCTGGTCAT	120
15	GAGAGGAGAC AGAGGCCTGA AGCAAAGACA TCTGGGTCAG AGAAAAAGTA TTTAAGGGCC	180
	ATGCAAGCCA ATCGTAGCCA ACTGCACAGT CCTCCAGGAA CTGGAAGCAG TGAGGATGCC	240
20	TCAACCCCTC AGTGTGTCCA CACAAGATTG ACAGGAGAGG GTTCTTGCCC TCATTCTGGA	300
	GATGTTTCATA TCCAGATAAA CTCCATACCT AAAGAATGTG CAGAAAAATGC AAGCTCCAGA	360
	AATATAAGGT CAGGTGTCCA TAGCTGTGCC CATGGATGTG TACACAGTCG CTTACGGGGT	420
25	CACTCCACCA GTGAAGCAAG GCTGACTGAT GATACTGCCG CAGAATCTGG AGATCATGGT	480
	AGTAGCTCCT TCTCAGAATT CCGCTATCTC TTCAAGTGGC TGCAAAAAAG TCTTCCATAT	540
30	ATTTTGATTG TGAGCGTCAA ACTTGTATG CAGCATATAA CAGGAATTTT TCTTGGAATT	600
	GGGCTGCTAA CAACCTTTAT GTATGCAAAC AAAAGCATTG TAAATCAGGT TTTTCTAAGA	660
	GAAAGTCTCT CAAAGATTCA GTGTGCTTGG TTAGTGGTAT TCTTAGCAGG ATCTTCTGTT	720
35	CTTTTATATT ACACCTTTCA TTCTCAGTCA CTTTATTACA GCTTAATTTT TTTAAATCCT	780
	ACTTTGGACC ATTTGAGCTT CTGGAAGTA TTTKGGATTG TTGGAATNAC AGACTTCATT	840
40	CTGAAATTCT TTTTCATGGG CTTAAATGC CTTATTTTAT TGGTGCCCTC TTTCATCATG	900
	CCTTTTAAAT CTAAGGGTTA CTGGTATATG CTTTGTAGAAG AATTGTGTCA ATACTACCGA	960
	ACTTTTGITC CCATACCACT TTGGTTTCGC TACCTTATAA GCTATGGGGA RPTTGGTMAC	1020
45	GTAAC TAGAT GGARTCTTGG GATACTGCTG GCTTTACTCT ACCTCATATT AAAACTTTTG	1080
	GAATTTTITG GGCATCTGAG AACTTTCAGA CAGGTTTAC GAATATTTTT TACACMACCM	1140
50	AGTTATGGAG TGGCTGCCAG CAAGAGACAG TGTTGAGATG TGGATGATAT TTGTTCAATA	1200
	TGTCAAGCTG AATTTTCAGAA GCCAATCTT CTCATTTGTC AGCATATATT TTGTGAAGAG	1260
	TGCATGACCT TATGGTTTAA CAGAGAGAAA ACATGTCCAC TCTGCAGAAC TGTGATTTC	1320
55	GACCATATAA ACAAATGGAA GGATGGAGCC ACTTCATCAC ACCTTCAAAT ATATTAAGTT	1380
	GTATAAACTA TCAAGGCCAC AAAATACTAA TGTCATTTGG TCATAATGAC TACTGATAAG	1440
60	GCATCAGAAT GGATTTTCAG GGCTACCAGA AAAATGTTTC CAGATGGTTT TAGAATGTAG	1500

5 GACTTATGAT CCAATTCACC AAAAGATTAA ATGAAACCAC CCTGTGTTTT AAAATATATA 1560
TAATGTTCAA CCTAATGTAT ATGCAACATT TATTCTATTG TAAITATTG ACAGGTAAC 1620
GCAGTGTAA ATTGTAAATG TGTTCCTTT ATGTTACCAA AACAGCAATT TGAAATTAGA 1680
ACTAGTGGTT TTAGAGAACT CAGGTATTCT TTCCTGACAT TGTTCCTCAGA ATAAAGAATA 1740
10 TTTTTCATAA TATTTTAAGA TACATACTAT CTAAAAGTAG AATTTTGTTG AGCATTGACT 1800
TTTATAATTC CCATCCTAAA AATTCCTAAT ATTTTCATAA AATTTGTATT TTTAAATGAA 1860
AATTCATAAT GTTGTATTTT ATCAGTAACA TTTTCTAAGT GAAGATTAAAT TTAGTGAGGA 1920
15 TGATACATTA TAGTATTGTA TTATTCTCTG TAGTAAGATT AGTAATAAGT GAAAATAAAT 1980
GATTTAAATT CAAAAAATAA AAAAAANTNA CTCGA 2015
20

(2) INFORMATION FOR SEQ ID NO: 79:

25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1213 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
30 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 79:

AGCCTAGTTA CAGATTGCAC TGCCTCAGAC TGTTCACAC CCAGAAGACG TCAGGTGACT 60
35 TCAGTCTGCG TGCAGTTGTG CAGCAGAGGA GACTGCAGAC TTCGGTTGAG GAAACGGGTA 120
TTTCATGTCT CAGGGAGTAG GTTTGTGCAG TTACAGCTTT TCTGTTGGTA TGCATAATTA 180
ATAATTGGAG CTGCAASCA GATCGTGACA AGAGATGGAC GGTGAGAAGA AAAATTGGAA 240
40 GGACAAGGTT GTTGACCTCC TGTACTGGAG AGACATTAAG AAGACTGGAG TGGTGTGTTG 300
TGCCAGCCTA TTCCTGCTGC TTTCATTGAC AGTATTCAGC ATTGTGAGCG TAACAGCCTA 360
45 CATTCGCTTG GCCCTGCTCT CTGTGACCAT CAGCTTTAGG ATATACAAGG GTGTGATCCA 420
AGCTATCCAG AAATCAGATG AAGGCCACCC ATTCAGGGCA TATCTGGAAT CTGAAGTTGC 480
TATATCTGAG GAGTTGGTTC AGAAGTACAG TAATTCTGCT CTTGGTCATG TGAAGTGCAC 540
50 GATAAAGGAA CTCAGGCGCC TCTTCTTAGT TGATGATTTA GTTGATTCTC TGAAGTTTGC 600
AGTGTGATG TGGGTATTTA CCTATGTTGG TGCCTTGTTT AATGGTCTGA CACTACTGAT 660
55 TTTGGCTCTC ATTCACCTCT TCAGTGTTC TGTATTATAT GAACGGCATC AGGCACAGAT 720
AGATCATTAT CTAGGACTTG CAAATAAGAA TGTAAAGAT GCTATGGCTA AAATCCAAGC 780
AAAAATCCCT GGATTGAAGC GCAAAGCTGA ATGAAAACGC CAAAATAAAT TAGTAGGAGT 840
60

TCATCTTTAA AGGGGATATT CATTTGATTA TACGGGGGAG GGTGAGGGAA GAACGAACCT 900
 TGACGTTGCA GTGCAGTTTC ACAGATCGTT GTTAGATCTT TATTTTITAGC CATGCACTGT 960
 5 TGTGAGGAAA AATTACCTGT CTGACTGCC ATGTGTTTCAT CATCTTAAGT ATTGTAAGCT 1020
 GCTATGTATG GATTTAAACC GTAATCATAT CTTTTTCCTA TCTGAGGCAC TGGTGAATA 1080
 AAAAACTGT ATATTTTACT TTGTTGAGA TAGTCTTGCC GCATCTTGGC AAGTTGAGA 1140
 10 GATGGTGGAG CTAGAAAAA AAAAAAAAA ANCTYGAGAC TAGCGGCACG AGGGGGGGCC 1200
 CGTACCCAAN ACG 1213

15

(2) INFORMATION FOR SEQ ID NO: 80:

20

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1391 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 80:

GCAGAGGCCG ACTGCTGAAG GTGGTTGCG TCGACATGCC GGTACCCCTG AGTCTCTTGC 60
 30 TGGGCGGGCG CGTTTGCGCG CCGTCACTCG CTGTGGGTTT GCGACCCGGG GGGTGGCGGG 120
 CCCAGGCCCT ATTGGCCGGG AGCCGGACCC CGATTCCGAC TGGGAGCCGG AGGAACGGGA 180
 35 GCTGCAGGAG GTGGAGAGCA CCCTGAAACG ACAGAAACAA GCAATCCGAT TCCAGAAAAT 240
 TCGGAGGCAA ATGGAGGCGC CTGGTGCCCC GCCCAGGACC CTGACGTGGG AAGCCATGGA 300
 GCAGATACGG TATTTACATG AGGAATTTCC AGAGTCTCGG TCAGTTCCCA GGTGGCTGA 360
 40 AGGCTTTGAT GTCAGCACTG ATGTGATCCG AAGAGTTTAA AAAAGCAAGT TTTTACCCAC 420
 ATTGGAGCAG AAGCTGAAGC AGGATCAAAA AGTCCTTAAG AAAGCTGGGC TTGCCCACTC 480
 45 GCTGCAGCAC CTCGGGGGCT CTGGAAATAC CTCAAAGCTG CTCCTTGCAG GCCACTCTGT 540
 ATCAGGCTCT TTGCTTATGC CAGGGCATGA AGCCTCATCT AAAGACCCAA ATCACAGCAC 600
 AGCTTTGAAA GTGATAGAGT CAGACACTCA CAGGACAAAT ACACCAAGGA GAAGGAAGGG 660
 50 AAGAAATAAA GAAATCCAGG ACCTGGAGGA GAGCTTTGTG CCTGTTGCTG CACCCCTAGG 720
 TCATCCAAGA GAGCTGCAGA AGTACTCCAG TGATTCTGAG AGCCCCAGAG GAACTGGCAG 780
 55 TGGTGCGTTG CCAAGTGGTC AGAAGCTGGA GGAGTTGAAG GCAGAGGAGC CAGATAACTT 840
 CAGCAGCAA GTAGTGCAGA GGGGCCGAGA GTTCTTTGAC AGCAACGGGA ACTTCTGTGA 900
 CAGAATTIGA GTCGGGGCTT GGCTTATGGA GATGCCTCGT GAAACACAGC TGGGCAAGTA 960
 60 TTAATGTATA TGAACAGCC TGGATTTCTG CATATGGATA AGCCACCTTG GAATAGGAAG 1020

5 AGGTGTTGAG CCTGGACTGT GGGAGGAAAG AGCTGCGTGG ATAGATTCAA ACTTCCTGTG 1080
 GTAGTGCTCC CAGTCTGACC TCTGTAGACC TTCAGTACTC ACTCTTCTTG CTTAGGCTCT 1140
 CTGTGTGTG AAAGCCATCC CGTGTTCAT GTGTGTGTAC AATTTTCTGT GATACTTGCA 1200
 ATTTATGTTT GAGAAGAAGT GAAAAGTTTG CCTTCTGACC TCATTTCCTT CTTGATCAGT 1260
 10 GAACACTAAC ATTTTGGGGA CAACCTAGTC AATTGGTTT CCTTACAACA AAATAAAGTA 1320
 AAATGTAGCA AAAAAAAAAA AAAAAAAAAACN CGGGGGGGGC CCGTCCCATT GCCCAAAGG 1380
 GGGCCGAATA A 1391
 15

20 (2) INFORMATION FOR SEQ ID NO: 81:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1008 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 25 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 81:

30 TGACATCGCC CTCATGAAGC TGCAGTCCC ACTCACTTC TCAGGCACAG TCAGGCCCAT 60
 CTGTCTGCCC TTCTTTGATG AGGAGCTCAC TCCAGCCACC CCACTCTGGA TCATTGGATG 120
 GGGCTTTACG AAGCAGAATG GAGGGAAGAT GTCTGACATA CTGCTGCAGG CGTCAGTCCA 180
 35 GGTCAATTGAC AGCACACGGT GMAATGCAGA CGATGCGTAC CAGGGGGAAG TCACCGAGAA 240
 GATGATGTGT GCAGGCATCC CGGAAGGGG TGTGGACACC TGCCAGGGTG ACAGTGGTGG 300
 GCCCCTGATG TACCAATCTG ACCAGTGGCA TGTGGTGGGC ATCGTTAGCT GGGGCTATGG 360
 40 CTGCGGGGGC CCGAGCACCC CAGGAGTATA CACCAAGGTC TCAGCCTATC TCAACTGGAT 420
 CTACAATGTC TGGAAGGCTG AGCTGTAATG CTGCTGCCCC TTTGCAGTGC TGGGAGCCGC 480
 45 TTCTTCTCTG CCCTGCCCCAC CTGGGGATYC CCCAAAGTCA GACACAGAGC AAGAGTCCCC 540
 TTGGGTACAM CCCTYTGCC ACAGCCTCAG CATTTCTTGG AGCAGCAAAG GGCCTCAATT 600
 CCTATAAGAG ACCCTCGCAG CCCAGAGGCG CCCAGAGGAA GTCAGCAGCC CTAGCTCGGC 660
 50 CACACTTGGT GCTCCAGCA TCCAGGGAG AGACACAGCC CACTGAACAA GGTCTCAGGG 720
 GTATGTCTAA GCCAAGAAGG AACTTTCCCA CACTACTGAA TGGAAGCAGG CTGTCTTGTA 780
 55 AAAGCCCAGA TCACTGTGGG CTGGAGAGGA GAAGGAAAGG GTCTGCGCCA GCCCTGTCCG 840
 TCTTCAACCA TCCCCAAGCC TACTAGAGCA AGAAACCACT TGTAATATAA AATGCACTGC 900
 CCTACTGTTG GTATGACTAC CGTTACCTAC TGTGTTCATT GTTATTACAG CTATGGCCAC 960
 60

TATTATTAAA GAGCTGTGTA ACATCAAAAA AAAAAAAAAA AAACTCGA

1008

5

(2) INFORMATION FOR SEQ ID NO: 82:

(i) SEQUENCE CHARACTERISTICS:

10

(A) LENGTH: 1261 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 82:

15

GTTTTCAAAC TCAATTCTAA GCCAAATAGT TTAGATAAAT ATTTACCCCTT ATATTTGGGG 60

GGAATTCAGG CTCACCATTT GCCGAGGCAA GCCCATCAAC AGTCTAGAGG CATATTCCTGT 120

20

GTCATTCTCT CCCGCTCCTT TCATAGAATA CTACTTTTTC CTTTGTCTTC CTGGCCATTTC 180

TCCATCATCT GCTGATTATT GCTAACCACA GGATGCTGGC AAAGCTTACA GTGATAGGCA 240

CATGTGTICA GTGATGTCCA ATACACTCTT ATCACAGTGG TTATGTCTTC TTAATCTTTT 300

25

CAAAATGCATT ATTCTACCCC TCAACCTAYA TCCAATCATT AGAACTATAC CTGACTGGAG 360

CCCAGAACTT GGGACCAATA CTTAATTCAA ATAGCAGGGG CTTGCTCACA AACATTAAGC 420

30

CCAAMAAGAA GCACAGCACT TTKGAAAAGT CAAATAGGSC TTTGGTAGCT CTGTACATTT 480

NGCAATTTAC ATTGTATTAT AGTTTATAGC ACTAATAACA CTTCACTCGT GAATCTACAG 540

TCTCAATATG ATAAGTCTTA GAACATGTTT TAGAAATAGT GGTACCTTGC TGCTATTATA 600

35

CTTAGTAACT TATACCCCAA TATAATAATA AGTATTAAAT ACAGATTGIG TATGCATTCT 660

TTGTGTGTAT ATGCCAATG TACTACTTAA CCTCACTGAT GAGCAATTAG AAAAAACAC 720

40

AAATGTGCAT AGTGAAAATA AGTCTTGGTC AATTCAGATG ATACGTGAAC CTGATAAATG 780

CTCTAATAGA TATGCTATTT TGTCTGTAT TGCTTGTMTT ACAGTATGGT GCATGTTGTT 840

TGCTAAGTAA AATGATAATA ATAATAAAGT ATACCCAATT TTAAGGTTAG AATTAAAATT 900

45

TTGCACATAT GCTTCTTGAT ATTCTGAAAT GTATTCTGTG GSTTMATTAT CTTATTCATA 960

CACATTKMGC TWGGCTTTTT ACCCCTAGGA AATAACTGTC CAAGTATATA TCTCGTCTTC 1020

50

TTTCTTGTA CTTTGATTAA ACTGCTTACT TCAACTTACA ACATTGTAAA GCCAGAATAC 1080

CTCATTTTAA CAGTGAAAAA AAATATTATG ACCTGATGTG TTCTCTTGT TTTGATTTGA 1140

ACTACCTAAA TAGGCTTAAC TGTAATAATA AATATACAAT TTTGGCAAAA AAAAAAAAAA 1200

55

AAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAGGCCGGC 1260

C 1261

60

(2) INFORMATION FOR SEQ ID NO: 83:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1045 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
10 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 83:

TCGAGTMTT TTTTMTTMTT TTTTAAGCAA CAGTTTATTG AGACGGAAAA AATATGATCC 60
15 AGCAAAGGCG AGGAGGGCAG CCGGGCCCCG AGCCAGCTGG TGTCATTGTC ACTGGCTCCC 120
AAACCTGACT CCTGTGGACG TGTCTGTACC CCAAACACAG CTGCCCCACC CAGCCCTGGC 180
ACAGAGCCCT TCTGAAAGAA AGAAAAAGA AGAAAGACGC GGCACCTGAC GCCAGCGGGT 240
20 AAAAGCAGGG CCCAGAGGC ATTTATTGAA AACACAGCAT CCAAAACAG ACATCTAGGC 300
CAGGCGCGAT GGTACAGTG ATGAGAGGGT CACTAGACAA TTATCCACAA TTCTACGACA 360
25 TGAGACAGAG ACTCAGCAAC AGTCACAGAC AGAAGGTCA TGTGTTCCCTT CCTGGGCAGG 420
GCTGAATGTG GCAGGTGCGG CGTGGAGGCT GCGTCCTGGC GGTTTGCTCC CAGGCAAGGG 480
GTACGGGGGG CCGGCTTGGC TGGGTGGGGA CCTCAAGTCT GAGGGTGAGG ATGGCTGAAT 540
30 CTACCTCGCT TATGTCTCAG GGACGGTCAC CCATACCTAG GATGACCCCA GCCAGACCCT 600
AGAAGGTCTG ATGGCCATCC CAAGTNCCCC CGCGAGGAGA AGAGTTCCCTT GGCAGGGGTG 660
35 ACACATTCCC GGTCAACAAG CCACAACACA GTGGTGCCCTG CACTCTCTCA GCTGTGCCA 720
CAACACTTGG TGCTGGAATT TTCTCCACGT AGTGAACTT TTAAGGGACA CATGAATAAT 780
TTAAAAAGTC ACACAAAACCT CTACGAAAGG CAGGAATCCT CACTCTGCTG AGAGCTACCT 840
40 CCTGAGATGT CGCTTCCGGA CCCCGGCAGA GGGCAGGAGC GACATCAGCT CGGCAGGAGG 900
ATCCTNGCCA GCGCGAGGGC TGGCTCTGGT TATTATAAAT AATCTAATTT AAATACGCAC 960
45 ATACACACAG ATGTCCTGCT TCTACCNAAC GCCAAGAAAA GCAGACATTA GCATCACACT 1020
GTCAACACTT CCTCGAGAAC NGAAG 1045

50

(2) INFORMATION FOR SEQ ID NO: 84:

- 55 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 2877 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

60 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 84:

	GAATTCGGCA CGAGACAAGA TGGCAGTCAA CAGCTTCCCA AAAGATAGGG ATTACAGAAG	60
	AGAGGTGATC ACAGACATGA AAAGATGCGA GACGCCGGAG ATCCTTCACC ACCAAATAAA	120
5	ATGTTGCGGA GATCTGATAG TCCTGAAAAC AAATACAGTG ACAGCACAGG TCACAGTAAG	180
	GCCAAAAATG TGCATACTCA CAGAGTTAGA GAGAGGGATG GTGGGACCAG TTA CTCTCCA	240
10	CAAGAAAATT CACACAACCA CAGTGCTCTT CATAGTTCAA ATTACATTTC TTCTAATCCA	300
	AGCAATAACC CAAGCAAAAC TTCAGATGCA CCTTATGATT CTGCAGATGA CTGGTCTGAG	360
	CATATTAGCT CTTCTGGGAA AAAGTACTAC TACAATTGTC GAACAGAAGT TTCACAATGG	420
15	GAAAAACCAA AAGAGTGGCT TGAAAGAGAA CAGAGACAAA AAGAAGCAAA CAAGATGGCA	480
	GTCAACAGCT TCCCAAAAGA TAGGGATTAC AGAAGAGAGG TGATGCAAGC AACAGCCACT	540
20	AGTGGGTTTG CCAGTGGAAT GGAAGACAAG CATTCCAGTG ATGCCAGTAG TTTGCTCCCA	600
	CAGAATATTT TGTCTCAAAC AAGCAGACAC AATGACAGAG ACTACAGACT GCCAAGAGCA	660
	GAGACTCACA GTAGTTCTAC GCCAGTACAG CACCCCATCA AACCAGTGGT TCATCCA ACT	720
25	GCTACCCCAA GCACTGTTCC TTCTAGTCCA TTTACGCTAC AGTCTGATCA CCAGCCAAAG	780
	AAATCATTTG ATGCTAATGG AGCATCTACT TTATCAAAAC TGCCTACACC CACATCTTCT	840
30	GTCCCTGCAC AGAAAACAGA AAGAAAAGAA TCTACATCAG GAGACAAACC CGTATCACAT	900
	TCTTGCACAA CTCCTTCCAC GTCTTCTGCC TCTGGACTGA ACCCCACATC TGCACCTCCA	960
	ACATCTGCTT CAGCGGTCCC TGTTTCTCCT GTTCCACAGT CGCCAATACC TCCCTTACTT	1020
35	CAGGACCCAA ATCTTCTTAG ACAATTGCTT CCTGCTTTGC AAGCCACGCT GCAGCTTAAT	1080
	AATTCTAATG TGGACATATC TAAAATAAAT GAAGTTCTTA CAGCAGCTGT GACACAAGCC	1140
40	TCACTGCAGT CTATAATTCA TAAGTTTCTT ACTGCTGGAC CATCTGCTTT CAACATAACG	1200
	TCTCTGATTT CTCAAGCTGC TCAGCTCTCT ACACAAGCCC AGCCATCTAA TCAGTCTCCG	1260
	ATGTCTTTAA CATCTGATGC GTCATCCCCA AGATCATATG TTTCTCCAAG AATAAGCACA	1320
45	CCTCAAACTA ACACAGTCCC TATCAAACCT TTGATCAGTA CTCCTCCTGT TTCATCACAG	1380
	CCAAAGGTTA GTACTCCAGT AGTTAAGCAA GGACCAGTGT CACAGTCAGC CACACAGCAG	1440
50	CCTGTAACTG CTGACAAGCM GCAAGGTCAT GAACCTGTCT CTCCTCGAAG TCTTCAGCGC	1500
	TCAAGTAGCC AGAGAAGTCC ATCACCTGGT CCCAATCATA CTTCTAATAG TAGTAATGCA	1560
	TCAAATGCAA CAGTTGTACC ACAGAACTCT TCTGCCCCGAT CCACGTGTTTCT ATTAACGCCT	1620
55	GCACTAGCAG CACACTTCAG TGAAAATCTC ATAAAACACG TTCAAGGATG GCCTGCAGAT	1680
	CATGCAGAGA AGCAGGCATC AAGATTACGC GAAGAAGCGC ATAACATGGG AACTATTAC	1740
60	ATGTCCGAAA TTTGTA CTGA ATTAAAAAAT TTAAGATCTT TAGTCCGAGT ATGTGAAATT	1800

	CAAGCAACTT TGCAGAGCA AAGGGATACT ATTTTGTAGA CAACAAATTA AGGAACCTGA	1860
5	AAAGCTAAAA AATCAGAAAT CCTTCATGGT GTGAAGATGT GAATAATTGC ACATGGTTTT	1920
	GAGAACAGGA ACTGTAAATC TGTTGCCCAA TCTTAACATT TTTGAGCTGC ATTTAAGTAG	1980
	ACTTTGGACC GTTAAGCTGG GCAAAGGAAA TGACAAGGGG ACGGGGTCTG TGAGAGTCAA	2040
10	TTCAGGGGAA AGATACAAGA TTGATTTGTA AAACCCCTGA AATGTAGATT TCTTGTAGAT	2100
	GTATCCTTCA CGTTGTAAAT ATGTTTTGTA GAGTGAAGCC ATGGGAAGCC ATGTGTAACA	2160
15	GAGCTTAGAC ATCCAAAAT AATCAATGCT GAGGTGGCTA AATACCTAGC CTTTACATG	2220
	TAAACCTGTC TGCAAAATTA GCTTTTTTAA AAAAAAAAAA AAAAAAATG GGGGGGTAA	2280
	TTTATCATTC AGAAATCTTG CATTTTCAAA AATTCAGTGC AAGCGCCAGG CGATTTGTGT	2340
20	CTAAGGATAC GATTTTGAAC CATATGGGCA GTGTACAAAA TATGAAACAA CTGTTTCCAC	2400
	ACTTGCACCT GATCAAGAGC AGTGCTTCTC CATTTGTTTT GCAGAGAAAT GTTTTTCATT	2460
25	TCCCGTGTGT TTCCATTTCC TTCTGAAAT CTGATTTTAT CCATTTTTTT AAGGCTCCTC	2520
	TTTATCTCCT TTCTTAAGGC ACTGTGCTA TGGCACTTT CTATAACCTT TTCATTCCTG	2580
	TGTACAGTAG CTTAAATTTG CAGTGATTGA GCATAACCTA CTTGTTTGTA TAAATTATG	2640
30	AAATCCATTT GCACCCCTGA AGAATGGACT TAAAAGTACT GCTGGACAGG CATGTGTGCT	2700
	CAAAGTACAT TGATTGCTCA AATATAAGGA AATGGCCCAA TGAACGTGGT TGTGGGAGGG	2760
35	GAAAGAGGAA ACAGAGCTAG TCAGATGTGA ATTGTATCTG TTGTAATAAA CATGTTAAAA	2820
	CAAAAAAAAA AAAAAAAGGG CGGCGGCTCG CGATCCTAGA ACTAGCGGAC GCGTGGG	2877

40

(2) INFORMATION FOR SEQ ID NO: 85:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1367 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

45

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 85:

50

	AATCATGAGC CTCAGAAGA GACAGATGGC CCACCAGGAG CTGTTGCTCT GGTGCTTC	60
	CTGCAGGCCT TGGAGAAGGA GGTGCCATA ATCGTTGACC AGAGAGCCTG GNAACTTGCA	120
55	CCARAAGATT GTTGAAGATG CTGTTGAGCA AGGTGTTCTG AAGACGAGA TCCCGATATT	180
	AACTTACCAA GGTGGATCAG TGGAAGCTGC TCAGGCATTC CTGTGCAAAA ATGGGGACCC	240
60	GCAGACACCT AGATTTGACC ACCTGGTGGC CATAGAGCGT GCCGGAAGAG CTGCTGATGG	300

CAATTACTAC AATGCAAGGA AGATGAACAT CAAGCACTTG GTTGACCCCA TTGACGATCT 360
TTTTCTTGCT GCGAAGAAGA TTCCTGGAAT CTCATCAACT GGAGTCGGTG ATGGAGGCAA 420
5 CGAGCTTGGG ATGGGTAAAG TCAAGGAGGC TGTGAGGAGG CACATACGGC ACGGGGATGT 480
CATCGCCTGC GACGTGGAGG CTGACTTTGC CGTCATTGCT GGTGTTTCTA ACTGGGGAGG 540
CTATGCCCTG GCCTGCGCAC TCTACATCCT GTACTCATGT GCTGTCCACA GTCAGTACCT 600
10 GAGGAAAGCA GTCGGACCCT CCAGGGCACC TGGAGATCAG GCCTGGACTC AGGCCCTCCC 660
GTCGGTCATT AAGGAAGAAA AAATGCTGGG CATCTTGGTG CAGCACAAG TCCGGAGTGG 720
15 CGTCTCGGGC ATCGTGGGCA TGGARGTGA TGGGCTGCCC TTCCACAACA MCCACGCCGA 780
GATGATCCAG AAGCTGGTGG ACGTCACCAC GGCACAGGTG TAACCGTCCA TGTTCCTGTG 840
GAGCAGAGTC CCTACCAACG GGCAGGTCTG CATCCGGGA GAATGCAGCT GCTTCTGGCG 900
20 ACAATCCTGC TAGTAAACAC TGGTCTTCGG TGAGCAACGA ACACTCGCCT GGCCTGGGAA 960
ACTGCATGCC CACTTTCTGG GAGGGGTTAG TGCAGGTGCC GTGGACAAAG GACAACATTT 1020
25 CTCTGGGGCT TTTTAACTTT TATTCCTAAG ACTCTAAAG CGTTGATTTT AACCCCTCCTT 1080
CACTCTGGCT TCTTCAGGCA ACCCAGTGG TCTCTGTGA GAATCTTCTC GACAGTTACT 1140
TATGGGGACA CTGTGAACA ATTAACTGCC AGGCAGAGCA TGAGAACAAA CATTTCCAGG 1200
30 CCATGTAGGA TAGGATACTC CAGACTCCAG TCATCCTCCC CCATCCATGG TTTCTGTTAC 1260
TCATGTTTC AGTTACTCAT AGCCAATGC AGACCGAAAA TACTAAATGA AAAATTTTAC 1320
35 AAATAAACAA CTCTTAAGTT TTAAAAAAA AAAAAAWAA ACTCGTA 1367

40 (2) INFORMATION FOR SEQ ID NO: 86:

(i) SEQUENCE CHARACTERISTICS:

45 (A) LENGTH: 1009 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 86:

50 GAATTCGGCA CGAGCTCGTG CCGAATTCTC GTGCCGAAC TAAACGTATC AAGAAATACC 60
TGGGCTTGAA GAATATTTCAC CTGAAATATA CCAAGAAACA TCCCAGCTTG AAGAATATTC 120
ACCTGAAATA TACCAAGAAA CACCGGGGCC TGAAGACCTC TCTACTGAGA CATATAAAAA 180
55 TAAGGATGTG CCTAAAGAAT GCTTTCCAGA ACCACACCAA GAAACAGGTG GGCCCCAAGG 240
CCAGGATCCT AAAGCACACC AGGAAGATGC TAAAGATGCT TATACTTTTC CTCAAGAAAT 300
60 GAAAGAAAAA CCCAAGAAG AGCCAGGAAT ACCAGCAATT CTGAATGAGA GTCATCCAGA 360

5 AAATGATGTC TATAGTTATG TTTTGTTTTA ACAATGCTCA ACCATAAAGT TGTGGTCCAA 420
 TGGAACATAC AGCTTAATAG TTTATGCGTG ATTTTCTCAA AATATTGTAA AACTTTTGAC 480
 AATGCTCATT AATATTATTT TTTCTATTTG TAGACCATAT CTGAAAGAAA TAACATTTTT 540
 TAAGGCTCTA CCACATAGAC AATATCATGC TAGAATGTGT GTGTGTGTGT GTGTGTGTGT 600
 10 GTGTGTATGT ATGTATAGGT CGGGGAGAGG ATAGTGGTGG GAACAGACAA ATAAGGAAGC 660
 GGGGAGGACT GGATAATTGG TTTTCCCCC TAAGAACATT TATTTACGTC TTAAGAGCAG 720
 ATAAGTGACT AAGACTGAAC ACATACATTT TGtGGAGTAT ATAGTTTCT TGTAAATGCT 780
 15 GTTCAATTAT TAATGTAACA GTAGCATCAA AATTTTATTG AGGCTTTAGT TGACTCTTTT 840
 GGTCAGTTTT AACAATTCTC CTAAAAGAT ATTTTGAGT GATGAATGTA GTTTACTTTT 900
 20 GTATTTGAAT TTTGATTTTC TATTTTATTT TTTTAAATAT TGTATTTGTG CACAATGTAC 960
 ATTAAATCAT TATTACATGC TTAACAAAAA AAAAAAATA AAAACTCGA 1009

25

(2) INFORMATION FOR SEQ ID NO: 87:

30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1367 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 87:

AATTCCAAAA CAAGGTAAAA GGAACCAGAA AAGAAAAAA ATGTAAATAA AGTTATAAAA 60
 ATAAAGAATT TTTTCAAGGT TAAAAAGCTG AAAAAGAAAT AATTTTATAT AAGAAAGAAT 120
 40 TTTATATGGT AAATTTAGTC CTAAAATAAA ATAACTGGTT GTTTAACAAG GAGGGATGTT 180
 CAGGACAAAC CAGAAAGTCC AAGCATGTCA TGAACATTGG TGTAAGTCAT GATAAGATTT 240
 45 TATATATATA TATACACACA CACACACACA CCCCAAAAGC TTTTATATAA TCAAGTTGTC 300
 MTATTATTAT TAAGTTTGGG TTTGCTTAGG GAAGAAAGAR CTAATTTTAA AAAAATCAAG 360
 GTTATTACAT CCATGTATCT TCCTGTGTAT GCTTTTAAAG TCCTTGTAAC ATTGAGTTAC 420
 50 AGGGCTTTAA CTCCTGTGTC TGAAAAATCA CAAACACTGA TGACAATCAA AGCCTCATCT 480
 TAAGGCCCCG TAGAAGATGC CAATCAAAAT AACTGCATT CCTGAGGCAC TAGGCAAGAA 540
 55 ATTAAAGCTA TTCAACTCCT CAAGGCCAG GACTATTGTC GGAAGAGGTG GGCGCGTAAG 600
 ATTGTAAGGG CCGATTTTGA AAGATCCAGT AAGTTTCAGT TCTCTATGAA CTAATCATTC 660
 AAGTCAAAGG CAACTGATG CAAAATCAGT ATATGGACCC CTGTGTCTGA TTAGCAAGGT 720
 60

TTTCTTGAAG CATTAACCAA CTCCTTCATA AAGGTTATAA AAGGCTTATG GRAGTTATAT 780
 TTTATAATCA AGATTAAATC TTATAGTTTG TTTACAAAAT TTGAAAATC AAATGTGATT 840
 5 GGCTTCAGGC TGTTTTTATT AGGGCTTCTT GTTTAGAAAG TTAAGTCACC TCTCTCAAAG 900
 AATGAAGGTT TTTGCTTTTT TTGAAATCCT TGAATTATCA CTGGRITAA ATAAATGACT 960
 10 TTACGATGAC CTGTAATTTT ATTTTGTAAT GTCAAGTGTT TTAAACCTTT TGTATTTGAC 1020
 AAGCTTTCCA AAATCAAATT ATAAATTATG TATTTTCTA ACCTAATTAA TCCTTTAAGA 1080
 TCTTAGTTTC CCTAAAGTCC TAAATGACA TAATTTGGCT TATTTGGTAT AAAAATTATA 1140
 15 TAGGAAGCAT TGTCAAATGT GAAATGGTGT TTGGTTTTCT TTGGGCTGTA TTTGTATAAA 1200
 TATGTTATTG GTGTATGTTT CAAAATTATG TGAAACTCCT ATAATTCTAA TATAACTTAG 1260
 TGTACATTAT CAGTAATAAT CATAATTGTT ATATTAAAT TATGTGTGC CACAGAGGTA 1320
 20 AAAAAAAGG AATTCGATAT CAAGCTTATC GATACCGTCG ACCTCGA 1367

25

(2) INFORMATION FOR SEQ ID NO: 88:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1088 base pairs
 30 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 88:
 35

GAATTCGGCA CGAGTGAAAT TTGTGCGATT TCAAAAATGG AAAATACATA ATATGCCAGG 60
 CACTTCCTGG GCAATACAGA TACCTGCACT AATGGAGTGA GCACCAGCAT CTTCCCTGAT 120
 40 GGCGTGTGCA GTGAGGTGAC TCGTCTGTAG TGTCCTCAAG GTCACGTAGA GAGCATACAG 180
 TAAATACTTG TTGACTCTTT CAACTTAAG TTAATGATAC AGTCAGGACT GATAGCCATT 240
 TTGTGTCTTT TCTTGAAAGT TTACGTGGAA GGCAGACCTT GTGTATGCTT TTCAAAGGGG 300
 45 CTCMTTAGC GCACTTGGCG CTTAAGAATT TGAGATCAGT AAGTGTGATG GTCCTAATCT 360
 TTTTAAATA GTATTGGAAG TTGAACYCM CCTGATGGGG TTGGTTTTTT TTTTTTTTTT 420
 50 TTCCAAAAA ATAATCATTC AAAATAATCG GTTAACATTT TCAATAAGAG CATTACATAC 480
 AAGGAGTTAG GGAACAAAGA GTTTTAAAT CTGGCTCTTT TTATCTCTAC TTAGGGCGTG 540
 CATCTCTCT TCTTACCCCA ACATATACTG ACTTTTAGG ACCTCCTTTA GGGAGATCTC 600
 55 AATATCCCGA ATTTTCTGT GTGAGAGGG GAAGGAATAT GTCTTTTTTT GCTTTGGTCA 660
 GAGTGGATAC ATTTTATAGT TTGTTTTTTC AAAGACGGGT CTCTGAGTC ASTTCTTTCA 720
 60 CTGCTGCCGT AAAGAACTG TATAAAGGTG ATTGAGCAGT GAAGGCATGG ATAAAAGGGG 780

AAATATTCAG CAGTTCTGAA CGTGCATGTC ATCAAATATA AAGGAGTGAG AACTTGATGT 840
ATAAGAAAAA ATGGAAGTTA AAAAAAWAA AAATCCAAGA ATGGGCTGCT TGTTCAGTA 900
5 GTGAACCTCT CGCTGGAGGT ACTAGAGCGG AGTCTGTCTC AAGGATGCTA TTGGAAGCAC 960
CCCAGCTGTG GGTGGAAGAAC TGCACCTTCT GAGCCTAGTC TTTTATAGCC TGGRGTTTTT 1020
10 GATGCTGATG CTTTACTAC TTGTCTTAG ACTWTTTTCG CATACGCTGC TCTGTTTTCT 1080
CACCTCCA 1088

15

(2) INFORMATION FOR SEQ ID NO: 89:

(i) SEQUENCE CHARACTERISTICS:
20 (A) LENGTH: 1861 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 89:

TCTCTGCCCC TCATCTTGGT AATTAGCCAG CCTCAGATAC TTCTGTGGGC CCTGAAGTGG 60
ACTCTCAAGG TCAGACCAAG GTTGTCTGATC TCAGTCCCAC TGTCTTCAGC CAGCTGAAGC 120
30 TGTGGGGCTG GGCTGGCAGC TTTATGTCA TCTTGCTTCA CCATTTTTTT TTCTCTCTCT
TTTCATTTCTA TTTTAAGTTT AGACCAAAAA AATACAGAGT CATCCCCTAC CCCCACCCCT 240
35 CTAGAGACCC TCCAGCTAAA AACAGAGCCT GAGTTCAGGG ACCCAAGTGG TGAGCGGCGT 300
CTTTTGGGGG TGAGGGAGCT TGGGTAGATG AGGCTCCTGG CTGAGCCCTC CCTGTGGTGA 360
TCCAGCCTA AGATGGCCCC TCTTCCCTCC TGGTGGGAGA CAGAGGACTG GACCCTGGGT 420
40 CTCAGGTTC AGCAAGTCAG GCTAGGGACC TGGGGGAGG AGACCCATGG ACTTCACCCA 480
TACTCAGTGA GGGGGCTCCT GCCGTCCTGA CGCCACCCCG CCCCATCAGC ACTTAAGCCA 540
45 CATGACACAA AGTCTGTACC GCACGGGAAA TGTTCAGCG CCTGGGCGGT GTGCATGGCC 600
TCCCGGGCTG TGGGGCAGCC GCATCTGTGA GGTGACYCGT GAAAGTAGGT GATTCCYTTG 660
CAGAACTTCA GGGACTGGGA GCAGAGGCC CTCACTCAAC GACGTTTGTG CGACATAGTA 720
50 TTGTATCCAC CTTAGTATTG TATCGAGCCT TTTCTGTGTT TTAATGAGAA AGCAGAACAC 780
TAGTTTCTTA TTAAAGACTT TAAGGGTTTG TGGGGCGGG CGGGATTAAC ACAACATTTG 840
55 GCTTTGTTTT CTTTTTCCTT TGATTTCAC ATCAGGTGTG TCGAGTGTG TGTGTGTGGA 900
GATGTTAAGA GCCTCACAAG GAAACTGGGT TATTGGAGGC CAAGGCGGCT TACAGTTCTC 960
TGCGTTCGTC ACTTAATTCC TGAATGTTTC AGAGAAACAG GAATCAGAAA ATAGCAGATA 1020
60

	TCATGTAGGA AAGAGAGGAT AAACAAAGAA AAAAGAAAA AAAATAAGCT CATACCCAAA	1080
	TTCACAAAGC CTATTTTTTA AACCAAAGCA CATTITGAAT GAGTATGGAA CCTCCATGGG	1140
5	CTCAGAAAAA AGATGCTAAT ATATTTATCT CATTGTTTAC ATAAGCTTTT ACAGTTTCAG	1200
	ACCTCAGCAG CTGTAAGGCC AGTCCAGGA ACCCTCCCCT GCTGCTGGAA ACCCTTCTGA	1260
	GTGCGCCCTG GAGTGGCTCA SGGGCAGAGA AGGGTAGCCC TGGGGCTGGG GGAGGGATTG	1320
10	GAAGCCTCCC TGGAGTCACC TGAGCCCTCG TCCCCATTCC CAGGGCCCCT CCAAGCCCAG	1380
	CTGGCACCAA ARAGCTTGGG CCCGTSCTGA CCAGCCCCCA AGGCCCTCTG GCCGGACCAT	1440
15	GCTGGTCTG ACCAGCTAGC CTACGCGGG ATGGCCGTCA GTTCTGGCCA CAGGACCGA	1500
	GTCTGGGCTT GGGTCCCCCT GCTGCTCTGC CCGTGACCCT TGGGGATGGG TTGATGCGAG	1560
	GGTCCCACTC AAGCCAAAAA GCCGGGACCT TTGCGCAGCT CTGTCGACTC TGGTGGGTCC	1620
20	CCACTCCTGG GGCCCCCTAA CCCACCCCA GGCAGCGAA GGGGCTGACT GGGTCTGGTC	1680
	CTTACCAACA TAGACGGTGC AAACACTCTT AACAGTGTG TTTTGTATC AATATGTTTG	1740
25	TGCAGTGATG AATGTATTTA TTCTCAGAC TTGGGGCGAG TGAGCGGGTG GCAGGCCGGC	1800
	TCCGCCACTG CAATGCTCCC GCCGGACCGA GCCCCAGCAA GGGCTCCTCC AGGATTGCAA	1860
30	A	1861

- (2) INFORMATION FOR SEQ ID NO: 90:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1259 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 90:

45	AATTGGCAC GAGCTCGTGG AGAGATTGAA GATGGCGGCT TCTCAGGCGG TGGAGGAAAT	60
	GCGGACCGCG TGGTCTGGG GGAGTTTGGG GTTCGCAATG TCCATACTAC TGACTTTCCC	120
	GGTAACTATT CCGGTTATGA TGATGCCTGG GACCAGGACC GCTTCGAGAA GAATTTCCGT	180
50	GTGGATGTAG TACACATGGA TGAAACTCA CTGGAGTTTG ACATGGTGGG AATTGACGCA	240
	GCCATTGCCA ATGCTTTTCG ACGAATTCCT CTAGCTGAGG TGCCAACTAT GGCTGTGGAG	300
	AAGGTCCCTG TGTACAATAA TACATCCATT GTTCAGGATG AGATTCTTGC TCACCGTCTG	360
55	GGGCTCATTC CCATTCATGC TGATCCCCGT CTTTTTGAGT ATCGGAACCA AGGAGATGAA	420
	GAAGGCACAG AGATAGATAC TCTACAGTTT CGTCTCCAGG TCAGATGCAC TCGGAACCCC	480
60	CATGCTGCTA AAGATTCCCTC TGACCCCAAC GAACTGTACG TGAACCACAA AGGCTGATCT	540

5 MTTTCAGAG GGCACATATCC GACCACTGCA TGATGATATC CTCATCGCTC AGCTGCGGCC 600
TGGCCAAGAA ATTGACCTGC TCATGCACTG TGTCAAGGGC ATTGGCAAAG ATCATGCCAA 660
GTTTTACCA GTGGCAACAG CCAGTTACAG GYTCTGCCA GACATCACCC TGCTTGAGCC 720
CGTGAAGGG GAGGCAGCTG AGGAGTTGAG CAGGTGYTTC TCAMCTGGTG TTATTGAGGT 780
10 GCAGGAAGTC CAAGGTAAAA AGGTGGCCAG AGTTGCCAAC CCCC GGCTGG ATACCTTCAG 840
CAGAGAAATC TTCCGGAATG AGAAGCTAAA GAAGTTGTG AGGCTTGCCC GGGTTCGAGA 900
TCATTATATC TTCTCTGTTG AGTCAACGGG GGTGTGCCA CCAGATGTGC TGGTGAGTGA 960
15 AGCCATCAA GTACTGATGG GGAAGTGCCG GCGCTTCTTG GATGAACTAG ATGCGGTCA 1020
GATGGACTGA GCTTGATGC TTCTGAGGCA AGCTGAAGCT TTGGGTCTG ACTGACCCAC 1080
20 CCTACAGGAC TGCTGAACAG AGAGCCAGT GTGACTAGGG ATCCTGAGTT TTCTGGGACA 1140
ATTCCAGCTT TAATCAATAC ATTTTGTTAA ATGTGCCATA AAATGAGACT TTTTACGCCT 1200
TTATAAGGCC TTAGATGTAA ATAACTCAC CCAAACAAAA AAAAAAAAAA AAAACTCGA 1259
25

30 (2) INFORMATION FOR SEQ ID NO: 91:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1566 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

35 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 91:

40 CTAGAAGAGC AAGCCGCCA GNANTGATGA AAAGTATTT TCCTGGAGAC CTTGGCAGTC 60
AGCGACAAGC TATTCCAACA ACTAAGAGAT CAGGACTCCA GTAGCAGTGA GTTCTGCACC 120
TTCTGGTGAC AGTGAGGGTG ATGAAGAGGA GACGACACAA GATGAAGTCT CTTCCACAC 180
45 ATCAGAGGAA GATGGAGGGG TGGTCAAAGT GGAGAAAGAG TTAGAAAATA CAGAACAGCC 240
TGTGTGGTGG AACGAAGKGT TAGAGCACGA GGTACAGGG AATTGGAATT CTGACCCCTT 300
GCTTGAACTC TGCCAGTGTC CCTCTGCCA GCTAGACTGC GGGACCGGGA GCAGTTGATT 360
50 GCTCACGTGT ACCAGCACAC TGCAGCAGTG GTGAGCGCCA AGAGCTACAT GTGTCTGTG 420
TGTGCGCGG CCCCTAGCTC CCGGGGTCA TTGGGTGCGC ACCTCTTAAT CCACTCGGAG 480
55 GACCAGCGAT CTAAGTGTG TGTGTGTGA GCCCGGTTCA CCAGCCATGC CACTTTTAAC 540
AGTGAGAAAC TTCTGAAGT ACTAAATATG GAATCCCTAC CCACAGTCCA CAATGAGGGT 600
CCCTCCAGTG CTGAGGGGAA GGATATGCC TTTAGTCCTC CAGTGTACCC TGCTGGAATT 660
60

	CTGCTTGTGT GCAACAACG TGCTGCCTAC CGTAAAMTGC TGAAGCCCA GACTCCCACT	720
	GTASGCAAGT GGGCTCTACG TCGACAGAAT GAGCCTTTGG AAGTACGGCT GCAGCGGCTG	780
5	GAACGAGAGC GCACGGCCAA GAAGAGCCGG CGGGACAATG AGACCCCGA GGAGCGGGAG	840
	GTGAGGCGCA TGAGGGACCG TGAAGCCAAG CGCTTGCAGC GCATGCAGGA GACAGACGAG	900
	CAGCGGGCAC GCCGGCTGCA GCGGGATCGG GAGGCCATGA GGCTGAAGCG GGCCAATGAA	960
10	ACCCCGGAAA AGCGGCAGGC CCGGCTCATC CGAGAGCGAG AGGCCAAGCG GCTCAAGAGG	1020
	AGGCTGGAGA AAATGGACAT GATGTTGCGA GCTCAGTTTG GCCAGGACCC TTCTGCCATG	1080
15	GCAGCCTTAG CAGCTGAAAT GAACTTCTTC CAGCTGCCTG TAAGTGGGT GGAGTTGGAC	1140
	ARCCAGCTTC TGGGCAAGAT GGCCTTTGAA GAGCAGAACA GCAGYTYTCT GCACTGAACC	1200
	ACACCTCCT GCTGCCCCC CTTCACCT ACCTACCCAC CCACCCACAC CCACAGCCAC	1260
20	GAGGACCACT GCTGCTGCCA CCCACGAGG CCTGTCTCTG CTGCCAGAGG CAGGCCTGGG	1320
	TTTATTCAG GTGGACCTGA GCAGCCCTTG CATATGGGAA CAGGATGATG GGTCAGGAG	1380
25	GGACCTGGCT CAAGGCAGCT CTGGACAAGG GAGCAGGCAG TCCAGAGAAC TGGCTCCCC	1440
	AGCCCACTGC CACAGGCTGT GCTTCTAGGA CTGTGGGCCC CTGTGTGGCC CATGAAGTTG	1500
	TGAAGTCAA TAAATTAATT TTATCTTTAA AAAAAAAAAA AAAAAAYVGG GGGGTTTTTT	1560
30	TGGGGG	1566

35

(2) INFORMATION FOR SEQ ID NO: 92:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1593 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 92:

45	GGCAGAGCC TCGGCCTCGG TGGCGGTGGT GGACACGTCG AGCCGGGTAG AAGTGGAGGG	60
	GCCGTTGAA GAGTCGTAG GGGGTGACGG GTTAAGATTC GGAGAGAGAG GTGCTAGTGG	120
50	CTGGACTTGA CCTGAAAGA ATCTTCTGCT GACTCTCAAC TTTCTCTGGA AAAAATGGAT	180
	CATTCCACC ATATGGGGAT GAGCTATATG GACTCCAACA GTACCATGCA ACCTTCTCAC	240
	CATCACCCAA CCACTTCAGC CTCACACTCC CATGGTGGAG GAGACAGCAG CATGATGATG	300
55	ATGCCTATGA CCTTCTACTT TGGCTTTAAG AATGTGGAAC TACTGTTTTC CGGTTTGGTG	360
	ATCAATACAG CTGGAGAAAT GGCTGGAGCT TTTGTGGCAG TGTTTTACT AGCAATGTTT	420
60	TATGAAGGAC TCAAGATAGC CCGAGAGAGC CTGCTGCGTA AGTCACAAGT CAGCATTCGC	480

	TACAATTCCA TGCCTGTCCC AGGACCAAAT GGAACCATCC TTATGGAGAC ACACAAAAC	540
	GTGGGGCAAC AGATGCTGAG CTTTCCTCAC CTCCTGCAA CAGTGCTGCA CATCATCCAG	600
5	GTGGTCATAA GCTACTTCCT CATGCTCATC TTCATGACCT ACAACGGGTA CCTCTGCATT	660
	GCAKKGACAG CAGGGGCCGG TACAGGATAC TTCCTCTTCA GCTGGAAGAA GGCAGTGGTA	720
10	GTGGATATCA CAGAGCATTG CCATTGACAT CAAACTCTAT GGCCTGGCCT TATCGATTGC	780
	AGTGGGAAGT TGTGAAGAC TTGAAGACGT GATTCTGCT CCAATCATCC CTTCTTGCTC	840
	CTCTTTGKGC ACGTACACAC ACACACACAC ACACACACAC ACACACCCGT GYTCAAACAG	900
15	AGGTTTAGTT TACAGTCTCT GAACTAAAGT AGTAACCTCC CAAATTGTTT TTTCTAATAA	960
	GCTGAGATTC CCATTTCTCT TAAGGAGAAG CCACCCATGA GATGCTTTT CTTCTCCAT	1020
20	CATCTTAGAG CCAAGTTATA TGTCTTGTC TAATCCATGT AGCTTTTGT TCAATGACTT	1080
	GATCATCTGC TTCTTTTGT AATTTTAAAC AGATAGTAAG TAAATTTGGT GGTTTTTC	1140
	CCTGGGTCAG TGATGGAAG GGGTTAACTT CAGCCAGGAT TGATGGCAGC TGAGGGAAAT	1200
25	TCTTGCCCAA CTAAACCCAG AACTCAAAC TAACATTAGA AAATAAGGTC CAGGGCCGGA	1260
	CACAGTGGCC CAAGCAAGTA ATCCAGCAC TTTGGGGGGC CAAGGCAGGC TGGATCACCT	1320
30	GAGGACAGGA GTTCGAGACC AGTCTGGCCA ACATGGGGAA ACCCGTCTC TACTAAAAAT	1380
	ACATAAATTA GCCGGCATG GTGGTGGCG CCGTAATCC CAGCTACTCA GAAGGCTGAG	1440
	GCAGGAGAAT CACTTGAACA TAGGAGGCG AGGTTCAGT GAGCCAAGAT GGCGCCATTG	1500
35	CACTCCAGCC TGGGTGACAA GNGTGAACT CCATCTCATA AAAAAAAAAA AAAATANTCG	1560
	AGGGGGGGCC CGGACCCAAA ACGCCGAAA GTG	1593
40		

(2) INFORMATION FOR SEQ ID NO: 93:

- 45 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 970 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

- 50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 93:

	CTCGTGCCGA ATTGGGCAG AGGTGCCAG GCTCTCAGG CAGAGGGTCC AGTGTGATCA	60
55	CTTTGCATGG CCTCTCTCCC CTCCTGAGCT TGTGCCAGG CCCCAGGGCT GACCTGGAGA	120
	GGAAAAGGC AGAGGTGAA GATGGGGTGT CTGGTTTGGG GACCATCCTG GCCCCCTTG	180
60	TCACTGTTGG CATCTCTTCT GCACAGTGGC ATTGCTGGGA GGTGCTTACT GTGCCTATTC	240

	AAGGGGCTGG CAGCCGAGC CTCACTGCAG ATCAGGGACT TGGCTTCCCG GTTGACCACA	300
	GGTCCAAGAA CCTGCAGGT CCAGCCTCCC CCCCATCCCC AGTCTTCCCC ACCCTGGCCC	360
5	GGCCCTCCAG GTGCAGAAAC ATGCAGGCCC CTCTCCAGGA CTGTGGGAGG AGTGTGTCCC	420
	TCAGACTGGC CTGTGTCTG GCTCCTCTTA CCACCTCTTC CAGAGGTTGT CACCTGCAGC	480
	TGCCCCAGGA TAAAGGCAAG GCCAGAGAGG ACTCCTGAAC TCCTGTGTGC CTGGGGTGGC	540
10	AGGGGCAAAC ATAGCCAACT GGTGGCCTGA GCGGGGCCAT GGTGARGACA CCCTTGGTGG	600
	CTTGTCCAC ATCAAGCTGG GARGTGACAC TGAGGATGCA TTAGTCTGCA GCGTATGATA	660
15	AAAACGGCAT TTCAGGCCAG GCGTGGTGGC TCATGCCTGT CACCCAGCA CCTTGGGAGG	720
	CCGAGGTGGG CAGATCACAT GAGGTCAGGA CTTTGAGACC AGCCTGGCCA ACATGGTGAA	780
	AACTCATCTG TACTAAAAA ACAAAAATTA TGTGGGTTGG TGGTGTGTGC CTGTAATCCC	840
20	AGCTACTTGG GAGGCTGAGG CAGGAGAATC ACTTGAACCT GGGAGGCGGA GGCTACAACG	900
	AGCCGAGATT GCACCACTGC ACTCCAGCCT GATCCGTCTC AAAAAAAAAA AAAAAAAAAA	960
25	AAAAACTCGA	970

30 (2) INFORMATION FOR SEQ ID NO: 94:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 934 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 94:

40	TCTCTCTCTC TCTCTCTCTC TCTGCTGTAA AGAACTCCCA AACTCAAAT GTATCAGGAA	60
	ATGTAAAGGT TAAGTCTGAC TACAAGAAGG CCAAAATTGC ACCAGCTTCC TAAGTGAAGA	120
	ATAATAGAAT AAAACATATA GAGGGCAGAA ATAAATGAG GTGTATCTGG AGAATTTTCAT	180
45	GATGAGCATT TAGATTTAGC AATGCCCAAT GTCATGCTGA CACTGTTTGT CATGACCTTG	240
	TCTTCAGCTA GTAATTTGGG GTTGTACTTT TTTAAATTTA ATTTTGAATG TTCTTGCA TG	300
50	TTTGGTACCT CTCTCCTCAC TGCTAAAGAT AAATTGTTTA TCTGTATAAC ATAACACAC	360
	CAATGTCATT TATTGTATAC GCTAGTACAC AAATGTGTTT TTTTATTAAG TAATGAARTA	420
	TTTGCTGTGA AAAATGTATT ATTTGTGCCA CCGTTTATAT CTGTGTTTAT TTTCTGTGTG	480
55	TATATGCGTG TGTATTCGAA TCTCAATTTT TCTTTTACTC TAGTTTAGAT TAAGACATAT	540
	TTAGATGAAA TTTTAAAAAT AACATTTGGA ATAGGAGGCT AAGTTTGTGT SAGTCTCATT	600
60	CCCTTGGGGG GAAATTCCTT TTGCCATTTT ATTTTCATGT ACAATAACCT AAAAAGGATC	660

TCCTACTGAC TTCCTTCTA ATTATTATG TTTTACACGA AAGAAAGGAA ATACGTTTTC 720
 AATTGAGTTG TTTGAAATCA TTCACTTTGT GTAGATTTCC CAGACTGATG TTTTCATTGTA 780
 5 AGAATATTAC ATTATAGACA GGTGCGCCAT TTCACAAGCA ACTAATCCAT AGTTTGGGAA 840
 GCCCGCTTTA AGAGACCTGA ATATCTTTGT TTTTAATAAA ATACTTAGAG TTTAAAAAAA 900
 10 AAAAAAAAAA AAAAAAAAAA AAAAAAAGG TAAA 934

15 (2) INFORMATION FOR SEQ ID NO: 95:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1392 base pairs

(B) TYPE: nucleic acid

20 (C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 95:

25 CAGCTCAGCT CTGCGCTGCT GCACGCCAAC CACACACTCA GCACCATTGA CCACCTGGTG 60
 TTGGAGACGG TGGAGAGGCT GGGCGAGGCG GTGAGGACAG AGCTGACCAC CCTGGAGGAG 120
 GTGCTCGANC CGCGCACGGA GCTGGTGGNT GCCGCCCGAG GGGCTCGACG GCAGGCGGAG 180
 30 GCTGCGGCCC AGCAGCTGCA GGGGCTGGCC TTCTGGCAGG GAGTGCSCCT GAGCCCCCTG 240
 CAGGTGGCTG AAAATGTGTC CTTGTGGAG GAGTACAGGT GGCTGGCCTA YGTCTCTCTG 300
 35 CTGCTCCTGG AGCTGCTGGT CTGCCTCTTC ACCCTCCTNG GCCTGGGGAA CAGAGCAAGT 360
 GGCTGGTGAT CGTGATGACA GTCATGAGTC TCCTGGTTCT CGTCTGAGC TGGGGCTCCA 420
 TGGGCTTGA GGCAGCCACG GCCGTGGGCC TCAGTGACTT CTGCTCCAAT CCAGACCCCT 480
 40 ATGTTCTGAA CCTGACCCAG GAGGAGACAG GGCTCAGCTC AGACATCCTG AGCTATTATC 540
 TCCTCTGCAA CGGGGCGTC TCCAACCCCT TCCAACAGAG GCTGACTCTG TCCCAGCGAG 600
 45 CTCTGGCCAA CATCCACTCC CAGCTGCTGG GCCTGGAGCG AGAAGCTGTG CCTCAGTTCC 660
 CTTCAGCGCA GAAGCCTCTG CTGTCTTTGG AGGAGACTCT GAATGTGACA GAAGGAAATT 720
 TCCACCAATT GGTGGCACTG CTACACTGCC GCAGCCTGCA CAAGGACTAT GGTGCAGCCC 780
 50 TGCGGGGCTT GTGCGAARAC GSCCTGGAAG GCCTGCTCTT CCTGCTGCTC TTCTCCCTGC 840
 TGTCTGCAGG AGCGCTGGCC ASTGCCCTMT GCAKCCTGCC CCGAGCSTGG GCCCTCTTCC 900
 55 CACCCAGGAA TCCAAGCGCT TTGTGCAGTG GCAGTCGTCT ATCTGAGCCC CTCTCCCGG 960
 CTGGAATGGA GCCTGGCTCC CCTCTCTGTT CCTTCCCTGG CTGCCGAGA GACCCCACTA 1020
 60 ACCCAGCCTG CCTGGGCTCT GACCACTAAC ACTCTTGGCC ATGGACAGCC TGCACAGGAC 1080

CGCCTCCCTG CTCTTGGCCA CTGTGCTCCC ATTCTGTGCC TTGGCCTTGG GAGTAGCTGA 1140
 GGGGGCAGAC TAGGGAGTAG GGCTGGCAGG GGAGGGGGCA GACAGCCTCG CCTCGCACCC 1200
 5 TTATCCCTG GCTGCCGGTC CCATCCTTGG AGGGACTAAG CTGGGGGTGG GACATGAGTC 1260
 CCCCTGCTGC CCCTGCCACA TCCCAGTGGG CTCTGACCCC CTGATCTCAA CTCGTGGCAC 1320
 TAACTTGAA AAGGTTGAT TTAAAATAAA AGGAAGACT ATTTTACAAA AAAAAAAAAA 1380
 10 AAAAAACTC GA 1392

15

(2) INFORMATION FOR SEQ ID NO: 96:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1963 base pairs
 20 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 96:
 25

GGTANCTGCA GTACGGTCCG ATTCCCGGGT CGACCCACGC GTCCGGAGAA ATGCAAATTA 60
 AAACAGTAAA GTGTCATTTT CACTTCCTGG ATTGGCAAAG GGTMTTATGT ATTTTACTGA 120
 30 CAGTGCTCAA CATTAGCAGT AAACAACAAA TGGTGAGTAA ATATGAGCTT CGGAACCTCA 180
 GGGAAATGAT CTCCTTATTT CAACCTGCAG ATTCCTTCCT ACAACCAGTG TAGAGCAGAG 240
 TACCAGGACG GGCCATTGAG CACCTGGTG TTGAGATCAA GTGCCCTCTA GTCAGAGTTG 300
 35 GGTGAGGGCC ACTGTGAGTG GGCTGCCCC AACATGAGTC AGCTGTCTAG GACTAGTTTA 360
 TCTCTGCTTC TCACTTTACT GGTATTATGG GGCAGCTCCT GCTGTCTTCC AATTGGTGT 420
 40 CTCCAAATC GGCACCGTCT TTAAAGTTG AGTTTCTTGT TATTCTCACC TGATATACCT 480
 TATTTATCCC ACACCCACCC CAATAACATA TCGTGCTCAG TGTATCTTT GAGACAACAC 540
 TTGAATTTTA CTCAGCCTGG AGCGCTCTTC ACATGTCTTG TCCAGATCCA GTTCGGACTC 600
 45 ATTCTTCAGC CGTGCATCAG TAAATGGGG CTAGGTTAAA CTGTGGTGAC AAACAACCTC 660
 CAAATTTTCA TGGCTCAAAA ATCTTCTTCC TCATTTATWT ACATTTTCATC ATGGGTCAGG 720
 50 TGAGAGGTAG CTCTGTGCTG TGTATCTCTA ACACAGGAAT CCAGACGGA GGAGGGACAA 780
 TCAATAAGAT CCCATTGCT ATAGAAAAGA RAAAAAGTA TCGGAATAR CACTCYGTTT 840
 CYTGAGAWT YCTCTGAAA AAGTCACATG TTATTTCTTC TCACCTCCAT TGGCAAAAAA 900
 55 AAAGTCATGT GGCCATGTGA AAATGTAAGT AGCGGGGATG GAACAGTCAG AATGCATCA 960
 TAAAATATGA ACTGAAATA TCTGGAGAAC AKCACCTATG ACTACCACGA ATGCCAACAT 1020
 60 GCATCCCTAA CAACCCAGTG CTGTACCCTT CCAAACCTTT TATGTCTTGC AAAGTATTAG 1080

5 AACTTCTTAT CTGAAGCCAT ACCACTCAGA GGGAANGCAA AATACATATT GACATCTCCT 1140
 TTAGGATGTC CTTAGAGAAT TCAAGGAAAA GAAGTTAAAT AATTTTAAAG TGCITTTGGG 1200
 TACAGCTATT TAGCACTAGA GGGTAAGATT AGACATAGAT TGTAAAGATA ATNATAGGGT 1260
 TAGGGATAGG ATTAGGATCT GGGTCAGAGT CAGGSCCAGA AGTATGGTTA GAGGTGGGGT 1320
 10 CATGGTCAGG GTSGAGATCA AAGTCAGGGT CAAAGTAAGG GTCAGAATTA GGGACCCAGG 1380
 ATAGGGATCA GGATTTAGGT TCAGTGTCAG AGTCTTGGGA CAAGGTTAGG GTTAGAATTA 1440
 15 GAACCAGAGC TTGTGTTCTCC TCAGGACCCA CCCGAGGGTG GGTCAACCATG GCTTTGGAGC 1500
 GCCTGGTAGT GTGGTGTGTC CACAGKGAAG ACCAGAGTTT CATTGTCCCT AAGACTGACY 1560
 TGGGGAGATG TGGCTGTAGS CCATTGAGGA AGGTGAGGCA ACAGCTTCCT GTCTGCTYCC 1620
 20 CCGTGTGCTG AGGAGGGAGT TCTGCCATGG GCTTTACTTT CACATGTTAT ATTCCACAAG 1680
 TCTTGTTTTA CAAAAGCATC CCTTCCTTGA GGCTTCGGCT GCTCATCGCT GCTCATCATM 1740
 25 ATAGCGTGCC ATAACATATA GTAAGATTTG GGTGTTGTTT TGGGGAGATA TCTTGGTATA 1800
 GAGAAAGGAG AAATGCTTAG AGCCACCATC AGGACAGTTG GGATGAAAGT TGGGTATAGG 1860
 CAGAGGCTGG AGGAAACATG TGCATCCCCT GTAAACACTT TTATTCATGT TTAAATTACT 1920
 30 CATTTTCTT ACAGTGTTAA ATTAGTAAAG ATAGTATTGA AAA 1963

35 (2) INFORMATION FOR SEQ ID NO: 97:

(i) SEQUENCE CHARACTERISTICS:

- 40 (A) LENGTH: 1052 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 97:

45 TCATTAACTT CAGACAACAT CATAAGCAA TGATAGCTCT TTTCTTTGTG ACCACAAYCT 60
 TAACTTGAGC TTGCTGGGT GTTTGCACA TAACAATGAG GGAATATTAG ACATAACATA 120
 ATTTTCATAG GTCATTGCC TGTCATGAT AGAGAAGATA ATTGCMAGAK AGTTWATTTT 180
 50 TGGTGTGIGT ATATGTGCAC AAATGTGCAG GGCCTCTACT TTGCAACTGG AATTATAGA 240
 CTAATGATAA AATATATCCC TTAAATATA CAAATGACAA TTGACTTCAA ACTTTCCCAA 300
 55 GCCACATAG AAATCCCTG AAAACATATA AAATATTGAG TTCTTCAACC TCAGCACTAT 360
 TGACATTTTG GACCARATAG TTCTGTWTGT KAAAGGCKGT CTTGCACTG TAGAATGTTT 420
 60 AGCAATATTC CAGGCCTCTA TCCACCTGAT ACCGGGCTG TATCCCCCTG ATACTGGTAG 480

TTCTTTTTC CCCCATCACA AATTGTGACA ACCCAGAAAT ATCTCCTTAT ACCTTTCCAG 540
AATGTTTTCC CTGGGGGACA AAAAGCACTC CCATTGAAAA ATCCACTGGT CCCAAATGGT 600
5 TAAAAATGG TTCCTTCCC ATTCTTTTA CCAGGTTGG GGCCAAGCCC CCTTCCCTTA 660
ATTTCCTCC CGAAATGAAC TGAAACCCAA CTGTWACTCT TAATGAAATA TTGAAGGKTT 720
GAAGCTTTAA AAAAAAAAAA AAAAKTACAG CTTGGCTGGG TGCAGTGGCT CAAGCCTGTA 780
10 ATCCTAGCAC TTTCGGAGGC CAAGGTGGGC AGATTGCCTG AGCTCAGGAG TTCGACACCA 840
GCGTGGGCAA CATGGTGAAA CTCGTCTCT ACTAAAATAC AAAAAGTTAA CCTGGCATGG 900
15 TGGCAGGTGC CTGTAGTCCC AGCTACTAGG GAGGCTGAGG CAGGAGAATT GCTTGAACCC 960
AGGAGGCAGA GGTGTCAGTG AGCCAAGATT GCCACTGCAC TCCAGCCTGG GCAACATAGC 1020
AAGACTCTGT CAAAAAAAAA AAAAAAATC GA 1052
20

25 (2) INFORMATION FOR SEQ ID NO: 98:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 929 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
30 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 98:

ATCCATCACA GCCTTTCTAT CTAGGCCACA CTATAAAATC TGGAGACCTT GAATATGTGG 60
35 GTATGGAAGG AGGAATGTCT TTAAGTGTAG AATCAATGAA AAGACTTAAC AGCCTTCTCA 120
ATATCCCAGA AAAGTGTCTT GAACAGGGAG GGATGATTTG GAAGATATCT GAAGATAAAC 180
40 AGCTAGCAGT TTGCCTGAAA TATGCTGGAG TATTTCAGAA AAATGCAGAA GATGCTGATG 240
GAAAAGATGT ATTTAATACC AAATCTGTG GGCCTTCTAT TAAAGAGGCA ATGACTTATC 300
ACCCCAACCA GGTAGTAGAA GGCTGTGTT CAGATATGGC TGTTACTTTT AATGGACTGA 360
45 CTCCAAATCA GATGATGTG ATGATGTATG GGTATACCG CCTTAGGGCA TTTGGGCATA 420
TTTTCAATGA TGCATTGGTT TTCCTACCTC CAAATGGTTC TGACAATGAC TGAGAAGTGG 480
50 TAGAAAAGCG TGAATATGAT CTTGTATAG GACGTGTGTT GTCATTATTT GTAGTAGTAA 540
CTACATATCC AATACAGCTG TATGTTTCTT TTTCTTTTCT AATTGGTGG CACTGGTATA 600
ACCACACATT AAAGTCAGTA GTACATTTTT AAATGAGGGT GGTTTTTTTC TTAAAACAC 660
55 ATGAACATTG TAAATGTGTT GGAAAGAAGT GTTTTAAGAA TAATAATTTT GCAAATAAAC 720
TATTAATAAA TATTATATGT GATAAATCT AAATTATGAA CATTAGAAAT CTGTGGGGCA 780
60 CATATTTTGG CTGATTGGTT AAAAAATTT AACAGGTCCT TAGCGTCTA AGATATGCAA 840

ATGATATCTC TAGTTGTGAA TTTGTGATTA AAGTAAACT TTTAGCTGTG TGTTCCTTT 900
ACTTCTGATA CTGATTATG TTNTAACCG 929

5

(2) INFORMATION FOR SEQ ID NO: 99:

10

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 359 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 99:

ATNGGANTCC CCCCGGCTG CAGGAAATTC CCCGGGCTGC ATGTCTAGTT CCACTCTGCA 60
CTGGAAAGAA TTCAAATATG CACCTGGCTC CCTTCACTAT TTTGCCCTAT CCTTTGTGCT 120
CATTCTTACT GAAATCTGTC TTGTCAGCTC AGGAATGGGA TTCCCCCAGG AAGGAAAGCA 180
CTTTTCTGTT CTGGGAAGCC CAGACTGTTT ACTTTGGGGC AGGGACGAAC ATGTGCCTCG 240
TGAATTTGCT TGAACACAGT CACCATCTTC TACCCCCATC ACTGTATAGT GAAAAACCTG 300
ATTAAAGTGG TATCTGAGAA CCAWAAAAAA AAAAAAAAAA ANCTCGAGGG GGGGCCCGG 359

30

(2) INFORMATION FOR SEQ ID NO: 100:

35

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 952 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 100:

GAATTCCTCCG GGGGATCAGG GCAGCCGGGG AGGTGGCCAG GCCAGTGGCA GGCCTGTGGA 60
GACAATCCCT YAGGACTAGG GACAGGGCTG TGCCGGCCTG GGCCAGGGCC CACGGACCCG 120
CAGCTCAGGG CGCTGCCCA CGTCGTCTGC CGGCGGTGCG CGCGGGCGT CCCTCGCGTC 180
TCTTCACTGC ACATTGCAAT GCATTTCGA TTCCCATTTT TCTGTAGGA GCCAGCCTGG 240
GTTGGCGCTG CTCCAGAGC CGTGGGTCC CAAGANCTTG CGTTCCCTTT TGTTCCTGTC 300
CCGTTTATCA AGAACACGGG CCCACCTGT TCACGTTGCC CGAAGGCCAC CCCAAGCCCA 360
ASCCTGCGGG GCGGTTCCCM MAYTGCCYTG RAATGCCCGG CTTNAAGTTY TTGCGCAACG 420
CMAGGAATTC AGTGTGGGGA CGCCCCCTGC CGGATTAGGC YTAGCCCTGG CCCAGGTGGT 480
GAGCGGTTTG CAGTGTCCGT TCTCATCCAC CTGATGGGCC CAGATAAAGG CCCCCGCTGT 540

55

60

5 CCAGCCTCCC TGGACGGCCC TCGCGGTCCC TGCAGCCCAA GATGGGACTC AGACCCCTGTG 600
 CCCCAGAGCT CCCCTGCCGC AGAATGGGCG CCCAGCCGGC CCCGACGGG TCCAGGAGCA 660
 CTGCTCGCCT GTACATACTG TTGCCCTAGC CCACCTGGTG CCGTGGGAGC CACCCCCAGG 720
 TGCNIGGCAC AGCCCCCTCC CACTCCGCCA CGCCCCCACC CACCCCGCGT GTTCTGCCC 780
 10 TGTGACTCCT GGAACCTGCG TCCTCCCCAA AGCCATGGGA GGGGTGTCCT CCTCAGACCA 840
 TGCCCCAGA TGATTTTTTT AAATAAGAA ACAAAATGCAC CTGCAAAAMA AAAAAAAAAA 900
 15 AAAAAAATC GAGGGGGGGC CCGGTACCCA ATTCGCCCTA TAGTGAGCGA TT 952

20 (2) INFORMATION FOR SEQ ID NO: 101:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 1545 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 101:

30 GAAAGACAAA AGGAAATAGA AGAAAGGGAA AAAAGGCGTA AAGACAGACA TGAAGCAAGT 60
 GGGTTTGCAA GGAGACCGAG ATCTCCAACC GGACCTAGCA CGGTGGCGCA CAAGATCATG 120
 CAGAAGTACG GCTTCCGGGA GGGCCAGGGT CTGGGGAAGC ATGAGCAGGG COTGAGCACT 180
 35 GCCTTGTCAG TGGAGAAGAC CAGCAAGCGT GGGGCAAGA TCATCGTGGG CGACGCCACA 240
 GAGAAAGGTG TGTCCCAGG GAAGCGTGTG ACTAGAGGGA AAGGACTGGC CCCATCCATA 300
 TCAGACATGG CCACTCTTGA TCCTCATGTG TCAGCAGGGG GACAATGAGG CGTGTGGCCA 360
 40 GAGGGAGAGG GCTGGCCCTG CCATCACTAG AACACAGGCC GTCCTGTTC AATGATGCAC 420
 TGCCACTTCC GTTTTGTGAA ACCAGGAATC CTGAGGCTCA TCTTTATTTT TTCAGAACAG 480
 45 ACGTAGAGAG ATGAAGGCTT GTGGAGGAAA AGATGGTGAG AGACTTGGGC AGAAAATGAG 540
 TAGTCTCAG GAAGAAATCT TGGTTATGTG TTTAGAGCAT GAAGGACAGA GCCATATAGT 600
 GTGGCAGTGA ATATACCTGC TATCTCCATC TCAGAGGTG TCTCTACTTT TCCCTTTTGC 660
 50 CCTTTCAGTA TAGATGTGAT TTCTGATTCT CTTACAGATT GTTGTCTTTG CGAGATCTGA 720
 TGTATGTTG CAGTCTCTTG GTAAATGATG CCTAGTTGGT GTTTTATTTT CATTTAATTT 780
 55 TTACAGTCTG TTCTGTGTTG AGGGAATTCA GGAAAGAGAC AAACATATGT TAGCATTTTA 840
 ATCAGGGAAT TAAGTTTGAG TCAGCCTAGC TGAACCTCCT TGGCTAAAGA AAGAAGAAAA 900
 60 CTTTCTGGC AGCCCCGTTT ATGCACAGCT TAGGATACAT CACGAGCCTG ACAGATGCAT 960

CCAAGAAGTC AGATTCAAAT CCGCTGACTG AAATACTTAA GTGTCCTACT AAAGTGGTCT 1020
 TACTAAGGAA CATGGTTGGT GCGGGAGAGG TGGATGAAGA CTGGGAAGT TGAAACCAAG 1080
 5 GAAGAATGTG NAAAAATATG GCAAAGTTGG AAAATGTGTG ATATTTGAAA TTCTGGTGC 1140
 CCCTGATGAT GAAGCAGTAC GGATATTTT AGAATTTGAG AGAGTTGAAT CAGCAATTAA 1200
 10 AGCGGTGTT GACTTGAATG GGAGGTATTT TGGTGGACGG GTGGTAAAAG CATGTTTCTA 1260
 CAATTTGGAC AAATTCAGGG TCTTGGATTT GGCAGAACAA GTTTGATTTT AAGAACTAGA 1320
 GCACGAGTCA TCTCCGGTGA TCCTTAAATG AACTGCAGGC TGAGAAAAGA AGGAAAAAGG 1380
 15 TCACAGCCTC CATGGCTGTT GCATACCAAG ACTCTTGAA GGAATCTAA GATATATGTT 1440
 GATTGATCCC TTTTATTTT TGTGTTTTT TAATATAGTA TAAAAATCCT TTTAAAAAA 1500
 20 CAAMAAAAAA AAAAAAACT CGAGGGGGG CCCGGTACCC AATTT 1545

(2) INFORMATION FOR SEQ ID NO: 102:

25

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1322 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

30

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 102:

CTCTGGGAG CGACCGCTCC GCTCGTCTCG TTGTTCCGG AGGTCGTGC GCGGTGGGA 60
 35 AATGCTGGCG CGCGCGGCG GNGGCACTGG GGCCCTTTTG CTGAGGGGCT CTCTACTGGC 120
 TTCTGGCCGC GCTCCGCSG CGCTCCTCT GGATGCCCC GAAACACCGT GGTACTGTT 180
 40 GTGCCGCAGC AGGAGGCTG GGTGGTGGAG CGAATGGGC GATTCCACCG GATCCTGGAG 240
 CCTGGTTGA ACATCCTCAT CCCTGTGTTA GACCGGATCC GATATGTGCA GAGTCTCAAG 300
 GAAATTGTCA TCAACGTGCC TGAGCAGTC GCTGTGACTC TCGACAATGT AACTCTGCAA 360
 45 ATCGATGGAG TCCTTTACCT GCGCATCATG GACCCTTACA AGGCAAGCTA CGGTGTGGAG 420
 GACCCTGAGT ATGCCGTCAC CCAGCTAGCT CAAACAACCA TGAGATCAGA GCTCGGCAAA 480
 50 CTCTCTCTGG ACAAGTCTT CCGGAACGG GAGTCCCTGA ATGCCAGCAT TGTGGATGCC 540
 ATCAACCAAG CTGCTGACTG CTGGGTATC CGCTGCCTCC GTTATGAGAT CAAGGATATC 600
 55 CATGTGCCAC CCCGGGTGAA AGAGTCTATG CAGATGCAGG TGGAGGCAGA GCGGCGGAAA 660
 CGGGCCACAG TTCTAGAGTC TGAGGGGACC CGAGAGTCGG CCATCAATGT GGCAGAAGGG 720
 AAGAAACAGG CCCAGATCCT GGCCTCCGAA GCAGAAAAGG CTGAACAGAT AAATCAGGCA 780
 60 GCAGGAGAGG CCAGTGCACT TCTGGCGAAG GCCAAGGCTA AAGCTGAAGC TATTCGAATC 840

CTGGCTGCAG CTCTGACACA ACATAATGGA GATGCAGCAG CTTCACTGAC TGTGGCCGAG 900
5 CAGTATGTCA GCGCGTTCTC CAAACTGGCC AAGGACTCCA AACTATCCT ACTGCCCTCC 960
AACCCTGGCG ATGTCACCAG CATGGTGGCT CAGGCCATGG GTGTATATGG AGCCCTCACC 1020
AAAGCCCCAG TGCCAGGGAC TCCAGACTCA CTCTCCAGTG GGAGCAGCAG AGATGTCCAG 1080
10 GGTACAGATG CAAGTCTTGA TGAGGAAGTT GATCGAGTCA AGATGAGTTA GTGGAGCTGG 1140
GCTTGGCCAG GGAGTCTGGG GACAAGGAAG CAGATTTTCC TGATTCTGGC TCTAGCTTCC 1200
15 CTGCCAAGAT TTTGGTTTTT ATTTTTTTAT TTGAACTTTA GTCGTGTAAT AAACTCACCA 1260
GTGGCAAACC AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAAN 1320
NN 1322
20

(2) INFORMATION FOR SEQ ID NO: 103:

25 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 276 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
30 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 103:

NNATAGCTCA ACCATGTTCC AGGAGTGTAT TCCAATCAGC TTGTTTTTTC TTAAGTGGTT 60
35 AAAGGAATGT TGCTCATTCA CCTGCCCCAA CTCACATATT AACAATGTGT TAACTGGGAT 120
TAGATAAAAG GAAAGCTGAC TTACAGATGA ACCAAGAGGG AGCTATTTAT GCCACAGCCC 180
40 CCAGCCCAGT AACTTTATGT TTCTGATCTC CTGCAAAATT TTTTATAAA AAAAGCTTAG 240
CCAGGAACTA GTAGAAAGAA TAAAGTAAAG ATGGTG 276

45 (2) INFORMATION FOR SEQ ID NO: 104:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 381 base pairs
50 (B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 104:

55 GATTAAGGTA GAAAAGTACA GAAAACACTA AATTTTCATT GTGCTGTTTC AATGTGGCAG 60
ATTCTTTAAA ATACTTCGAC ACGCTACAAT AATTAAAGGT TTTAAGAACA TTAAGATACT 120
60 TAAAAAATAA AAGCCACAA TTGAATAACA AAAATGAACT TTGTTTTATT TTTTATTGGC 180

ATTAATGTAG GTTGCCGTGG TGAAAATAGT TTGAAATACT TCACAGTAAC AGTTTTKTGC 240
AGCCCTAGAG ATTA AAAACA GCAAAGTAAA TAAGCAGGAC TCTCAACGAC TCATACTCAC 300
5 AGACTGTTTA ATGTWATCCT ARCACTTCSG GARGCTGARG CGGGAGGATT ACTTGAGCCT 360
AGGATTGAG ACCAGCCTGG G 381

10

(2) INFORMATION FOR SEQ ID NO: 105:

15

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 638 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 105:

TGTGGAAAAC AGTAGGAAAG CAATGAAAGA AGCTGGTAAG GGAGGCGTCG CTGATTCCAG 60
25 AGAGCTAAAG CCGATGGTAG GTGGAGATGA RGARGTGGCC GCCCTCCAAG AATTTCACTT 120
TCACTTCCTC TCTCTCTCTG TCTTCACTGA CTGCACCTTCT TCAGGAGAAG CTTTGTGTAT 180
CTGTATCACG CAGACATGCT GCTCTTTCTG TTTGTGTGCT TACCCATCAC TTGGATGGCA 240
30 GAATTCTTGT CACAACAGAG ACACCTYCTA TAAAAGTAAG CTGAAAGGAA CAGCATCCTC 300
GTCAGTGCTC GGCAGGGGCG GGTAGGGGAT GATGGTTTTT TCCCTAAGGT AAAACTGCTG 360
35 TTGCTCTTGT TTCCTTTTTA ACTGTCAGTG TTTGGCTTTC ATCAGACTGA ACATTTTGGT 420
GTACACTTGA ACTGACGGTT TGATTTTTAT CATTTTGGAA GGTGATCATA GCAATTCCTT 480
TCAACTTGCT AAAATTCATA CTCCTCTTTT TAAAAGTATG GTTCTGCTTA CATTCGCTGC 540
40 CTTTCCCTT GGTGACTTT TTCTTCTGTT GCCTAGGTTG TACTTTTTTN TTTTNT 600
TTTTCAGTAG CAAACAAGGC TGTTCATC AATACCCA 638

45

(2) INFORMATION FOR SEQ ID NO: 106:

50

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 2246 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 106:

GGCAGGAGGC CGGGGAGAG TCACGCAAAT GACTTGGAGT GTTCAGGAAA AGGAAAATGC 60
60 ACCACGAAGC CGTCAGAGGC AACTTTTCC TGTACCTGTG AGGAGCAGTA CGTGGTACT 120

	TTCTGTGAAG AATACGATGC TTGCCAGAGG AAACCTTGCC AAAACAACGC GAGCTGTATT	180
5	GATGCAAATG AAAAGCAAGA TGGGAGCAAT TTCACCTGTG TTTGCCCTTCC TGGTTATACT	240
	GGAGAGCTTT GCCAGTCCAA GATTGATTAC TGCATCCTAG ACCCATGCAG AAATGGAGCA	300
	ACATGCATTT CCAGTCTCAG TGGATTACCC TGCCAGTGTG CAGAAGGATA CTTCGGATCT	360
10	GCTTGTGAAG AAAAGGTGGA CCCCTGCGCC TCGTCTCCGT GCCAGAACAA CGGCACCTGC	420
	TATGTGGACG GGGTACACTT TACCTGCAAC TGCAGCCCGG GCTTCACAGG GCCGACCTGT	480
15	GCCCAGCTTA TTGACTTCTG TGCCCTCAGC CCCTGTGCTC ATGGCACGTG CCGCAGCGTG	540
	GGCACCAGCT ACAAATGCCT CTGTGATCCA GGTACCATG GCCTCTACTG TGAGGAGGAA	600
	TATAATGAGT GCCTCTCCGC TCCATGCCTG AATGCAGCCA CCTGCAGGGA CCTCGTTAAT	660
20	GGCTATGAGT GTGTGTGCCT GGCAGAATAC AAAGGAACAC ACTGTGAATT GTACAAGGAT	720
	CCCTGCGCTA ACGTCAGCTG TCTGAACGGA GCCACCTGTG ACAGCGACGG CCTGAATGGC	780
25	ACGTGCATCT GTGCACCCGG GTTTACAGGT GAAGAGTGGC ACATTGACAT AAATGAATGT	840
	GACAGTAACC CCTGCCACCA TGGTGGGAGC TGCCCTGGACC AGCCCAATGG TTATAACTGC	900
	CACTGCCCCG ATGGTTGGGT GGGAGCAAAC TGTGAGATCC ACCTCCAATG GAAGTCCGGG	960
30	CACATGGCGG AGAGCCTCAC CAACATGCCA CGGCACTCCC TCTACATCAT CATTTGGAGCC	1020
	CTCTGCGTGG CCTTCATCCT TATGCTGATC ATCCTGATCG TGGGGATTTG CCGCATCAGC	1080
35	CGCATTGAAT ACCAGGGTTC TTCCAGGCCA GCCTATGAGG AGTTCTACAA CTGCCGCAGC	1140
	ATCGACAGCG AGTTTCAGCA TGCCATTGCA TCCATCCGGC ATGCCAGGTT TGGAAAGAAA	1200
	TCCCGGCCCTG CAATGTATGA TGTGAGCCCC ATCGCCTATG AAGATTACAG TCCTGATGAC	1260
40	AAACCOCTTG TCACACTGAT TAAAACTAAA GATTGTGAAT CTTTMTTGG ATTATTTTTC	1320
	AAAAAGATGA GATACTACAC TCATTTAAAT ATTTTAAAGG AAATTAAGAA GCTTAAGAAA	1380
45	TTTAAATGC TAGCTGCTCA AGRGTTTCA GTAGAATATT TAAGAACTAA TTTTCTGCAG	1440
	CTTTTAGTTT GGAAAAATA TTTTAAAAAC AAAATTGTG AAACCTATAG ACGATGTTT	1500
	AATGTACCTT CAGCTCTCTA AACTGTGTGC TTCTACTAGT GTGTGCTCTT TTCACTGTAG	1560
50	ACACTATCAC GAGACCCAGA TTAATTCTG TGGTTGTAC AGAATAAGTC TAATCAAGGA	1620
	GAAGTTTCTG TTTGACGTTT GAGTGCCGGC TTTCTGAGTA GAGTTAGGAA AACCACGTAA	1680
55	CGTAGCATAT GATGTATAAT AGAGTATACC CGTTACTTAA AAAGAAGTCT GAAATGTTTCG	1740
	TTTTGTGGAA AAGAACTAG TTAAATTAC TATTCCTAAC CCGAATGAAA TTAGCCTTTG	1800
	CCTTATTCTG TGCATGGGTA AGTAACCTAT TTCTGCACTG TTTTGTGAA CTTTGTGGAA	1860
60	ACATTCTTTC GAGTTTGTTC TTGTCAATTT CGTAACAGTC GTCGAAC TAG GCCTCAAAAA	1920

CATACGTAAC GAAAAGGCCT AGCGAGGCAA ATTCTGATTG ATTTGAATCT ATATTTTCT 1980
 5 TTAAAAAGTC AAGGGTTCTA TATTGTGAGT AAATTAAATT TACATTTGAG TTGTTTGTG 2040
 CTAAGAGSTA GTAAATGTAA GAGAGTACTG GTTCCTTCAG TAGTGAGTAT TTCTCATAGT 2100
 GCAGCTTTAT TTATCTCCAG GATGTTTTTG TGGCTGTATT TGATTGATAT GTGCTTCTTC 2160
 10 TGATTCTTGC TAATTTCCAA CCATATTGAA TAAATGTGAT CAAGTCAAAA AAAAAAAAAA 2220
 AAAAAAATT ACTCGGTGCG AAGGGA 2246

15

(2) INFORMATION FOR SEQ ID NO: 107:

20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1105 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: double
 (D) TOPOLOGY: linear

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 107:

GAATTCGGCA GAGCCCACTT AGAGGAGCTA AAATAGCTAA AGGTTACATG CTTTGCCTCA 60
 AATAATAGAC TTAGTGAAGA GGGTAGAAGT AGAAATRAGG TCAGCCCCC AGAGCAGTCT 120
 30 GGTTGGCCTT AGCAACCAGG AAGGTAAAGC CGGTACCTCA GTTAAATCAC CAAGTTTACT 180
 GGAAGTGCAT ATTTTTCATG TGCCAAATTC AGTAAGTCAT GGAGCAAATG TTTATTTTGC 240
 35 TATGCTTTAA AAAGTTGCTT GCTTCTTGTA AGTTTCTCA GTGGAAGGGT TCCAAGTTAT 300
 GACTTAATCT ATGTTTGCAG CATTGCACTG GAAACAGGAT TTGTCTGTGA AATGGCTCTG 360
 TCATTTGTGG ACCACTTCTG TAGGGAGATT GTGGATTAG GAAGGGCAGA AGCAACAGCA 420
 40 GATATGCCTG GTGTTTGAAT GGATGTGCCT CTYTGGAGG CAGCAAGCAG CATACCCATA 480
 TTATAAAGTT TTTGATTTTC TAACATCTGA AGACAGGCAT CCAGCCTTGC AGAACAGCCA 540
 45 GGTGTCTGTT CTATAGACTA CAGTTCCTTG TTTCCAGAAT TACGGTAACC AAATAATACA 600
 CAAGGTCACC TGATTGCACT TCCAACAAC CTGAACAAAG AGCACCTTTC CGCTTGCTGG 660
 TAGGTGCTGT ACCAGACTCT TTGTAATCTG CCTTAGKTCA GRGAAGAACA AGCCATTACC 720
 50 AGTATGGGAG TCCATCCYTA GTCAGGGCTA GTTGCTAITA TCCCTTGAAT ACTCTGCAGG 780
 CATCCACAA GACATTTGAG ACTTCATATT TGTCAAATAA TAGAAATSTG GCTGGCCTAG 840
 55 TGGCTCATGC CTGTAATCCT AACCTTTGG GAGGCTGATG TGGGCAGATT GCTTGAGGCC 900
 AGGAGTTTGA GACCCACCTG GGCAACACAG TGACATGTTG TCTCTACAAA AAATTTAAAA 960
 ATTAAGTAGG CATGGTAGTG TGCCTATAGT CCCAGCTACT CCAGAGGCTG AGGCAGGAAG 1020
 60

ATCCCTTGAG CCCAGTAATT CAAGGCTACA GTTAGCTCTG ATCCTGCCAC TGCACTCCTG 1080
TCCTGGTAAA GGAGCTAAAC CCACT 1105

5

(2) INFORMATION FOR SEQ ID NO: 108:

10

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 505 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 108:

ATTTACACA GGAAACAGCT ATGACCATGA TTCCGCCAAG CNGGAAATTA ACCNTCACTA 60
AAGGGAACAA AACTGGAGCT CCACCGCGGT GCGGCGCGCT CTAGAACTAG TGGATCCCCC 120
GGGCTCAGGA ATTCCGCACG AGTTCTTCCA CATGTGTGCA CCCCCAGCTT GGCCAACCTT 180
CAGCCTTGCG GTGGGGCCCG AAGCATCTTC CCTTCCGCTT GCGGTCTCTG GGATTGGGAT 240
GAGTGCCTGG CTCCCATCTC CTCCTCACTT TTTGTGTGTA TCGGCAGCTG CTGGCTCAGG 300
GGCATCCAC CTCCGGGCTC TGGGTTCTCT TGCCCTGGAA GGGCTCCAGG ACCCGTCCCA 360
ATAACCACCC ACGGCCAGGA GRGCCAAGGC CCCGTGCTGG ATATTTAAAT TTAGGGGCCG 420
GTCTCCAGGG CCGGTAGATA AATAAATACA CTCAGCGTCA AAAAAAAAAA AAAAAAAAAA 480
AAAAAAAAA AAAAAAAAAA CTCGA 505

35

(2) INFORMATION FOR SEQ ID NO: 109:

40

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1380 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 109:

AATCATGAGC CTCCAGAAGA GACAGATGGC CCACCAGGAG CTGTTGCTCT GGTTCGCTTC 60
CTGCAGGCCT TGGAGAAGGA GGTGCGCATA ATCGTTGACC AGAGAGCCTG GAACTTGCAC 120
CARAAGATTG TTGAAGATGC TGTGAGCAA GGTGTTCTGA AGACGCAGAT CCCGATATTA 180
ACTTACCAAG GTGGATCAGT GGAAGCTGCT CAGGCATTCC TGTGCAAAA TGGGGACCCG 240
CAGACACCTA GATTTGACCA CCTGGTGGCC ATAGAGCGTG CCGGAAGAGC TGCTGATGGC 300
AATTACTACA ATGCAAGGAA GATGAACATC AAGCACTTGG TTGACCCCAT TGACGATCTT 360

60

	TTTCTTGCTG CGAAGAAGAT TCCTGGAATC TCATCAACTG GAGTCGGTGA TGGAGGCAAC	420
	GAGCTTGCGA TGGGTAAAGT CAAGGAGGCT GTGAGGAGGC ACATACGGCA CGGGRATGTC	480
5	ATCGCCTGCG ACGTGGAGGC TGACTTTGCC GTCATTGCTG GTGTTTCTAA CTGGGGAGGC	540
	TATGCCCTGG CCTGCGCACT CTACATCCTG TACTCATGTG CTGTCCACAG TCAGTACCTG	600
	AGGAAAGCAG TCGGACCCCTC CAGGGCACCT GGAGATCAGG CCTGGACTCA GGCCCTCCCG	660
10	TCGGTCATTA AGGAAGAAAA AATGCTGGGC ATCTTGGTGC AGCACAAAGT CCGGAGTGGC	720
	GTCTCGGGCA TCGTGGGCAT GGAGGTGGAT GGGCTGCCCT TCCACAACAC CCACGCCGAG	780
15	ATGATCCAGA AGCTGGTGGG CGTCACCACG GCACAGGTGT AACCGTCCAT GTTCCGTGTG	840
	AGCAGAGTCC CTACCAACGG GCAGGTCTGC ATCCGGGGAG AATGCAGCTG CTTCTGGCGA	900
	CAATCCTGCT AGTAAACACT GGTCTTCGGT GAGCAACGAA CACTCGCCTG GCCTGGGAAA	960
20	CTGCATGCCC ACTTTCTGGG AGGGTTAGT GCAGGTGCCG TGGACAAAGG ACAACATTTC	1020
	TCTGGGGCTT TTTAACTTTT ATTCCTAAGA CTCTAAAGGC GTTGATTICA ACCCTCCTTC	1080
25	ACTCTGGCTT CTTCAGGCAA CCCACGTGGT CTCCTGTGAG AATCTTCTCG ACAGTTACTT	1140
	ATGGGGACAC TTGTGAACAA TTAACGCCA GGCAGAGCAT GAGAACAAC ATTCCCAGGC	1200
	CATGTAGGAT AGGATACTCC AGACTCCAGT CATCCTCCCC CATCCATGGT TTCTGTACT	1260
30	CATGGTTTCA GTTACTCATA GCCAACTGCA GACCGAAAAT ACTAAATGAA AAATTTGAGA	1320
	AATAACAAC TCTTAAGTTT TAAAAA AAAA AAAA GGGCGGCCGC	1380
35		

(2) INFORMATION FOR SEQ ID NO: 110:

- 40 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 646 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: double
 - (D) TOPOLOGY: linear
- 45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 110:

	CAGATGCCAG GGACTTGNC TTCCTCCCGT TGAACCACAG GTTCCAAGAA ACCTGCAGGG	60
50	TCCAGCCTCC CCCCCATCC CAGTYTTC CACCCTGGC CGGCCCTCCA GGTGCAGAAA	120
	CATGCAGGCC CCTCTCCAGG ACTGTGGAG GAGTGTGTC CTCAGACTGG CCTGTGTCCT	180
	GGCTCCTCTT ACCACCTCTT CCAGAGGTTG TCACCTGCAG CTGCCCCAGG ATAAAGGCAA	240
55	GGCCAGARAG GACTCCTGAA CTCCTGTGTG CCTGGGGTGG CAGGGGCAAA CATAGCCAAC	300
	TGGTGGCCTG AGCGGGGCCA TGGTGARGAC ACCCTTGGTG GCTTGTCCTA CATCAAGCTG	360
60	GGARGTGACA CTTAGGATGC ATTTTCAAT ATTTTAGTGT TTGAATAACG GGCTAWCTTG	420

5 AGAAAAAAT AATTTGAATC ACACATCACA CCAAAAATAA ATTCTAGGTG GATTTTAACA 480
 CTTTCCAAAA ATTATTATTA GTTTAGAGAC AGGGTCTCAC TCCGTCGCCT AGGCTGGAGT 540
 GCANGGGTAT GATCATGGTT CACTGCAACC TTAAACTCCC TGGCCTCATA TGATCCCCCC 600
 GGGCTCCAGC CCTTCCAAAG TTACTGGGAA ACTACCAAAC ATGCCC 646

10

(2) INFORMATION FOR SEQ ID NO: 111:

15

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 111:

20

Met Asp Ser Tyr Trp His Ser Arg Cys Leu Lys Cys Ser Cys Cys Gln
 1 5 10 15

25

Ala Xaa Trp Ala Thr Ser Ala Arg Pro Val Thr Pro Lys Val Ala Xaa
 20 25 30

30

(2) INFORMATION FOR SEQ ID NO: 112:

35

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 36 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 112:

40

Ile Tyr Ser Ser Gly Tyr Phe Gln Ile Tyr Asn Met Leu Leu Leu Thr
 1 5 10 15

Ile Leu Ile Leu Leu Cys Asn Arg Thr Pro Glu Leu Ile Pro Gly Phe
 20 25 30

45

Tyr Ile Arg Xaa
 35

50

(2) INFORMATION FOR SEQ ID NO: 113:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 220 amino acids

55

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 113:

60

Met Ser His Lys Leu Gly Asp Pro Gly Phe Val Val Phe Ala Thr Leu
 1 5 10 15

265

Val Val Ile Val Ala Leu Ile Leu Ile Phe Val Val Gly Pro Arg His
 20 25 30

5 Gly Gln Thr Asn Ile Leu Val Tyr Ile Thr Ile Cys Ser Val Ile Gly
 35 40 45

Ala Phe Ser Val Ser Cys Val Lys Gly Leu Gly Ile Ala Ile Lys Glu
 50 55 60

10 Leu Phe Ala Gly Lys Pro Val Leu Arg His Pro Leu Ala Trp Ile Leu
 65 70 75 80

15 Leu Leu Ser Leu Ile Val Cys Val Ser Thr Gln Ile Asn Tyr Leu Asn
 85 90 95

Arg Ala Leu Asp Ile Phe Asn Thr Ser Ile Val Thr Pro Ile Tyr Tyr
 100 105 110

20 Val Phe Phe Thr Thr Ser Val Leu Thr Cys Ser Ala Ile Leu Phe Lys
 115 120 125

Glu Trp Gln Asp Met Pro Val Asp Asp Val Ile Gly Thr Leu Ser Gly
 130 135 140

25 Phe Phe Thr Ile Ile Val Gly Ile Phe Leu Leu His Ala Phe Lys Asp
 145 150 155 160

30 Val Ser Phe Ser Leu Ala Ser Leu Pro Val Ser Phe Arg Lys Asp Glu
 165 170 175

Lys Ala Met Asn Gly Asn Leu Ser Asn Met Tyr Glu Val Leu Asn Asn
 180 185 190

35 Asn Glu Glu Ser Leu Thr Cys Gly Ile Glu Gln His Thr Gly Glu Asn
 195 200 205

Val Ser Arg Arg Asn Gly Asn Leu Thr Ala Phe Xaa
 210 215 220

40

(2) INFORMATION FOR SEQ ID NO: 114:

45

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 114:

Met Thr Ile Trp Glu Arg Lys Tyr Ile Trp Met Leu Gln Ile Cys Val
 1 5 10 15

55

Phe Leu Glu Pro Arg Ala Lys Pro Ser Leu Gly Asp Leu Asp Trp Xaa
 20 25 30

60

(2) INFORMATION FOR SEQ ID NO: 115:

- 5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 27 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 115:

10 Met Leu Thr Phe Leu Leu Phe Ile Pro Val Ala Pro Thr Glu Thr Ser
 1 5 10 15
 Gln Lys Asn Arg Ser Val Phe Leu Pro Pro Xaa
 20 25

15

(2) INFORMATION FOR SEQ ID NO: 116:

- 20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 132 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 116:

25 Met Leu Phe Val Phe Cys Cys Thr Val Phe Phe Val Cys Leu Phe Val
 1 5 10 15
 Tyr Leu Val Gly Phe Leu Glu Arg Glu Ile Trp Lys Arg Asp Ile His
 30 20 25 30
 Lys Ser Tyr Thr Pro Thr Phe Pro Phe Tyr His Asp Ile Gln Glu Glu
 35 35 40 45
 Thr Ser Arg Ala Lys Asn Gly Val Lys Lys Gly Ser Met Ala Gly Thr
 50 55 60
 Ser Lys Glu Leu Arg Ala Val Ala Leu Lys Asn Tyr Phe Phe Tyr Tyr
 40 65 70 75 80
 Tyr Phe Glu Ser Met Glu Val Phe His Ser Leu Gly Lys Gly Gly Lys
 85 90 95
 Ser Ala Phe Ile Phe Ile Gln Ser Tyr Leu Ile Thr Ser Lys Thr His
 45 100 105 110
 Met Leu Glu Ile Ala Phe Ala Gly Ala Lys Tyr Ile Asn Glu Gln Glu
 115 120 125
 50 Tyr Ile His Xaa
 130

55 (2) INFORMATION FOR SEQ ID NO: 117:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 65 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 117:

Met Trp Tyr Phe Met Ser Leu Ile Ser Met Val Leu Leu Leu Ser Pro
 1 5 10 15
 5 Ser Cys Ser Asp Leu Leu Val Ile Ser Val Leu Asn Leu Glu Gln Arg
 20 25 30
 10 Arg Gln Ser Lys Val Gly Phe Glu Pro Phe Thr Ser Pro Leu Cys Gly
 35 40 45
 Xaa Trp His His Leu Ser Pro Asp Arg Leu Pro Gln Asp Gly Thr Phe
 50 55 60
 15 Xaa
 65

(2) INFORMATION FOR SEQ ID NO: 118:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 118:

Leu Leu Leu Phe Cys Ile Leu Gly Xaa
 1 5
 30

(2) INFORMATION FOR SEQ ID NO: 119:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 50 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 119:

Met Gly Val Leu Phe Val Pro Gln Glu Thr Ser Xaa Lys Val Xaa Xaa
 1 5 10 15
 45 Asp Ile Xaa Gly Leu Ser Gln Phe Val Met Gly Glu Lys Arg Thr Thr
 20 25 30
 Ser Ile Arg Gly Ile Gln Ala Arg Tyr Gln Val Asp Arg Gly Leu Glu
 35 40 45
 50 Tyr Cys
 50

(2) INFORMATION FOR SEQ ID NO: 120:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 76 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 120:

5 Met Leu Leu Leu Leu Leu Leu Leu Leu Leu Leu Trp Thr Cys Gln
 1 5 10 15
 Lys Ala Leu Val Arg Arg Gln Phe Cys Leu Phe Asn Leu Ile Ala Arg
 20 25 30
 10 Asn Ser Ser Leu Met Leu Gln Lys Asp Glu Lys Lys Gly Lys Lys Arg
 35 40 45
 Asp Asn Ser Gln Ala Gln Arg Glu Lys Lys Gly Gly Gly Lys Glu Pro
 50 55 60
 15 Gln Gly Asp Leu Gln Glu Arg Pro Gly Pro Gly Xaa
 65 70 75

20 (2) INFORMATION FOR SEQ ID NO: 121:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 121:

30 Met His Asn Ala Phe Asn Leu Asn Val Leu Thr Leu Phe Leu Ser Val
 1 5 10 15
 Leu Cys Cys Thr Phe Ser Asp Ser Glu Leu Xaa
 20 25

35 (2) INFORMATION FOR SEQ ID NO: 122:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 122:

45 Met Ser Trp Leu Phe Leu Leu Phe Ala Leu Leu Cys Lys Phe Gln His
 1 5 10 15
 Lys Leu Xaa Phe His Asn Ile Xaa
 20

50 (2) INFORMATION FOR SEQ ID NO: 123:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 123:

60 Met Leu Leu Phe Leu Thr Val Ile Asn Phe Met Ala Leu Ala Lys Met

1 5 10 15

Asn Phe Cys Gly Asp Xaa
20

5

(2) INFORMATION FOR SEQ ID NO: 124:

10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 55 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 124:

Met Val Xaa Asn Leu Gln Val Ile Ser Ile Trp Xaa Xaa Ser Thr Thr
1 5 10 15

Cys Phe Tyr Ala Cys Ile Trp Xaa Gln Gly Cys Leu Met Leu Arg Xaa
20 20 25 30

Phe Xaa Thr Leu Asn Asn Val Thr Arg Leu Pro Ser Ser Gln Lys Pro
35 40 45

25 Ile Lys Cys Tyr Leu Leu Xaa
50 55

30 (2) INFORMATION FOR SEQ ID NO: 125:

 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 318 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 125:

Met Leu Ser Glu Ser Ser Ser Phe Leu Lys Gly Val Met Leu Gly Ser
1 5 10 15

40 Ile Phe Cys Ala Leu Ile Thr Met Leu Gly His Ile Arg Ile Gly His
20 25 30

Gly Asn Arg Met His His His Glu His His His Leu Gln Ala Pro Asn
45 35 40 45

Lys Glu Asp Ile Leu Lys Ile Ser Glu Asp Glu Arg Met Glu Leu Ser
50 55 60

50 Lys Ser Phe Arg Val Tyr Cys Ile Ile Leu Val Lys Pro Lys Asp Val
65 70 75 80

Ser Leu Trp Ala Ala Val Lys Glu Thr Trp Thr Lys His Cys Asp Lys
85 90 95

55 Ala Glu Phe Phe Ser Ser Glu Asn Val Lys Val Phe Glu Ser Ile Asn
100 105 110

60 Met Asp Thr Asn Asp Met Trp Leu Met Met Arg Lys Ala Tyr Lys Tyr
115 120 125

270

Ala Phe Xaa Lys Tyr Arg Asp Gln Tyr Asn Trp Phe Phe Leu Ala Arg
 130 135 140

5 Pro Thr Thr Phe Ala Ile Ile Glu Asn Leu Lys Tyr Phe Leu Leu Lys
 145 150 155 160

Lys Asp Pro Ser Gln Pro Phe Tyr Leu Gly His Thr Ile Lys Ser Gly
 165 170 175

10 Asp Leu Glu Tyr Val Gly Met Glu Gly Gly Ile Val Leu Ser Val Glu
 180 185 190

Ser Met Lys Arg Leu Asn Ser Leu Leu Asn Ile Pro Glu Lys Cys Pro
 195 200 205

15 Glu Gln Gly Gly Met Ile Trp Lys Ile Ser Glu Asp Lys Gln Leu Ala
 210 215 220

20 Val Cys Leu Lys Tyr Ala Gly Val Phe Ala Glu Asn Ala Glu Asp Ala
 225 230 235 240

Asp Gly Lys Asp Val Phe Asn Thr Lys Ser Val Gly Leu Ser Ile Lys
 245 250 255

25 Glu Ala Met Thr Tyr His Pro Asn Gln Val Val Glu Gly Cys Cys Ser
 260 265 270

30 Asp Met Ala Val Thr Phe Asn Gly Leu Thr Pro Asn Gln Met His Val
 275 280 285

Met Met Tyr Gly Val Tyr Arg Leu Arg Ala Phe Gly His Ile Phe Asn
 290 295 300

35 Asp Ala Leu Val Phe Leu Pro Pro Asn Gly Ser Asp Asn Asp
 305 310 315

40 (2) INFORMATION FOR SEQ ID NO: 126:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 59 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 126:

Met Thr Trp Pro Pro Ser Cys Leu Val Ala Leu Leu Leu Ser Thr Val
 1 5 10 15

50 Thr Gln Lys Met Thr Pro Leu Asn Leu Met Arg Thr Thr Gly Pro Ile
 20 25 30

Asn Ser Phe Cys Leu Leu Pro Thr Phe Phe Phe Phe Pro Ser Tyr Leu
 35 40 45

55 Pro Ser Leu Met Pro Thr Pro Thr Asp Pro Xaa
 50 55

60

(2) INFORMATION FOR SEQ ID NO: 127:

5 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 99 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 127:

10 Ile Leu Phe Ser Phe Leu Ile Pro Ser Asn Leu Ser Phe Ser Pro Val
 1 5 10 15

Ile Phe Phe Leu Cys Gly Pro Phe Lys Val Val Ile Ile Cys Thr Glu
 20 25 30

15 Leu Gln Asn Val Ser Arg Ser Pro Gln Thr Thr Leu Ala Thr Val Tyr
 35 40 45

20 Cys Asn Lys Ile Thr Ser Tyr Ile Cys Arg Asn Ser Phe Gly Val Ile
 50 55 60

Leu Phe Phe Pro Leu Asn Ile Tyr Asn Trp Thr Asn Ala Gly Lys Lys
 65 70 75 80

25 Lys Lys Met Val Ser Lys Lys Pro Lys Ile Lys Phe Arg Gly His Gln
 85 90 95

Ala Phe Xaa

30

(2) INFORMATION FOR SEQ ID NO: 128:

35 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 29 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 128:

40 Met Ser Ile Leu Leu Leu Xaa Phe Pro Ser Ala Pro Ala Pro Val Val
 1 5 10 15

45 Ser Gly Gly Leu Gln Pro Trp Leu His Ser Cys Ile Xaa
 20 25

50 (2) INFORMATION FOR SEQ ID NO: 129:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 129:

Met Gly Thr Ser Leu Asn Leu Gln Ile Met Ala Leu Phe Ser Gly Gln
 1 5 10 15

60 Ala Met Ala Pro Arg Xaa

20

5 (2) INFORMATION FOR SEQ ID NO: 130:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 112 amino acids

(B) TYPE: amino acid

10 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 130:

Met Leu Trp Leu Pro Leu Leu Ala Ala Leu Ser Pro Ser Pro Pro Gly
 1 5 10 15
 Val Ser Ser Glu Glu Glu Gln His Trp Ser Gln Ala Glu Ala Leu Pro
 20 25 30
 Cys Trp Asp Pro Gly Ser Glu Ser Ser Pro Arg Ile Pro Gly Cys Arg
 35 40 45
 Glu Leu Gln Ser Cys Pro Pro Pro Thr Ala Pro Ser Ala His Thr Gln
 50 55 60
 Ser Pro Gly Gly Leu Gly Ala Lys Ala Gly Ala Ala Leu Val Pro Phe
 65 70 75 80
 Pro Gly Pro Ser Phe Pro Thr Ser Lys Pro Lys Lys Gly Glu Ala Gly
 85 90 95
 Ala Pro Val Pro Gln Pro His Ser Ala Leu Thr Val Pro Ser Ser Xaa
 100 105 110

35

40 (2) INFORMATION FOR SEQ ID NO: 131:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 114 amino acids

(B) TYPE: amino acid

45 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 131:

Met Glu Lys Pro Leu Phe Pro Leu Val Pro Leu His Trp Phe Gly Phe
 1 5 10 15
 Gly Tyr Thr Ala Leu Val Val Ser Gly Gly Ile Val Gly Tyr Val Lys
 20 25 30
 Thr Gly Ser Val Pro Ser Leu Ala Ala Gly Leu Leu Phe Gly Ser Leu
 35 40 45
 Ala Gly Leu Gly Ala Tyr Gln Leu Tyr Gln Asp Pro Arg Asn Val Trp
 50 55 60
 Gly Phe Leu Ala Ala Thr Ser Val Thr Phe Val Gly Val Met Gly Met
 65 70 75 80

Arg Ser Tyr Tyr Tyr Gly Lys Phe Met Pro Val Gly Leu Ile Ala Gly
85 90 95

5 Ala Ser Leu Leu Met Ala Ala Lys Val Gly Val Arg Met Leu Met Thr
100 105 110

Ser Asp

10

(2) INFORMATION FOR SEQ ID NO: 132:

15

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 132:

20

Met Ile Thr Leu Leu Ile Trp Met Leu Ala Gly Phe Ile Ala Arg Ile
1 5 10 15

Xaa Val Ala Leu Gln Xaa
20

25

(2) INFORMATION FOR SEQ ID NO: 133:

30

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 52 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 133:

35

Met Ala Gly Val Ser Glu Ile Ser Val Cys Phe Xaa Leu Leu Ser Leu
1 5 10 15

40

Phe Ser Leu Phe Cys Ser Phe Tyr Phe Pro Lys Gln Ala Thr Pro Lys
20 25 30

Arg Asp Leu Phe Val Gln Glu Ser Gly Lys Gly Lys Arg Asn Thr Glu
35 40 45

45

Ser Trp Glu Xaa
50

50

(2) INFORMATION FOR SEQ ID NO: 134:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 99 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 134:

55

Met Thr Ser Ala Leu Thr Gln Gly Leu Glu Arg Ile Pro Asp Gln Leu
1 5 10 15

60

Gly Tyr Leu Val Leu Ser Glu Gly Ala Val Leu Ala Ser Ser Gly Asp
 20 25 30

5 Leu Glu Asn Asp Glu Gln Ala Ala Ser Ala Ile Ser Glu Leu Val Ser
 35 40 45

Thr Ala Cys Gly Phe Arg Leu His Arg Gly Met Asn Val Pro Phe Lys
 50 55 60

10 Arg Leu Ser Val Val Phe Gly Glu His Thr Leu Val Thr Val Ser
 65 70 75 80

15 Gly Gln Arg Val Phe Val Val Lys Arg Gln Asn Arg Gly Arg Glu Pro
 85 90 95

Ile Asp Val

20

(2) INFORMATION FOR SEQ ID NO: 135:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 176 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 135:

30 Met Gly Ser Ala Ala Leu Glu Ile Leu Gly Leu Val Leu Cys Leu Val
 1 5 10 15

Gly Trp Gly Gly Leu Ile Leu Ala Cys Gly Leu Pro Met Trp Gln Val
 20 25 30

35 Thr Ala Phe Leu Asp His Asn Ile Val Thr Ala Gln Thr Thr Trp Lys
 35 40 45

40 Gly Leu Trp Met Ser Cys Val Val Gln Ser Thr Gly His Met Gln Cys
 50 55 60

Lys Val Tyr Asp Ser Val Leu Ala Leu Ser Thr Glu Val Gln Ala Ala
 65 70 75 80

45 Arg Ala Leu Thr Val Ser Ala Val Leu Leu Ala Phe Val Ala Leu Phe
 85 90 95

Val Thr Leu Ala Gly Ala Gln Cys Thr Thr Cys Val Ala Pro Gly Pro
 100 105 110

50 Ala Lys Ala Arg Val Ala Leu Thr Gly Gly Val Leu Tyr Leu Phe Cys
 115 120 125

55 Gly Leu Leu Ala Leu Val Pro Leu Cys Trp Phe Ala Asn Ile Val Val
 130 135 140

Arg Glu Phe Tyr Asp Pro Ser Val Pro Val Ser Gln Lys Tyr Glu Leu
 145 150 155 160

60 Gly Ala Xaa Cys Thr Ser Ala Gly Arg Pro Pro Arg Cys Ser Trp Xaa

165

170

175

5

(2) INFORMATION FOR SEQ ID NO: 136:

10

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 187 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 136:

15

Met Val Leu Leu Trp Val Val Thr Cys Pro Ala Thr Met Leu Thr Glu
 1 5 10 15

20

Pro Gln Asn Pro His Leu Ile Gly Phe Val Ala Tyr Ser Gly Pro Ser
 20 25 30

His Thr Thr Gln Pro His Lys Tyr Trp Leu Leu Leu Asp Gly Gln Ala
 35 40 45

25

Asp Pro Ala Ala Ala Glu Gly Pro Val Lys Arg Lys Ala Ala Ser Val
 50 55 60

Val Trp Trp Pro Gln Ala Leu Arg His Leu Ser Leu Leu Val His Cys
 65 70 75 80

30

Trp Glu Glu Ser Tyr Glu Met Asn Ile Gly Cys Gln Ser Leu Trp Ala
 85 90 95

35

Gly Gly Leu Ala Ser Ser Gly Asn Gly Trp Asp Leu Gly Val Ala Phe
 100 105 110

Arg Arg Asp Thr Cys Met Ser Ser Ser Ser Leu His Trp Lys Glu Phe
 115 120 125

40

Lys Tyr Ala Pro Gly Ser Leu His Tyr Phe Ala Leu Ser Phe Val Leu
 130 135 140

Ile Leu Thr Glu Ile Cys Leu Val Ser Ser Gly Met Gly Phe Pro Gln
 145 150 155 160

45

Glu Gly Lys His Phe Ser Val Leu Gly Ser Pro Asp Cys Ser Leu Trp
 165 170 175

50

Gly Arg Asp Glu His Val Pro Arg Glu Phe Ala
 180 185

(2) INFORMATION FOR SEQ ID NO: 137:

55

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 288 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 137:

276

Met Pro Ala His Arg Phe Val Leu Ala Val Gly Ser Ala Val Phe Asn
 1 5 10 15
 5 Ala Met Phe Asn Gly Gly Met Ala Thr Thr Ser Thr Glu Ile Glu Leu
 20 25 30
 Pro Asp Val Glu Pro Ala Ala Phe Leu Ala Leu Leu Lys Phe Leu Tyr
 35 40 45
 10 Ser Asp Glu Val Gln Ile Gly Pro Glu Thr Val Met Thr Thr Xaa Tyr
 50 55 60
 Thr Ala Lys Lys Tyr Ala Val Pro Ala Leu Glu Ala His Cys Val Glu
 65 70 75 80
 15 Phe Leu Lys Lys Asn Leu Arg Ala Asp Asn Ala Phe Met Leu Leu Thr
 85 90 95
 20 Gln Ala Arg Leu Phe Asp Glu Pro Gln Leu Ala Ser Leu Cys Leu Glu
 100 105 110
 Asn Ile Asp Lys Asn Thr Ala Asp Ala Ile Thr Ala Glu Gly Phe Thr
 115 120 125
 25 Asp Ile Asp Leu Asp Thr Leu Val Ala Val Leu Glu Arg Asp Thr Leu
 130 135 140
 Gly Ile Arg Glu Val Arg Leu Phe Asn Ala Val Val Arg Trp Ser Glu
 145 150 155 160
 30 Ala Glu Cys Gln Arg Gln Gln Leu Gln Val Thr Pro Glu Asn Arg Arg
 165 170 175
 35 Lys Val Leu Gly Lys Ala Leu Gly Leu Ile Arg Phe Pro Leu Met Thr
 180 185 190
 Ile Glu Glu Phe Ala Ala Gly Pro Ala Gln Ser Gly Ile Leu Val Asp
 195 200 205
 40 Arg Glu Val Val Ser Leu Phe Cys Thr Ser Pro Ser Thr Pro Ser His
 210 215 220
 Glu Trp Ser Ser Leu Thr Gly Pro Ala Ala Ala Cys Val Gly Arg Ser
 225 230 235 240
 45 Ala Ala Ser Thr Ala Ser Ser Arg Trp Arg Val Ala Gly Ala Thr Xaa
 245 250 255
 50 Gly Pro Val Thr Ala Ser Gly Ser Gln Ser Thr Ser Ala Ser Ser Trp
 260 265 270
 Trp Asp Leu Gly Cys Met Asp Pro Ser Thr Gly Pro Pro Thr Thr Lys
 275 280 285
 55
 60

(2) INFORMATION FOR SEQ ID NO: 138:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 114 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 138:

5
10 Met Pro Arg Cys Arg Trp Leu Ser Leu Ile Leu Leu Thr Ile Pro Leu
1 5 10 15
Ala Leu Val Ala Arg Lys Asp Pro Lys Lys Asn Glu Thr Gly Val Leu
20 25 30
15 Arg Lys Leu Lys Pro Val Asn Ala Phe Xaa Cys Gln Arg Gly Ser Ser
35 40 45
Val Xaa Gly Phe Ala Met Gln Glu Tyr Asn Lys Glu Ser Glu Asp Lys
50 55 60
20 Tyr Val Phe Leu Val Val Lys Thr Leu Gln Ala Gln Leu Gln Val Thr
65 70 75 80
25 Asn Leu Leu Glu Tyr Leu Ile Asp Val Glu Ile Ala Arg Ser Asp Cys
85 90 95
Arg Lys Pro Leu Ser Thr Asn Glu Ile Ala Pro Phe Lys Xaa Thr Pro
100 105 110
30 Ser Xaa

35 (2) INFORMATION FOR SEQ ID NO: 139:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 120 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 139:

40
45 Met Ser Pro His Pro Thr Ala Leu Leu Gly Leu Val Leu Cys Leu Ala
1 5 10 15
Gln Thr Ile His Thr Gln Glu Glu Asp Leu Pro Arg Pro Ser Ile Ser
20 25 30
50 Ala Glu Pro Gly Thr Val Ile Pro Leu Gly Ser His Val Thr Phe Val
35 40 45
Cys Arg Gly Pro Val Gly Val Gln Thr Phe Arg Leu Glu Arg Glu Ser
50 55 60
55 Arg Ser Thr Tyr Asn Asp Thr Glu Asp Val Ser Gln Ala Ser Pro Ser
65 70 75 80
Glu Ser Glu Ala Arg Phe Arg Ile Asp Ser Val Ser Glu Gly Asn Ala
85 90 95
60

Gly Pro Tyr Arg Cys Ile Tyr Tyr Lys Pro Pro Lys Trp Ser Glu Gln
 100 105 110

5 Ser Asp Tyr Trp Ser Cys Trp Xaa
 115 120

(2) INFORMATION FOR SEQ ID NO: 140:

10

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 438 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 140:

Met Asn Thr Pro Asn Gly Asn Ser Leu Ser Ala Ala Glu Leu Thr Cys
 1 5 10 15

20 Gly Met Ile Met Cys Leu Ala Arg Gln Ile Pro Gln Ala Thr Ala Ser
 20 25 30

Met Lys Asp Gly Lys Trp Glu Arg Lys Lys Phe Met Gly Thr Glu Leu
 35 40 45

25

Asn Gly Lys Thr Leu Gly Ile Leu Gly Leu Gly Arg Ile Gly Arg Glu
 50 55 60

30 Val Ala Thr Arg Met Gln Ser Phe Gly Met Lys Thr Ile Gly Tyr Asp
 65 70 75 80

Pro Ile Ile Ser Pro Glu Val Ser Ala Ser Phe Gly Val Gln Gln Leu
 85 90 95

35 Pro Leu Glu Glu Ile Trp Pro Leu Cys Asp Phe Ile Thr Val His Thr
 100 105 110

Pro Leu Leu Pro Ser Thr Thr Gly Leu Leu Asn Asp Asn Thr Phe Ala
 115 120 125

40

Gln Cys Lys Lys Gly Val Arg Val Val Asn Cys Ala Arg Gly Gly Ile
 130 135 140

45 Val Asp Glu Gly Ala Leu Leu Arg Ala Leu Gln Ser Gly Gln Cys Ala
 145 150 155 160

Gly Ala Ala Leu Asp Val Phe Thr Glu Glu Pro Pro Arg Asp Arg Ala
 165 170 175

50 Leu Val Asp His Glu Asn Val Ile Ser Cys Pro His Leu Gly Ala Ser
 180 185 190

Thr Lys Glu Ala Gln Ser Arg Cys Gly Glu Glu Ile Ala Val Gln Phe
 195 200 205

55

Val Asp Met Val Lys Gly Lys Ser Leu Thr Gly Val Val Asn Ala Gln
 210 215 220

60 Ala Leu Thr Ser Ala Phe Ser Pro His Thr Lys Pro Trp Ile Gly Leu
 225 230 235 240

279

Ala Glu Ala Leu Gly Thr Leu Met Arg Ala Trp Ala Gly Ser Pro Lys
 245 250 255

5 Gly Thr Ile Gln Val Ile Thr Gln Gly Thr Ser Leu Lys Asn Ala Gly
 260 265 270

Asn Cys Leu Ser Pro Ala Val Ile Val Gly Leu Leu Lys Glu Ala Ser
 275 280 285

10 Lys Gln Ala Asp Val Asn Leu Val Asn Ala Lys Leu Leu Val Lys Glu
 290 295 300

Ala Gly Leu Asn Val Thr Thr Ser His Ser Pro Ala Ala Pro Gly Glu
 15 305 310 315 320

Gln Gly Phe Gly Glu Cys Leu Leu Ala Val Ala Leu Ala Gly Ala Pro
 325 330 335

20 Tyr Gln Ala Val Gly Leu Val Gln Gly Thr Thr Pro Val Leu Gln Gly
 340 345 350

Leu Asn Gly Ala Val Phe Arg Pro Glu Val Pro Leu Arg Arg Asp Leu
 355 360 365

25 Pro Leu Leu Leu Phe Arg Thr Gln Thr Ser Asp Pro Ala Met Leu Pro
 370 375 380

Thr Met Ile Gly Leu Leu Ala Glu Ala Gly Val Arg Leu Leu Ser Tyr
 30 385 390 395 400

Gln Thr Ser Leu Val Ser Asp Gly Glu Thr Trp His Val Met Gly Ile
 405 410 415

35 Ser Ser Leu Leu Pro Ser Leu Glu Ala Trp Lys Gln His Val Thr Glu
 420 425 430

Ala Phe Gln Phe His Phe
 40 435

(2) INFORMATION FOR SEQ ID NO: 141:

- 45 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 164 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 141:

50 Met Ser Arg Pro Thr His Thr Pro Leu Ser Pro Ala Thr Ile Ser Pro
 1 5 10 15

Thr Ile Thr Val Ala Val Phe Phe Ala Val Phe Val Ala Ala Ala Ala
 55 20 25 30

Ala Thr Ala Val Val Ala Val Ala Ala Ala Thr Thr Ser Ser Gly Arg
 35 40 45

60 Arg Thr Xaa Asp Lys Ser Pro Ile Ala Thr Gln Ser Ser Val Thr His

280

50 55 60

Ile Ala Ala Lys Arg Cys His Asn Tyr Thr Glu Cys Leu Ser Leu Ile
65 70 75 80

5 Arg Xaa Thr Arg Ile Pro Thr Trp Xaa Xaa Xaa Thr Thr Cys Pro Ser
 85 90 95

10 Arg Ile Pro Ser Thr His Val Ala Ala Gly Ala Gly Phe Ile Arg Glu
 100 105 110

Arg Ala Cys Leu Gln Cys Gly Ala Val Gly Pro Pro Gly Cys Ile Leu
115 120 125

15 Ala Ser Leu Pro Pro Pro Ser Leu Tyr Leu Ser Pro Glu Leu Arg Cys
130 135 140

Met Pro Lys Arg Val Glu Ala Arg Ser Glu Leu Arg Leu Cys Pro Pro
145 150 155 160

20 Gly Val Xaa Xaa

25

(2) INFORMATION FOR SEQ ID NO: 142:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 73 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 142:

35 Met Gln Arg Trp Val Cys Ile Leu Glu Phe Lys Glu Asn Leu Phe Gln
1 5 10 15

Ile Pro Ser Ser Leu Val Ala Leu Leu Asn Thr Leu Phe Leu Asp Ile
20 25 30

40 Leu His Pro Gln Asn Ser Leu Ser Pro His Gly Ser Phe Ser Leu Ser
35 40 45

Ser Leu Ser Phe Pro Pro Leu Pro Val Ser Ser Leu Gln Pro Phe Leu
50 55 60

45 Phe Leu Arg Ser Leu Leu Cys Arg Xaa
65 70

50

(2) INFORMATION FOR SEQ ID NO: 143:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 123 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 143:

60 Phe Gly Thr Arg Phe Leu Ala Asn Leu Leu Leu Glu Glu Asp Asn Lys
1 5 10 15

Phe Cys Ala Asp Cys Gln Ser Lys Gly Pro Arg Trp Ala Ser Trp Asn
 20 25 30
 5 Ile Gly Val Phe Ile Cys Ile Arg Cys Ala Xaa Ile His Arg Asn Leu
 35 40 45
 Gly Val His Ile Ser Arg Val Lys Ser Val Asn Leu Asp Gln Trp Thr
 50 55 60
 10 Gln Val Gln Ile Gln Cys Met Gln Xaa Met Gly Asn Gly Lys Ala Asn
 65 70 75 80
 Arg Leu Tyr Glu Ala Tyr Leu Pro Glu Thr Phe Arg Arg Pro Gln Ile
 15 85 90 95
 Asp Pro Ala Val Glu Gly Phe Ile Arg Asp Xaa Tyr Glu Lys Lys Lys
 100 105 110
 20 Tyr Met Asp Arg Ser Leu Gly His Gln Cys Leu
 115 120

25 (2) INFORMATION FOR SEQ ID NO: 144:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 138 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 144:

Met Ser Leu Tyr Asp Asp Leu Gly Val Glu Thr Ser Asp Ser Lys Thr
 1 5 10 15
 35 Glu Gly Trp Ser Lys Asn Phe Lys Leu Leu Gln Ser Gln Leu Gln Val
 20 25 30
 40 Lys Lys Ala Ala Leu Thr Gln Ala Lys Ser Gln Arg Thr Lys Gln Ser
 35 40 45
 Thr Val Leu Ala Pro Val Ile Asp Leu Lys Arg Gly Gly Ser Ser Asp
 50 55 60
 45 Asp Arg Gln Ile Val Asp Thr Pro Pro His Val Ala Ala Gly Leu Lys
 65 70 75 80
 Asp Pro Val Pro Ser Gly Phe Ser Ala Gly Glu Val Leu Ile Pro Leu
 85 90 95
 50 Ala Asp Glu Tyr Asp Pro Met Phe Pro Asn Asp Tyr Glu Lys Val Val
 100 105 110
 55 Lys Arg Ala Lys Arg Gly Thr Thr Glu Thr Ala Gly Val Xaa Lys Thr
 115 120 125
 Lys Gly Asn Arg Arg Lys Gly Lys Lys Ala
 130 135

60

(2) INFORMATION FOR SEQ ID NO: 145:

(i) SEQUENCE CHARACTERISTICS:

5

(A) LENGTH: 356 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 145:

10 Met Leu Ala Arg Ala Ala Arg Gly Thr Gly Ala Leu Leu Leu Arg Gly
1 5 10 15

Ser Leu Leu Ala Ser Gly Arg Ala Pro Arg Arg Ala Ser Ser Gly Leu
20 25 30

15 Pro Arg Asn Thr Val Val Leu Phe Val Pro Gln Gln Glu Ala Trp Val
35 40 45

20 Val Glu Arg Met Gly Arg Phe His Arg Ile Leu Glu Pro Gly Leu Asn
50 55 60

Ile Leu Ile Pro Val Leu Asp Arg Ile Arg Tyr Val Gln Ser Leu Lys
65 70 75 80

25 Glu Ile Val Ile Asn Val Pro Glu Gln Ser Ala Val Thr Leu Asp Asn
85 90 95

Val Thr Leu Gln Ile Asp Gly Val Leu Tyr Leu Arg Ile Met Asp Pro
100 105 110

30 Tyr Lys Ala Ser Tyr Gly Val Glu Asp Pro Glu Tyr Ala Val Thr Gln
115 120 125

Leu Ala Gln Thr Thr Met Arg Ser Glu Leu Gly Lys Leu Ser Leu Asp
130 135 140

35 Lys Val Phe Arg Glu Arg Glu Ser Leu Asn Ala Ser Ile Val Asp Ala
145 150 155 160

40 Ile Asn Gln Ala Ala Asp Cys Trp Gly Ile Arg Cys Leu Arg Tyr Glu
165 170 175

Ile Lys Asp Ile His Val Pro Pro Arg Val Lys Glu Ser Met Gln Met
180 185 190

45 Gln Val Glu Ala Glu Arg Arg Lys Arg Ala Thr Val Leu Glu Ser Glu
195 200 205

Gly Thr Arg Glu Ser Ala Ile Asn Val Ala Glu Gly Lys Lys Gln Ala
210 215 220

50 Gln Ile Leu Ala Ser Glu Ala Glu Lys Ala Glu Gln Ile Asn Gln Ala
225 230 235 240

55 Ala Gly Glu Ala Ser Ala Val Leu Ala Lys Ala Lys Ala Lys Ala Glu
245 250 255

Ala Ile Arg Ile Leu Ala Ala Ala Leu Thr Gln His Asn Gly Asp Ala
260 265 270

60

283

Ala Ala Ser Leu Thr Val Ala Glu Gln Tyr Val Ser Ala Phe Ser Lys
 275 280 285

5 Leu Ala Lys Asp Ser Asn Thr Ile Leu Leu Pro Ser Asn Pro Gly Asp
 290 295 300

Val Thr Ser Met Val Ala Gln Ala Met Gly Val Tyr Gly Ala Leu Thr
 305 310 315 320

10 Lys Ala Pro Val Pro Gly Thr Pro Asp Ser Leu Ser Ser Gly Ser Ser
 325 330 335

Arg Asp Val Gln Gly Thr Asp Ala Ser Leu Asp Glu Glu Leu Asp Arg
 340 345 350

15 Val Lys Met Ser
 355

20

(2) INFORMATION FOR SEQ ID NO: 146:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 40 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 146:

30 Met Tyr Ile Leu Leu Phe Trp Gly Gly Xaa Phe His Arg Cys Leu Ser
 1 5 10 15

Xaa Leu Phe Asp Pro Glu Leu Xaa Ser Xaa Pro Gly Ile Ser Xaa Phe
 20 25 30

35 Thr Val Xaa Leu Gln Met Thr Xaa
 35 40

40

(2) INFORMATION FOR SEQ ID NO: 147:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 147:

50 Met Pro Ser Pro Lys Tyr Cys Met His Thr Asn Asp Val Gln Ser Val
 1 5 10 15

Glu Tyr Asn Gly Asp Thr Leu Phe Gln Lys Leu Ser Ser Ser Xaa Leu
 20 25 30

55 Ser Phe Lys Ser Ile His Ile Tyr Pro Asn Glu Xaa Lys Thr Cys Xaa
 35 40 45

Xaa Ile Phe Ile Ser Lys Val Tyr Met Ile Ser Lys Thr Trp Lys Xaa
 50 55 60

60 Pro Arg Phe Thr Ser Xaa Gly

65 70

5 (2) INFORMATION FOR SEQ ID NO: 148:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33 amino acids

(B) TYPE: amino acid

10 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 148:

Met Asn Phe Val Leu Phe Phe Ile Gly Ile Asn Val Gly Cys Arg Gly
1 5 10 15

15 Glu Asn Ser Leu Lys Tyr Phe Thr Val Thr Val Leu Cys Ser Pro Arg
 20 25 30

20 Asp

25 (2) INFORMATION FOR SEQ ID NO: 149:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 78 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 149:

Met Lys Glu Ala Gly Lys Gly Gly Val Ala Asp Ser Arg Glu Leu Lys
1 5 10 15

35 Pro Met Val Gly Gly Asp Glu Glu Val Ala Ala Leu Gln Glu Phe His
 20 25 30

Phe His Phe Leu Ser Leu Ser Val Phe Thr Asp Cys Thr Ser Ser Gly
35 40 45

Glu Ala Phe Val Ile Cys Ile Thr Gln Thr Cys Cys Ser Phe Cys Leu
50 55 60

45 Cys Ala Tyr Pro Ser Leu Gly Trp Gln Asn Ser Cys His Asn
65 70 75

50 (2) INFORMATION FOR SEQ ID NO: 150:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 150:

Met Phe Ser Ser Lys Ser Leu Leu Val Leu Pro Phe Cys Phe Arg Ser
1 5 10 15

60 Ala Ala His Leu Glu Leu Ser Val Trp Cys Val Cys Gly Val Arg Xaa

285

20

25

30

5

(2) INFORMATION FOR SEQ ID NO: 151:

10

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 464 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 151:

15

Met Leu Ala Leu Gly Asn Asn His Phe Ile Gly Phe Val Asn Asp Ser
 1 5 10 15

20

Val Thr Lys Ser Ile Val Ala Leu Arg Leu Thr Leu Val Val Lys Val
 20 25 30

Ser Thr Xaa Pro Gly Glu Ser His Ala Asn Asp Leu Glu Cys Ser Gly
 35 40 45

25

Lys Gly Lys Cys Thr Thr Lys Pro Ser Glu Ala Thr Phe Ser Cys Thr
 50 55 60

Cys Glu Glu Gln Tyr Val Gly Thr Phe Cys Glu Glu Tyr Asp Ala Cys
 65 70 75 80

30

Gln Arg Lys Pro Cys Gln Asn Asn Ala Ser Cys Ile Asp Ala Asn Glu
 85 90 95

35

Lys Gln Asp Gly Ser Asn Phe Thr Cys Val Cys Leu Pro Gly Tyr Thr
 100 105 110

Gly Glu Leu Cys Gln Ser Lys Ile Asp Tyr Cys Ile Leu Asp Pro Cys
 115 120 125

40

Arg Asn Gly Ala Thr Cys Ile Ser Ser Leu Ser Gly Phe Thr Cys Gln
 130 135 140

Cys Pro Glu Gly Tyr Phe Gly Ser Ala Cys Glu Glu Lys Val Asp Pro
 145 150 155 160

45

Cys Ala Ser Ser Pro Cys Gln Asn Asn Gly Thr Cys Tyr Val Asp Gly
 165 170 175

50

Val His Phe Thr Cys Asn Cys Ser Pro Gly Phe Thr Gly Pro Thr Cys
 180 185 190

Ala Gln Leu Ile Asp Phe Cys Ala Leu Ser Pro Cys Ala His Gly Thr
 195 200 205

55

Cys Arg Ser Val Gly Thr Ser Tyr Lys Cys Leu Cys Asp Pro Gly Tyr
 210 215 220

His Gly Leu Tyr Cys Glu Glu Glu Tyr Asn Glu Cys Leu Ser Ala Pro
 225 230 235 240

60

286

Cys Leu Asn Ala Ala Thr Cys Arg Asp Leu Val Asn Gly Tyr Glu Cys
 245 250 255
 5 Val Cys Leu Ala Glu Tyr Lys Gly Thr His Cys Glu Leu Tyr Lys Asp
 260 265 270
 Pro Cys Ala Asn Val Ser Cys Leu Asn Gly Ala Thr Cys Asp Ser Asp
 275 280 285
 10 Gly Leu Asn Gly Thr Cys Ile Cys Ala Pro Gly Phe Thr Gly Glu Glu
 290 295 300
 Cys Asp Ile Asp Ile Asn Glu Cys Asp Ser Asn Pro Cys His His Gly
 305 310 315 320
 15 Gly Ser Cys Leu Asp Gln Pro Asn Gly Tyr Asn Xaa His Cys Pro His
 325 330 335
 Gly Trp Val Gly Ala Asn Cys Glu Ile His Leu Gln Trp Lys Ser Gly
 340 345 350
 20 His Met Ala Glu Ser Leu Thr Asn Met Pro Arg His Ser Leu Tyr Ile
 355 360 365
 25 Ile Ile Gly Ala Leu Cys Val Ala Phe Ile Leu Met Leu Ile Ile Leu
 370 375 380
 Ile Val Gly Ile Cys Arg Ile Ser Arg Ile Glu Tyr Gln Gly Ser Ser
 385 390 395 400
 30 Arg Pro Ala Tyr Xaa Glu Phe Tyr Asn Cys Arg Ser Ile Asp Ser Glu
 405 410 415
 Phe Ser Asn Ala Ile Ala Ser Ile Arg His Ala Arg Phe Gly Lys Lys
 420 425 430
 35 Ser Arg Pro Ala Met Tyr Asp Val Ser Pro Ile Ala Tyr Glu Asp Tyr
 435 440 445
 40 Ser Pro Asp Asp Lys Pro Leu Val Thr Leu Ile Lys Thr Lys Asp Leu
 450 455 460

45

(2) INFORMATION FOR SEQ ID NO: 152:

50

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 151 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 152:

55

Met His His Gln Met Thr Arg Thr Thr Leu Met Thr Lys Gln His Glu
 1 5 10 15

60

Leu Gly Gly Leu Leu Ala Leu Val Gln Asn Cys Gln Ser Glu Met Asn
 20 25 30

Ile Lys Asp Ser Arg Ala Val Gly Leu Ser Val Lys Arg Leu Cys Ile
 35 40 45

5 Ser Phe Val Asp Glu Phe Cys Glu Arg Thr Glu Arg Pro Leu Tyr Leu
 50 55 60

Ala Gln Gly Leu Phe Met Lys Arg Glu Thr Tyr Trp Glu Val Gln Asp
 65 70 75 80

10 Ser Gly Ile Ser Pro Leu Leu Leu Leu Ser Thr Ala Leu Asp Cys
 85 90 95

Ser Pro Glu Ala Glu Thr Arg Gln Ser Pro Gly Gly Arg Lys Met Leu
 100 105 110

Gln Glu Pro Thr Leu Ser Met Ser Leu Gln Ile Leu Thr Gly Phe Leu
 115 120 125

20 Trp Val Gln Leu Trp Asn Trp Glu Thr Phe Leu Arg Ile Arg Thr His
 130 135 140

Ser Thr Asp Ala Ser Cys Pro
 145 150

25

(2) INFORMATION FOR SEQ ID NO: 153:

30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 299 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 153:

35 Met Ala Gln Asn Leu Lys Asp Leu Ala Gly Arg Leu Pro Ala Gly Pro
 1 5 10 15

Arg Gly Met Gly Thr Ala Leu Lys Leu Leu Leu Gly Ala Gly Ala Val
 20 25 30

Ala Tyr Gly Val Arg Glu Ser Val Phe Thr Val Glu Gly Gly His Arg
 35 40 45

45 Ala Ile Phe Phe Asn Arg Ile Gly Gly Val Gln Gln Asp Thr Ile Leu
 50 55 60

Ala Glu Gly Leu His Phe Arg Ile Pro Trp Phe Gln Tyr Pro Ile Ile
 65 70 75 80

50 Tyr Asp Ile Arg Ala Arg Pro Arg Lys Ile Ser Ser Pro Thr Gly Ser
 85 90 95

Lys Asp Leu Gln Met Val Asn Ile Ser Leu Arg Val Leu Ser Arg Pro
 100 105 110

Asn Ala Gln Glu Leu Pro Ser Met Tyr Gln Arg Leu Gly Leu Asp Tyr
 115 120 125

60 Glu Glu Arg Val Leu Pro Ser Ile Val Asn Glu Val Leu Lys Ser Val

288

35 (2) INFORMATION FOR SEQ ID NO: 154:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 398 amino acids

(B) TYPE: amino acid

40 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 154:

Met Leu Arg Gly Pro Trp Arg Gln Leu Trp Leu Phe Xaa Leu Leu Leu
1 5 10 15

45 Leu Pro Gly Ala Pro Glu Pro Arg Gly Ala Ser Arg Pro Trp Glu Gly
20 25 30

50 Thr Asp Glu Pro Gly Ser Ala Trp Ala Trp Pro Gly Phe Gln Arg Leu
35 40 45

Gln Glu Gln Leu Arg Ala Ala Gly Ala Leu Ser Lys Arg Tyr Trp Thr
50 55 60

55 Leu Phe Ser Cys Gln Val Trp Pro Asp Asp Cys Asp Glu Asp Glu Glu
65 70 75 80

Ala Ala Thr Gly Pro Leu Gly Trp Arg Leu Pro Leu Leu Gly Gln Arg
85 90 95

60

289

Tyr Leu Asp Leu Leu Thr Thr Trp Tyr Cys Ser Phe Lys Asp Cys Cys
 100 105 110

5 Pro Arg Gly Asp Cys Arg Ile Ser Asn Asn Phe Thr Gly Leu Glu Trp
 115 120 125

Asp Leu Asn Val Arg Leu His Gly Gln His Leu Val Gln Gln Leu Val
 130 135 140

10 Leu Arg Thr Val Arg Gly Tyr Leu Glu Thr Pro Gln Pro Glu Lys Ala
 145 150 155 160

Leu Ala Leu Ser Phe His Gly Trp Ser Gly Thr Gly Lys Asn Phe Val
 165 170 175

15 Ala Arg Met Leu Val Glu Asn Leu Tyr Arg Asp Gly Leu Met Ser Asp
 180 185 190

20 Cys Val Arg Met Phe Ile Ala Thr Phe His Phe Pro His Pro Lys Tyr
 195 200 205

Val Asp Leu Tyr Lys Glu Gln Leu Met Ser Gln Ile Arg Glu Thr Gln
 210 215 220

25 Gln Leu Cys His Gln Thr Leu Phe Ile Phe Asp Glu Ala Glu Lys Leu
 225 230 235 240

His Pro Gly Leu Leu Glu Val Leu Gly Pro His Leu Glu Arg Arg Ala
 245 250 255

30 Pro Xaa Gly His Arg Ala Glu Ser Pro Trp Thr Ile Phe Leu Phe Leu
 260 265 270

35 Ser Asn Leu Arg Gly Asp Ile Ile Asn Glu Val Val Leu Lys Leu Leu
 275 280 285

Lys Ala Gly Trp Ser Arg Glu Glu Ile Thr Met Glu His Leu Glu Pro
 290 295 300

40 His Leu Gln Ala Glu Ile Val Glu Thr Ile Asp Asn Gly Phe Gly His
 305 310 315 320

Ser Arg Leu Val Lys Glu Asn Leu Ile Asp Tyr Phe Ile Pro Phe Leu
 325 330 335

45 Pro Leu Glu Tyr Arg His Val Arg Leu Cys Ala Arg Asp Ala Phe Leu
 340 345 350

50 Ser Gln Glu Leu Leu Tyr Lys Glu Glu Thr Leu Asp Glu Ile Ala Gln
 355 360 365

Met Met Val Tyr Val Pro Lys Glu Glu Gln Leu Phe Ser Ser Gln Gly
 370 375 380

55 Cys Lys Ser Ile Ser Gln Arg Ile Asn Tyr Phe Leu Ser Xaa
 385 390 395

60 (2) INFORMATION FOR SEQ ID NO: 155:

290

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 83 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 155:

Met Ala Phe Thr Leu Tyr Ser Leu Leu Gln Ala Xaa Leu Leu Cys Val
 1 5 10 15

Asn Ala Ile Ala Val Leu His Glu Glu Arg Phe Leu Lys Asn Ile Gly
 20 25 30

Trp Gly Thr Asp Gln Gly Ile Gly Gly Phe Gly Glu Glu Pro Gly Ile
 35 40 45

Lys Ser Gln Leu Met Asn Leu Ile Arg Ser Val Arg Thr Val Met Arg
 50 55 60

Val Pro Leu Ile Ile Val Asn Ser Ile Ala Ile Val Leu Leu Leu Leu
 65 70 75 80

Phe Gly Xaa

(2) INFORMATION FOR SEQ ID NO: 156:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 50 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 156:

Met Ala Pro Arg Asn Gln Gly Ser Phe Ser Phe Gly Asn Phe Met Leu
 1 5 10 15

Phe Leu Val Leu Ile Glu Arg Arg Tyr Leu Pro Phe Leu Ser Pro Ile
 20 25 30

Leu Phe Cys Cys Ser Thr His Asn Arg Ser Ala Val Thr Ala Thr Asn
 35 40 45

Leu Xaa
 50

(2) INFORMATION FOR SEQ ID NO: 157:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 51 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 157:

Met Asp Val Leu Thr Val Ala Phe Leu Ser Ile Leu Ile Thr Ala Pro
 1 5 10 15

Ile Gly Ser Leu Leu Ile Gly Leu Leu Gly Pro Arg Leu Leu Gln Lys
 20 25 30
 5 Val Glu His Gln Asn Lys Asp Glu Glu Val Gln Gly Glu Thr Ser Val
 35 40 45
 Gln Val Xaa
 50

10

(2) INFORMATION FOR SEQ ID NO: 158:

15 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 17 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 158:

20 Pro Asn Ser Phe Ser Cys Leu Gly Leu Ala Gly Thr Gly Ala Gly Ile
 1 5 10 15
 Xaa

25

(2) INFORMATION FOR SEQ ID NO: 159:

30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 53 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 159:

35 Met Gly Arg Tyr His Phe Val Phe Leu Thr Phe Phe Phe Ser Thr Tyr
 1 5 10 15

40 Ser Ser Cys Phe Tyr Pro Val Val Ser Gln Val Leu Tyr Leu Val Cys
 20 25 30

Ser Cys Thr Ala Asp Arg Pro Leu Met Ala Pro Val Gly Ser Cys Leu
 35 40 45

45 Gly Gly Arg Asn Xaa
 50

50 (2) INFORMATION FOR SEQ ID NO: 160:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 64 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 160:

60 Met Phe Val Thr Leu Ser Ile Leu Asn Ile Thr Ile Glu Lys Asp Lys
 1 5 10 15

Ser Thr Asn Arg Phe Arg Asp Val Phe Leu Gln His Ile Leu Val Ile
 20 25 30
 5 Leu Met Pro Ser Leu Thr Tyr Cys Leu Ile Gly Gln His Leu Cys Ser
 35 40 45
 Phe Thr Arg Tyr Val Ser Leu Cys Tyr Ser Arg Cys His Ser Trp Xaa
 50 55 60

10

15 (2) INFORMATION FOR SEQ ID NO: 161:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 161:

Met Ser Ile Cys Pro Leu Leu Val Met Leu Ile Leu Ile Thr Trp Val
 1 5 10 15

25

Arg Cys Pro Val Ser Pro Val Tyr Arg Tyr Cys Phe Ser Phe Cys Asn
 20 25 30

30

Xaa

35 (2) INFORMATION FOR SEQ ID NO: 162:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 95 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 162:

Met Gln Asp Ile Val Tyr Lys Leu Val Pro Gly Leu Gln Glu Gly Glu
 1 5 10 15

45

Cys Leu Thr Val Leu Leu Ile Pro Glu Val Pro Ala Trp Pro Leu Gln
 20 25 30

50

Pro Leu Leu Ser Trp Lys Phe Gly Ser Arg Met Gly Gly Pro Phe Pro
 35 40 45

Phe Gly Arg Ile Thr Val Phe Ser Ser Leu Leu Ser Ala Gln Leu His
 50 55 60

55

Leu Leu Gly Trp Ser Leu Leu Ser Ser Lys Met Arg Xaa His Leu Phe
 65 70 75 80

60

Thr Pro Tyr Val Tyr Ser Phe Ser Lys Tyr Gly Ser His Val Xaa
 85 90 95

(2) INFORMATION FOR SEQ ID NO: 163:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 58 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 163:

5
10 Met Lys Val Leu Ala Thr Ser Phe Val Leu Gly Ser Leu Gly Leu Ala
1 5 10 15
Phe Tyr Leu Pro Leu Val Val Thr Thr Pro Lys Thr Leu Ala Ile Pro
20 25 30
15 Xaa Glu Ala Ala Arg Ser Cys Gly Glu Ser Tyr His Gln Cys His Asn
35 40 45
20 Leu Tyr Cys His Leu Trp Pro Trp Leu Xaa
50 55

(2) INFORMATION FOR SEQ ID NO: 164:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 44 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 164:

25
30 Met Asp Tyr Gly Tyr Tyr Ser Ala Gly Gln Phe Leu Leu His Leu Phe
1 5 10 15
35 Leu Ala Asp Leu Thr Gln Ala Thr Thr Gln Gln Lys Thr Asn Thr Ser
20 25 30
Glu Asn Gly Cys Lys Phe Val Cys Ala Val Phe Xaa
35 40
40

(2) INFORMATION FOR SEQ ID NO: 165:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 18 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 165:

50
Gly Ile Val Leu Leu Ile Gly Val Leu Val Gln Val Ser Ala Val Asp
1 5 10 15
55 Asp Xaa

(2) INFORMATION FOR SEQ ID NO: 166:

60

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 166:

Met Gly Asn Ala Phe Glu Val Thr Gly Leu Met Leu Ala Leu Leu Cys
1 5 10 15

10 Tyr Val Val Asp Gly Gln Lys Pro Lys Xaa Gly Phe Xaa Xaa
20 25 30

15 (2) INFORMATION FOR SEQ ID NO: 167:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 37 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 167:

Met Ser His Glu Lys Ser Asn Glu Leu Val Leu Leu Ile Val Thr Val
1 5 10 15

25 Met Arg Ser Leu Thr Tyr Asn Ile Ala Val Val Ala Ala Trp Phe Asn
20 25 30

30 Gly Cys Ile Arg Xaa
35

35 (2) INFORMATION FOR SEQ ID NO: 168:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 40 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 168:

Met Tyr Leu Leu Tyr Leu Pro Ser Ala Leu Leu Pro Pro Tyr Pro Thr
1 5 10 15

45 Cys Pro Tyr Glu His Gly Ser Pro Trp Pro His Thr Pro Ala Lys Leu
20 25 30

Leu Cys Cys Phe Ala Phe Leu Xaa
35 40

50

(2) INFORMATION FOR SEQ ID NO: 169:

55 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 47 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

60 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 169:

Met Lys Phe Ile Val Trp Arg Arg Phe Lys Trp Val Ile Ile Gly Leu
 1 5 10 15
 5 Leu Phe Leu Leu Ile Leu Leu Leu Phe Val Ala Val Leu Leu Tyr Ser
 20 25 30
 Leu Pro Asn Tyr Leu Ser Met Lys Ile Val Lys Pro Asn Val Xaa
 35 40 45

10

(2) INFORMATION FOR SEQ ID NO: 170:

(i) SEQUENCE CHARACTERISTICS:
 15 (A) LENGTH: 34 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 170:

20 Ile Glu Trp Ser Gly Tyr Asn Lys Pro Glu Arg Lys Gly Pro Leu Ala
 1 5 10 15
 Leu Phe Leu Val Phe Leu Phe Leu Asp Thr Pro Pro Leu Gln Gly Asp
 20 25 30
 25 Leu Xaa

30

(2) INFORMATION FOR SEQ ID NO: 171:

(i) SEQUENCE CHARACTERISTICS:
 35 (A) LENGTH: 5 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 171:

40 Met Ser Leu Leu Xaa
 1 5

45 (2) INFORMATION FOR SEQ ID NO: 172:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 25 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 172:

Met Gln Leu Leu Ile Val Trp Asn Glu Ser Leu Thr Asn Ser Val Pro
 1 5 10 15
 55 Ala Ser Val Asp Thr Ser Gln Cys Xaa
 20 25

60 (2) INFORMATION FOR SEQ ID NO: 173:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 262 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 173:

5 Met Ala Leu Gly Leu Lys Cys Phe Arg Met Val His Pro Thr Phe Arg
 1 5 10 15
 10 Asn Tyr Leu Ala Ala Ser Ile Arg Pro Val Ser Glu Val Thr Leu Lys
 20 25 30
 15 Thr Val His Glu Arg Gln His Gly His Arg Gln Tyr Met Ala Tyr Ser
 35 40 45
 Ala Val Pro Val Arg His Phe Ala Thr Lys Lys Ala Lys Ala Lys Gly
 50 55 60
 20 Lys Gly Gln Ser Gln Thr Arg Val Asn Ile Asn Ala Ala Leu Val Glu
 65 70 75 80
 Asp Ile Ile Asn Leu Glu Glu Val Asn Glu Glu Met Lys Ser Val Ile
 85 90 95
 25 Glu Ala Leu Lys Asp Asn Phe Asn Lys Thr Leu Asn Ile Arg Thr Ser
 100 105 110
 30 Pro Gly Ser Leu Asp Lys Ile Ala Val Val Thr Ala Asp Gly Lys Leu
 115 120 125
 Ala Leu Asn Gln Ile Ser Gln Ile Ser Met Lys Ser Pro Gln Leu Ile
 130 135 140
 35 Leu Val Asn Met Ala Ser Phe Pro Glu Cys Thr Ala Ala Ala Ile Lys
 145 150 155 160
 Ala Ile Arg Glu Ser Gly Met Asn Leu Asn Pro Glu Val Glu Gly Thr
 165 170 175
 40 Leu Ile Arg Val Pro Ile Pro Gln Val Thr Arg Glu His Arg Glu Met
 180 185 190
 45 Leu Val Lys Leu Ala Lys Gln Asn Thr Asn Lys Ala Lys Asp Ser Leu
 195 200 205
 Arg Lys Val Arg Thr Asn Ser Met Asn Lys Leu Lys Lys Ser Lys Asp
 210 215 220
 50 Thr Val Ser Glu Asp Thr Ile Arg Leu Ile Glu Lys Gln Ile Ser Gln
 225 230 235 240
 Met Ala Asp Asp Thr Val Ala Glu Leu Asp Arg His Leu Ala Val Lys
 245 250 255
 55 Thr Lys Glu Leu Leu Gly
 260
 60

(2) INFORMATION FOR SEQ ID NO: 174:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 967 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 174:

5 Met Gln Arg Ala Val Pro Glu Gly Phe Gly Arg Arg Lys Leu Gly Ser
 10 1 5 10 15
 Asp Met Gly Asn Ala Glu Arg Ala Pro Gly Ser Arg Ser Phe Gly Pro
 20 25 30
 15 Val Pro Thr Leu Leu Leu Leu Xaa Ala Ala Leu Leu Xaa Val Ser Asp
 35 40 45
 Ala Leu Gly Arg Pro Ser Glu Glu Asp Glu Glu Leu Val Val Pro Glu
 50 55 60
 20 Leu Glu Arg Ala Pro Gly His Gly Thr Thr Arg Leu Arg Leu His Ala
 65 70 75 80
 Phe Asp Gln Gln Leu Asp Leu Glu Leu Arg Pro Asp Ser Ser Phe Leu
 25 85 90 95
 Ala Pro Gly Phe Thr Leu Gln Asn Val Gly Arg Lys Ser Gly Ser Glu
 100 105 110
 30 Thr Pro Leu Pro Glu Thr Asp Leu Ala His Cys Phe Tyr Ser Gly Thr
 115 120 125
 Val Asn Gly Asp Pro Ser Ser Ala Ala Ala Leu Ser Leu Cys Glu Gly
 130 135 140
 35 Val Arg Gly Ala Phe Tyr Leu Leu Gly Glu Ala Tyr Phe Ile Gln Pro
 145 150 155 160
 Leu Pro Ala Ala Ser Glu Arg Leu Xaa Thr Ala Ala Pro Gly Glu Lys
 40 165 170 175
 Pro Pro Ala Pro Leu Gln Phe His Leu Leu Arg Arg Asn Arg Gln Gly
 180 185 190
 45 Asp Val Gly Gly Thr Cys Gly Val Val Asp Asp Glu Pro Arg Pro Thr
 195 200 205
 Gly Lys Ala Glu Thr Glu Asp Glu Asp Glu Gly Thr Glu Gly Glu Asp
 210 215 220
 50 Glu Gly Pro Gln Trp Ser Pro Gln Asp Pro Ala Leu Gln Gly Val Gly
 225 230 235 240
 Gln Pro Thr Gly Thr Gly Ser Ile Arg Lys Lys Arg Phe Val Ser Ser
 245 250 255
 55 His Arg Tyr Val Glu Thr Met Leu Val Ala Asp Gln Ser Met Ala Glu
 260 265 270
 60 Phe His Gly Ser Gly Leu Lys His Tyr Leu Leu Thr Leu Phe Ser Val

	275	280	285
	Ala Ala Arg Leu Xaa Lys His Pro Xaa Ile Arg Asn Ser Val Ser Leu		
	290	295	300
5	Val Val Val Lys Ile Leu Val Ile His Asp Glu Gln Lys Gly Pro Glu		
	305	310	315 320
	Val Thr Ser Asn Ala Ala Leu Thr Leu Arg Asn Phe Cys Asn Trp Gln		
10	325	330	335
	Lys Gln His Asn Pro Pro Ser Asp Arg Asp Ala Glu His Tyr Asp Thr		
	340	345	350
15	Ala Ile Leu Phe Thr Arg Gln Asp Leu Cys Gly Ser Gln Thr Cys Asp		
	355	360	365
	Thr Leu Gly Met Ala Asp Val Gly Thr Val Cys Asp Pro Ser Arg Ser		
20	370	375	380
	Cys Ser Val Ile Glu Asp Asp Gly Leu Gln Ala Ala Phe Thr Thr Ala		
	385	390	395 400
	His Glu Leu Gly His Val Phe Asn Met Pro His Asp Asp Ala Lys Gln		
25	405	410	415
	Cys Ala Ser Leu Asn Gly Val Asn Gln Asp Ser His Met Met Ala Ser		
	420	425	430
30	Met Leu Ser Asn Leu Asp His Ser Gln Pro Trp Ser Pro Cys Ser Ala		
	435	440	445
	Tyr Met Ile Thr Ser Phe Leu Asp Asn Gly His Gly Glu Cys Leu Met		
35	450	455	460
	Asp Lys Pro Gln Asn Pro Ile Gln Leu Pro Gly Asp Leu Pro Gly Thr		
	465	470	475 480
	Ser Tyr Asp Ala Asn Arg Gln Cys Gln Phe Thr Phe Gly Glu Asp Ser		
40	485	490	495
	Lys His Cys Pro Asp Ala Ala Ser Thr Cys Ser Thr Leu Trp Cys Thr		
	500	505	510
45	Gly Thr Ser Gly Gly Val Leu Val Cys Gln Thr Lys His Phe Pro Trp		
	515	520	525
	Ala Asp Gly Thr Ser Cys Gly Glu Gly Lys Trp Cys Ile Asn Gly Lys		
50	530	535	540
	Cys Val Xaa Lys Thr Asp Arg Lys His Phe Asp Thr Pro Phe His Gly		
	545	550	555 560
	Ser Trp Gly Met Trp Gly Pro Trp Gly Asp Cys Ser Arg Thr Cys Gly		
55	565	570	575
	Gly Gly Val Gln Tyr Thr Met Arg Glu Cys Asp Asn Pro Val Pro Lys		
	580	585	590
60	Asn Gly Gly Lys Tyr Cys Glu Gly Lys Arg Val Arg Tyr Arg Ser Cys		

	595	600	605
	Asn Leu Glu Asp Cys Pro Asp Asn Asn Gly Lys Thr Phe Arg Glu Glu		
	610	615	620
5	Gln Cys Glu Ala His Asn Glu Phe Ser Lys Ala Ser Phe Gly Ser Gly		
	625	630	635 640
10	Pro Ala Val Glu Trp Ile Pro Lys Tyr Ala Gly Val Ser Pro Lys Asp		
	645	650	655
	Arg Cys Lys Leu Ile Cys Gln Ala Lys Gly Ile Gly Tyr Phe Phe Val		
	660	665	670
15	Leu Gln Pro Lys Val Val Asp Gly Thr Pro Cys Ser Pro Asp Ser Thr		
	675	680	685
	Ser Val Cys Val Gln Gly Gln Cys Val Lys Ala Gly Cys Asp Arg Ile		
20	690	695	700
	Ile Asp Ser Lys Lys Lys Phe Asp Lys Cys Gly Val Cys Gly Gly Asn		
	705	710	715 720
25	Gly Ser Thr Cys Lys Lys Ile Ser Gly Ser Val Thr Ser Ala Lys Pro		
	725	730	735
	Gly Tyr His Asp Ile Ile Thr Ile Pro Thr Gly Ala Thr Asn Ile Glu		
	740	745	750
30	Val Lys Gln Arg Asn Gln Arg Gly Ser Arg Asn Asn Gly Ser Phe Leu		
	755	760	765
	Ala Ile Lys Ala Ala Asp Gly Thr Tyr Ile Leu Asn Gly Asp Tyr Thr		
35	770	775	780
	Leu Ser Thr Leu Glu Gln Asp Ile Met Tyr Lys Gly Val Val Leu Arg		
	785	790	795 800
40	Tyr Ser Gly Ser Ser Ala Ala Leu Glu Arg Ile Arg Ser Phe Ser Pro		
	805	810	815
	Leu Lys Glu Pro Leu Thr Ile Gln Val Leu Thr Val Gly Asn Ala Leu		
	820	825	830
45	Arg Pro Lys Ile Lys Tyr Thr Tyr Phe Val Lys Lys Lys Lys Glu Ser		
	835	840	845
	Phe Asn Ala Ile Pro Thr Phe Ser Ala Trp Val Ile Glu Glu Trp Gly		
50	850	855	860
	Glu Cys Ser Lys Ser Cys Glu Leu Gly Trp Gln Arg Arg Leu Val Glu		
	865	870	875 880
55	Cys Arg Asp Ile Asn Gly Gln Pro Ala Ser Glu Cys Ala Lys Glu Val		
	885	890	895
	Lys Pro Ala Ser Thr Arg Pro Cys Ala Asp His Pro Cys Pro Gln Trp		
	900	905	910
60	Gln Leu Gly Glu Trp Ser Ser Cys Ser Lys Thr Cys Gly Lys Gly Tyr		

300

915 920 925
 Lys Lys Arg Ser Leu Lys Cys Leu Ser His Asp Gly Gly Val Leu Ser
 930 935 940
 5 His Glu Ser Cys Asp Pro Leu Lys Lys Pro Lys His Phe Ile Asp Phe
 945 950 955 960
 10 Cys Thr Met Ala Glu Cys Ser
 965

15 (2) INFORMATION FOR SEQ ID NO: 175:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 175:

Met Leu Lys Ile Pro Thr His Leu Glu Gly Lys Ile Lys Ile Thr Lys
 1 5 10 15
 25 Val Tyr Xaa

30 (2) INFORMATION FOR SEQ ID NO: 176:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 205 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 176:

Met Tyr Glu Thr Met Lys Leu Asp Ala Cys Xaa His Gln Gln Arg Pro
 1 5 10 15
 40 Thr Leu Gln Ala Gly Pro Lys Leu Leu Thr Leu Ala Pro Arg Glu Glu
 20 25 30
 Pro Arg Gly Gln Ser Gly Arg Gly Ser Glu Leu Thr Ala Arg Gln Arg
 45 35 40 45
 His Ser Thr Gly Asp Pro Gln Gly Glu Gln Ala Leu Pro Arg Ala Gly
 50 55 60
 50 Cys Val Thr Gly Pro Pro Ala Thr Pro His Arg Pro Ser Glu Pro Gln
 65 70 75 80
 Leu Leu Arg Thr His Pro Asp Ala Arg Pro Lys Ser Ala Met Ala Gln
 85 90 95
 55 Thr Phe Val His Gln Gly Pro Val Ala Leu Gln Gln Leu Thr Thr Asn
 100 105 110
 Arg Arg Val Glu Thr Ser Met Ser Ser Asp Gly His Gly Gln Asn Pro
 60 115 120 125

Thr Pro Ser Pro Trp Ala Asp Val Cys Ala Ser Arg Ala Asp Ala Val
 130 135 140
 5 Ala Phe Pro Ala Ser Gly Xaa Cys His Ser Pro Trp Leu Met Xaa Pro
 145 150 155 160
 Ser Ser His Pro Leu Asn Pro His Ser Pro Leu Asn Leu Pro Pro Pro
 165 170 175
 10 Ser Phe His Cys Lys Asp Pro Val Met Thr Leu His Pro Gln Thr Leu
 180 185 190
 Val Thr Gln Gly His Leu Ser Thr Ser Gly Arg Leu Thr
 15 195 200 205

20 (2) INFORMATION FOR SEQ ID NO: 177:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 54 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 177:

Met Asp Ser Met Pro Glu Pro Ala Ser Arg Cys Leu Leu Leu Leu Pro
 1 5 10 15
 30 Leu Leu Leu Leu Leu Leu Leu Leu Pro Ala Pro Glu Leu Gly Pro
 20 25 30
 Ser Gln Ala Gly Ala Glu Glu Asn Asp Trp Val Arg Leu Pro Ser Lys
 35 35 40 45
 Cys Glu Gly Thr Cys Gly
 50

40 (2) INFORMATION FOR SEQ ID NO: 178:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 436 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 178:

Met Pro Leu Phe Leu Leu Ser Leu Pro Thr Pro Pro Ser Ala Ser Gly
 50 1 5 10 15
 His Glu Arg Arg Gln Arg Pro Glu Ala Lys Thr Ser Gly Ser Glu Lys
 20 25 30
 55 Lys Tyr Leu Arg Ala Met Gln Ala Asn Arg Ser Gln Leu His Ser Pro
 35 40 45
 Pro Gly Thr Gly Ser Ser Glu Asp Ala Ser Thr Pro Gln Cys Val His
 50 55 60
 60

Thr Arg Leu Thr Gly Glu Gly Ser Cys Pro His Ser Gly Asp Val His
 65 70 75 80
 5 Ile Gln Ile Asn Ser Ile Pro Lys Glu Cys Ala Glu Asn Ala Ser Ser
 85 90 95
 Arg Asn Ile Arg Ser Gly Val His Ser Cys Ala His Gly Cys Val His
 100 105 110
 10 Ser Arg Leu Arg Gly His Ser His Ser Glu Ala Arg Leu Thr Asp Asp
 115 120 125
 Thr Ala Ala Glu Ser Gly Asp His Gly Ser Ser Ser Phe Ser Glu Phe
 130 135 140
 15 Arg Tyr Leu Phe Lys Trp Leu Gln Lys Ser Leu Pro Tyr Ile Leu Ile
 145 150 155 160
 Leu Ser Val Lys Leu Val Met Gln His Ile Thr Gly Ile Ser Leu Gly
 165 170 175
 20 Ile Gly Leu Leu Thr Thr Phe Met Tyr Ala Asn Lys Ser Ile Val Asn
 180 185 190
 25 Gln Val Phe Leu Arg Glu Arg Ser Ser Lys Ile Gln Cys Ala Trp Leu
 195 200 205
 Leu Val Phe Leu Ala Gly Ser Ser Val Leu Leu Tyr Tyr Thr Phe His
 210 215 220
 30 Ser Gln Ser Leu Tyr Tyr Ser Leu Ile Phe Leu Asn Pro Thr Leu Asp
 225 230 235 240
 His Leu Ser Phe Trp Glu Val Phe Xaa Ile Val Gly Xaa Thr Asp Phe
 245 250 255
 35 Ile Leu Lys Phe Phe Phe Met Gly Leu Lys Cys Leu Ile Leu Leu Val
 260 265 270
 40 Pro Ser Phe Ile Met Pro Phe Lys Ser Lys Gly Tyr Trp Tyr Met Leu
 275 280 285
 Leu Glu Glu Leu Cys Gln Tyr Tyr Arg Thr Phe Val Pro Ile Pro Val
 290 295 300
 45 Trp Phe Arg Tyr Leu Ile Ser Tyr Gly Glu Phe Gly Xaa Val Thr Arg
 305 310 315 320
 Trp Xaa Leu Gly Ile Leu Leu Ala Leu Leu Tyr Leu Ile Leu Lys Leu
 325 330 335
 50 Leu Glu Phe Phe Gly His Leu Arg Thr Phe Arg Gln Val Leu Arg Ile
 340 345 350
 55 Phe Phe Thr Xaa Pro Ser Tyr Gly Val Ala Ala Ser Lys Arg Gln Cys
 355 360 365
 Ser Asp Val Asp Asp Ile Cys Ser Ile Cys Gln Ala Glu Phe Gln Lys
 370 375 380
 60

303

Pro Ile Leu Leu Ile Cys Gln His Ile Phe Cys Glu Glu Cys Met Thr
 385 390 395 400

5 Leu Trp Phe Asn Arg Glu Lys Thr Cys Pro Leu Cys Arg Thr Val Ile
 405 410 415

Ser Asp His Ile Asn Lys Trp Lys Asp Gly Ala Thr Ser Ser His Leu
 420 425 430

10 Gln Ile Tyr Xaa
 435

15 (2) INFORMATION FOR SEQ ID NO: 179:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 175 amino acids

(B) TYPE: amino acid

20 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 179:

Val Val Phe Gly Ala Ser Leu Phe Leu Leu Leu Ser Leu Thr Val Phe
 1 5 10 15

25 Ser Ile Val Ser Val Thr Ala Tyr Ile Ala Leu Ala Leu Leu Ser Val
 20 25 30

Thr Ile Ser Phe Arg Ile Tyr Lys Gly Val Ile Gln Ala Ile Gln Lys
 35 40 45

Ser Asp Glu Gly His Pro Phe Arg Ala Tyr Leu Glu Ser Glu Val Ala
 50 55 60

35 Ile Ser Glu Glu Leu Val Gln Lys Tyr Ser Asn Ser Ala Leu Gly His
 65 70 75 80

Val Asn Cys Thr Ile Lys Glu Leu Arg Arg Leu Phe Leu Val Asp Asp
 85 90 95

40 Leu Val Asp Ser Leu Lys Phe Ala Val Leu Met Trp Val Phe Thr Tyr
 100 105 110

Val Gly Ala Leu Phe Asn Gly Leu Thr Leu Leu Ile Leu Ala Leu Ile
 115 120 125

45 Ser Leu Phe Ser Val Pro Val Ile Tyr Glu Arg His Gln Ala Gln Ile
 130 135 140

50 Asp His Tyr Leu Gly Leu Ala Asn Lys Asn Val Lys Asp Ala Met Ala
 145 150 155 160

Lys Ile Gln Ala Lys Ile Pro Gly Leu Lys Arg Lys Ala Glu Xaa
 165 170 175

55

(2) INFORMATION FOR SEQ ID NO: 180:

60 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 219 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 180:

5 Met Glu Ala Pro Gly Ala Pro Pro Arg Thr Leu Thr Trp Glu Ala Met
1 5 10 15

10 Glu Gln Ile Arg Tyr Leu His Glu Glu Phe Pro Glu Ser Trp Ser Val
20 25 30

Pro Arg Leu Ala Glu Gly Phe Asp Val Ser Thr Asp Val Ile Arg Arg
35 40 45

15 Val Leu Lys Ser Lys Phe Leu Pro Thr Leu Glu Gln Lys Leu Lys Gln
50 55 60

Asp Gln Lys Val Leu Lys Lys Ala Gly Leu Ala His Ser Leu Gln His
65 70 75 80

20 Leu Arg Gly Ser Gly Asn Thr Ser Lys Leu Leu Pro Ala Gly His Ser
85 90 95

Val Ser Gly Ser Leu Leu Met Pro Gly His Glu Ala Ser Ser Lys Asp
100 105 110

25 Pro Asn His Ser Thr Ala Leu Lys Val Ile Glu Ser Asp Thr His Arg
115 120 125

30 Thr Asn Thr Pro Arg Arg Arg Lys Gly Arg Asn Lys Glu Ile Gln Asp
130 135 140

Leu Glu Glu Ser Phe Val Pro Val Ala Ala Pro Leu Gly His Pro Arg
145 150 155 160

35 Glu Leu Gln Lys Tyr Ser Ser Asp Ser Glu Ser Pro Arg Gly Thr Gly
165 170 175

Ser Gly Ala Leu Pro Ser Gly Gln Lys Leu Glu Glu Leu Lys Ala Glu
180 185 190

40 Glu Pro Asp Asn Phe Ser Ser Lys Val Val Gln Arg Gly Arg Glu Phe
195 200 205

45 Phe Asp Ser Asn Gly Asn Phe Leu Tyr Arg Ile
210 215

50 (2) INFORMATION FOR SEQ ID NO: 181:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 6 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 181:

Trp Lys Ala Glu Leu Xaa
1 5

60

(2) INFORMATION FOR SEQ ID NO: 182:

5

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 44 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 182:

10

Met Ser Asn Thr Leu Leu Ser Gln Trp Leu Leu Leu Thr Leu Phe
1 5 10 15

15

Lys Cys Ile Ile Leu Pro Leu Asn Leu Xaa Pro Ile Ile Arg Thr Ile
20 25 30

Pro Asp Trp Ser Pro Glu Leu Gly Thr Asn Thr Xaa
35 40

20

(2) INFORMATION FOR SEQ ID NO: 183:

25

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 59 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 183:

30

Met Trp Gln Val Arg Arg Gly Gly Cys Val Leu Ala Val Cys Ser Gln
1 5 10 15

Ala Arg Gly Thr Gly Gly Arg Leu Gly Trp Val Gly Thr Ser Ser Leu
20 25 30

35

Arg Val Arg Met Ala Glu Ser Thr Ser Leu Met Ser Gln Gly Arg Ser
35 40 45

Pro Ile Pro Arg Met Thr Pro Ala Arg Pro Xaa
50 55

40

(2) INFORMATION FOR SEQ ID NO: 184:

45

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 588 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 184:

Met Arg Asp Ala Gly Asp Pro Ser Pro Pro Asn Lys Met Leu Arg Arg
1 5 10 15

55

Ser Asp Ser Pro Glu Asn Lys Tyr Ser Asp Ser Thr Gly His Ser Lys
20 25 30

Ala Lys Asn Val His Thr His Arg Val Arg Glu Arg Asp Gly Gly Thr
35 40 45

60

306

Ser Tyr Ser Pro Gln Glu Asn Ser His Asn His Ser Ala Leu His Ser
 50 55 60
 5 Ser Asn Ser His Ser Ser Asn Pro Ser Asn Asn Pro Ser Lys Thr Ser
 65 70 75 80
 Asp Ala Pro Tyr Asp Ser Ala Asp Asp Trp Ser Glu His Ile Ser Ser
 85 90 95
 10 Ser Gly Lys Lys Tyr Tyr Tyr Asn Cys Arg Thr Glu Val Ser Gln Trp
 100 105 110
 Glu Lys Pro Lys Glu Trp Leu Glu Arg Glu Gln Arg Gln Lys Glu Ala
 115 120 125
 15 Asn Lys Met Ala Val Asn Ser Phe Pro Lys Asp Arg Asp Tyr Arg Arg
 130 135 140
 Glu Val Met Gln Ala Thr Ala Thr Ser Gly Phe Ala Ser Gly Met Glu
 145 150 155 160
 Asp Lys His Ser Ser Asp Ala Ser Ser Leu Leu Pro Gln Asn Ile Leu
 165 170 175
 25 Ser Gln Thr Ser Arg His Asn Asp Arg Asp Tyr Arg Leu Pro Arg Ala
 180 185 190
 Glu Thr His Ser Ser Ser Thr Pro Val Gln His Pro Ile Lys Pro Val
 195 200 205
 30 Val His Pro Thr Ala Thr Pro Ser Thr Val Pro Ser Ser Pro Phe Thr
 210 215 220
 Leu Gln Ser Asp His Gln Pro Lys Lys Ser Phe Asp Ala Asn Gly Ala
 225 230 235 240
 Ser Thr Leu Ser Lys Leu Pro Thr Pro Thr Ser Ser Val Pro Ala Gln
 245 250 255
 40 Lys Thr Glu Arg Lys Glu Ser Thr Ser Gly Asp Lys Pro Val Ser His
 260 265 270
 Ser Cys Thr Thr Pro Ser Thr Ser Ser Ala Ser Gly Leu Asn Pro Thr
 275 280 285
 45 Ser Ala Pro Pro Thr Ser Ala Ser Ala Val Pro Val Ser Pro Val Pro
 290 295 300
 Gln Ser Pro Ile Pro Pro Leu Leu Gln Asp Pro Asn Leu Leu Arg Gln
 305 310 315 320
 Leu Leu Pro Ala Leu Gln Ala Thr Leu Gln Leu Asn Asn Ser Asn Val
 325 330 335
 55 Asp Ile Ser Lys Ile Asn Glu Val Leu Thr Ala Ala Val Thr Gln Ala
 340 345 350
 Ser Leu Gln Ser Ile Ile His Lys Phe Leu Thr Ala Gly Pro Ser Ala
 355 360 365
 60

307

Phe Asn Ile Thr Ser Leu Ile Ser Gln Ala Ala Gln Leu Ser Thr Gln
 370 375 380
 5 Ala Gln Pro Ser Asn Gln Ser Pro Met Ser Leu Thr Ser Asp Ala Ser
 385 390 395 400
 Ser Pro Arg Ser Tyr Val Ser Pro Arg Ile Ser Thr Pro Gln Thr Asn
 405 410 415
 10 Thr Val Pro Ile Lys Pro Leu Ile Ser Thr Pro Pro Val Ser Ser Gln
 420 425 430
 Pro Lys Val Ser Thr Pro Val Val Lys Gln Gly Pro Val Ser Gln Ser
 435 440 445
 15 Ala Thr Gln Gln Pro Val Thr Ala Asp Lys Xaa Gln Gly His Glu Pro
 450 455 460
 Val Ser Pro Arg Ser Leu Gln Arg Ser Ser Ser Gln Arg Ser Pro Ser
 20 465 470 475 480
 Pro Gly Pro Asn His Thr Ser Asn Ser Ser Asn Ala Ser Asn Ala Thr
 485 490 495
 25 Val Val Pro Gln Asn Ser Ser Ala Arg Ser Thr Cys Ser Leu Thr Pro
 500 505 510
 Ala Leu Ala Ala His Phe Ser Glu Asn Leu Ile Lys His Val Gln Gly
 515 520 525
 30 Trp Pro Ala Asp His Ala Glu Lys Gln Ala Ser Arg Leu Arg Glu Glu
 530 535 540
 Ala His Asn Met Gly Thr Ile His Met Ser Glu Ile Cys Thr Glu Leu
 35 545 550 555 560
 Lys Asn Leu Arg Ser Leu Val Arg Val Cys Glu Ile Gln Ala Thr Leu
 565 570 575
 40 Arg Glu Gln Arg Asp Thr Ile Phe Glu Thr Thr Asn
 580 585

45 (2) INFORMATION FOR SEQ ID NO: 185:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 166 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 185:

Met Asn Ile Lys His Leu Val Asp Pro Ile Asp Asp Leu Phe Leu Ala
 1 5 10 15
 55 Ala Lys Lys Ile Pro Gly Ile Ser Ser Thr Gly Val Gly Asp Gly Gly
 20 25 30
 Asn Glu Leu Gly Met Gly Lys Val Lys Glu Ala Val Arg Arg His Ile
 35 40 45
 60

308

Arg His Gly Asp Val Ile Ala Cys Asp Val Glu Ala Asp Phe Ala Val
 50 55 60
 5 Ile Ala Gly Val Ser Asn Trp Gly Gly Tyr Ala Leu Ala Cys Ala Leu
 65 70 75 80
 Tyr Ile Leu Tyr Ser Cys Ala Val His Ser Gln Tyr Leu Arg Lys Ala
 85 90 95
 10 Val Gly Pro Ser Arg Ala Pro Gly Asp Gln Ala Trp Thr Gln Ala Leu
 100 105 110
 Pro Ser Val Ile Lys Glu Glu Lys Met Leu Gly Ile Leu Val Gln His
 115 120 125
 Lys Val Arg Ser Gly Val Ser Gly Ile Val Gly Met Glu Val Asp Gly
 130 135 140
 20 Leu Pro Phe His Asn Xaa His Ala Glu Met Ile Gln Lys Leu Val Asp
 145 150 155 160
 Val Thr Thr Ala Gln Val
 165
 25

(2) INFORMATION FOR SEQ ID NO: 186:

30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 186:

Met Leu Ile Leu Phe Leu Lys Lys Xaa
 1 5

40

(2) INFORMATION FOR SEQ ID NO: 187:

45 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 187:

50 Thr His Thr His Thr His Pro Lys Ser Phe Tyr Ile Ile Lys Leu Ser
 1 5 10 15
 Tyr Tyr Tyr Xaa
 20
 55

(2) INFORMATION FOR SEQ ID NO: 188:

60 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 188:

5 Met Ile Gln Ser Gly Leu Ile Ala Ile Leu Leu Ser Phe Leu Lys Val
 1 5 10 15

Tyr Val Glu Gly Arg Pro Cys Val Cys Phe Ser Lys Gly Leu Xaa Xaa
 20 25 30

10

15

(2) INFORMATION FOR SEQ ID NO: 189:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 19 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 189:

20

25

Tyr Ile Tyr Leu Ile Val Tyr Ile Ser Phe Tyr Ser Phe Arg Pro Gln
 1 5 10 15

Gln Leu Xaa

30

(2) INFORMATION FOR SEQ ID NO: 190:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 190:

35

40

Met Arg Phe Leu Leu Thr Val Trp Gly Ser Phe Pro Phe Met Leu Ile
 1 5 10 15

Pro Val Phe Leu Ser Ile Gly Thr Lys Glu Met Lys Lys Ala Gln Arg
 20 25 30

45

Xaa

50

(2) INFORMATION FOR SEQ ID NO: 191:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 84 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 191:

55

60

Met Arg Val Pro Pro Val Leu Arg Gly Arg Ile Leu Pro Leu Val Leu
 1 5 10 15

310

5 Gln Cys Thr Leu Leu Glu Phe Cys Leu Cys Ala Thr Thr Val Leu Pro
 20 25 30
 10 Thr Val Xaa Cys Trp Lys Pro Arg Leu Pro Val Xaa Ala Ser Gly Leu
 35 40 45
 15 Tyr Val Asp Arg Met Ser Leu Trp Lys Tyr Gly Cys Ser Gly Trp Asn
 50 55 60
 20 Glu Ser Ala Arg Pro Arg Arg Ala Gly Gly Thr Met Arg Pro Pro Arg
 65 70 75 80
 25 Ser Gly Arg Xaa

(2) INFORMATION FOR SEQ ID NO: 192:

20

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 123 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 192:

Met Ala Gly Ala Phe Val Ala Val Phe Leu Leu Ala Met Phe Tyr Glu
 1 5 10 15
 30 Gly Leu Lys Ile Ala Arg Glu Ser Leu Leu Arg Lys Ser Gln Val Ser
 20 25 30
 35 Ile Arg Tyr Asn Ser Met Pro Val Pro Gly Pro Asn Gly Thr Ile Leu
 35 40 45
 40 Met Glu Thr His Lys Thr Val Gly Gln Gln Met Leu Ser Phe Pro His
 50 55 60
 45 Leu Leu Gln Thr Val Leu His Ile Ile Gln Val Val Ile Ser Tyr Phe
 65 70 75 80
 50 Leu Met Leu Ile Phe Met Thr Tyr Asn Gly Tyr Leu Cys Ile Ala Xaa
 85 90 95
 55 Ala Ala Gly Ala Gly Thr Gly Tyr Phe Leu Phe Ser Trp Lys Lys Ala
 100 105 110
 60 Val Val Val Asp Ile Thr Glu His Cys His Xaa
 115 120

(2) INFORMATION FOR SEQ ID NO: 193:

55

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 143 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 193:

311

Met Gly Cys Leu Val Trp Gly Pro Ser Trp Pro Pro Leu Ser Leu Leu
 1 5 10 15

Ala Ser Leu Leu His Ser Gly Ile Ala Gly Arg Cys Leu Leu Cys Leu
 5 20 25 30

Phe Lys Gly Leu Ala Ala Ala Ala Ser Leu Gln Ile Arg Asp Leu Ala
 35 40 45

Ser Arg Leu Thr Thr Gly Pro Arg Thr Cys Arg Val Gln Pro Pro Pro
 10 50 55 60

His Pro Gln Ser Ser Pro Trp Pro Gly Pro Pro Gly Ala Glu Thr
 65 70 75 80

Cys Arg Pro Leu Ser Arg Thr Val Gly Gly Val Cys Pro Ser Asp Trp
 15 85 90 95

Pro Val Ser Trp Leu Leu Leu Pro Pro Leu Pro Glu Val Val Thr Cys
 20 100 105 110

Ser Cys Pro Arg Ile Lys Ala Arg Pro Glu Arg Thr Pro Glu Leu Leu
 115 120 125

Cys Ala Trp Gly Gly Arg Gly Lys His Ser Gln Leu Val Ala Xaa
 25 130 135 140

30 (2) INFORMATION FOR SEQ ID NO: 194:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 51 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 194:

Met Pro Asn Val Met Leu Thr Leu Phe Val Met Thr Leu Ser Ser Ala
 1 5 10 15

Ser Asn Leu Gly Leu Tyr Phe Phe Lys Phe Asn Phe Glu Cys Ser Cys
 20 25 30

Met Phe Gly Thr Ser Leu Leu Thr Ala Lys Asp Lys Leu Phe Ile Cys
 45 35 40 45

Ile Thr Xaa
 50

50

(2) INFORMATION FOR SEQ ID NO: 195:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 222 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 195:

60 Met Ser Leu Leu Val Leu Val Leu Ser Trp Gly Ser Met Gly Leu Glu

312

1 5 10 15

Ala Ala Thr Ala Val Gly Leu Ser Asp Phe Cys Ser Asn Pro Asp Pro
20 25 30

5 Tyr Val Leu Asn Leu Thr Gln Glu Glu Thr Gly Leu Ser Ser Asp Ile
35 40 45

10 Leu Ser Tyr Tyr Leu Leu Cys Asn Arg Ala Val Ser Asn Pro Phe Gln
50 55 60

Gln Arg Leu Thr Leu Ser Gln Arg Ala Leu Ala Asn Ile His Ser Gln
65 70 75 80

15 Leu Leu Gly Leu Glu Arg Glu Ala Val Pro Gln Phe Pro Ser Ala Gln
85 90 95

Lys Pro Leu Leu Ser Leu Glu Glu Thr Leu Asn Val Thr Glu Gly Asn
100 105 110

20 Phe His Gln Leu Val Ala Leu Leu His Cys Arg Ser Leu His Lys Asp
115 120 125

25 Tyr Gly Ala Ala Leu Arg Gly Leu Cys Glu Xaa Xaa Leu Glu Gly Leu
130 135 140

Leu Phe Leu Leu Leu Phe Ser Leu Leu Ser Ala Gly Ala Leu Ala Xaa
145 150 155 160

30 Ala Leu Cys Xaa Leu Pro Arg Ala Trp Ala Leu Phe Pro Pro Arg Asn
165 170 175

Pro Ser Ala Leu Cys Ser Gly Ser Arg Leu Ser Glu Pro Leu Leu Pro
180 185 190

35 Ala Gly Leu Glu Pro Gly Ser Pro Leu Arg Ser Phe Pro Gly Cys Arg
195 200 205

40 Arg Asp Pro Thr Asn Pro Ala Cys Leu Gly Ser Asp His Xaa
210 215 220

(2) INFORMATION FOR SEQ ID NO: 196:

45

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 102 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 196:

Met Ser Gln Leu Ser Arg Thr Ser Leu Ser Leu Leu Thr Leu Leu
1 5 10 15

55 Val Leu Trp Gly Ser Ser Cys Cys Leu Pro Ile Trp Cys Leu Pro Asn
20 25 30

Arg His Arg Leu Leu Lys Leu Ser Phe Leu Leu Phe Ser Pro Asp Ile
35 40 45

60

Pro Tyr Leu Ser His Thr His Pro Asn Asn Ile Ser Cys Ser Val Leu
 50 55 60

5 Ser Leu Arg Gln His Leu Asn Phe Thr Gln Pro Gly Ala Leu Phe Thr
 65 70 75 80

Cys Leu Val Gln Ile Gln Phe Gly Leu Ile Leu Gln Pro Cys Ile Ser
 85 90 95

10 Lys Trp Gly Leu Gly Xaa
 100

15 (2) INFORMATION FOR SEQ ID NO: 197:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 13 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 197:

Met Ile Ala Leu Phe Phe Val Thr Thr Xaa Leu Thr Xaa
 1 5 10

25

(2) INFORMATION FOR SEQ ID NO: 198:

30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 61 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 198:

35 Met Thr Tyr His Pro Asn Gln Val Val Glu Gly Cys Cys Ser Asp Met
 1 5 10 15

40 Ala Val Thr Phe Asn Gly Leu Thr Pro Asn Gln Met His Val Met Met
 20 25 30

Tyr Gly Val Tyr Arg Leu Arg Ala Phe Gly His Ile Phe Asn Asp Ala
 35 40 45

45 Leu Val Phe Leu Pro Pro Asn Gly Ser Asp Asn Asp Xaa
 50 55 60

50 (2) INFORMATION FOR SEQ ID NO: 199:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 199:

Met Ser Ser Ser Ser Leu His Trp Lys Glu Phe Lys Tyr Ala Pro Gly
 1 5 10 15

60

Ser Leu His Tyr Phe Ala Leu Ser Phe Val Leu Ile Leu Thr Glu Ile
 20 25 30
 Cys Leu Val Ser Ser Gly Met Gly Phe Pro Gln Glu Gly Lys His Phe
 5 35 40 45
 Ser Val Leu Gly Ser Pro Asp Cys Ser Leu Trp Gly Arg Asp Glu His
 50 55 60
 10 Val Pro Arg Glu Phe Ala Xaa
 65 70

15 (2) INFORMATION FOR SEQ ID NO: 200:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 200:

Met His Leu Arg Phe Pro Phe Leu Cys Xaa
 1 5 10

25

(2) INFORMATION FOR SEQ ID NO: 201:

30 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 50 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 201:

35

Met Arg Arg Val Ala Arg Gly Arg Gly Leu Ala Leu Pro Ser Leu Glu
 1 5 10 15

40

His Arg Pro Ser Cys Ser Tyr Asp Ala Leu Pro Leu Pro Phe Cys Glu
 20 25 30

Thr Arg Asn Pro Glu Ala His Leu Tyr Phe Phe Arg Thr Asp Val Glu
 35 40 45

45

Arg Xaa
 50

50 (2) INFORMATION FOR SEQ ID NO: 202:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 202:

Ala Lys Ile Leu Val Phe Ile Phe Leu Phe Glu Leu Xaa
 1 5 10

60

(2) INFORMATION FOR SEQ ID NO: 203:

5

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 38 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 203:

Met Phe Gln Glu Cys Ile Pro Ile Ser Leu Phe Phe Leu Asn Trp Leu
 1 5 10 15

15

Lys Glu Cys Cys Ser Phe Thr Cys Pro Asn Ser His Ile Asn Asn Cys
 20 25 30

Leu Thr Gly Ile Arg Xaa
 35

20

(2) INFORMATION FOR SEQ ID NO: 204:

25

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 34 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 204:

30

Met Asn Phe Val Leu Phe Phe Ile Gly Ile Asn Val Gly Cys Arg Gly
 1 5 10 15

Glu Asn Ser Leu Lys Tyr Phe Thr Val Thr Val Xaa Cys Ser Pro Arg
 20 25 30

35

Asp Xaa

40

(2) INFORMATION FOR SEQ ID NO: 205:

45

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 205:

50

Met Leu Leu Phe Leu Phe Val Cys Leu Pro Ile Thr Trp Met Ala Glu
 1 5 10 15

Phe Leu Ser Gln Leu Arg His Leu Leu Xaa
 20 25

55

(2) INFORMATION FOR SEQ ID NO: 206:

60

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 105 amino acids

316

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 206:

5 Met Pro Arg His Ser Leu Tyr Ile Ile Ile Gly Ala Leu Cys Val Ala
 1 5 10 15
 Phe Ile Leu Met Leu Ile Ile Leu Ile Val Gly Ile Cys Arg Ile Ser
 20 25 30
 10 Arg Ile Glu Tyr Gln Gly Ser Ser Arg Pro Ala Tyr Glu Glu Phe Tyr
 35 40 45
 Asn Cys Arg Ser Ile Asp Ser Glu Phe Ser Asn Ala Ile Ala Ser Ile
 15 50 55 60
 Arg His Ala Arg Phe Gly Lys Lys Ser Arg Pro Ala Met Tyr Asp Val
 65 70 75 80
 20 Ser Pro Ile Ala Tyr Glu Asp Tyr Ser Pro Asp Asp Lys Pro Leu Val
 85 90 95
 Thr Leu Ile Lys Thr Lys Asp Leu Xaa
 100 105
 25

(2) INFORMATION FOR SEQ ID NO: 207:

30 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 64 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 207:

35 Leu Lys Ser Cys Leu Leu Leu Val Ser Phe Leu Ser Gly Arg Val Pro
 1 5 10 15
 40 Ser Tyr Asp Leu Ile Tyr Val Cys Ser Ile Ala Leu Glu Thr Gly Phe
 20 25 30
 Val Cys Glu Met Ala Leu Ser Phe Val Asp His Phe Cys Arg Glu Ile
 35 40 45
 45 Val Asp Leu Gly Arg Ala Glu Ala Thr Ala Asp Met Pro Gly Val Xaa
 50 55 60

50

(2) INFORMATION FOR SEQ ID NO: 208:

55 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 42 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 208:

60

Met Ser Ala Trp Leu Pro Ser Pro Pro His Leu Leu Leu Leu Ser Ala
 1 5 10 15

5 Ala Ala Gly Ser Gly Ala Ser His Leu Arg Ala Leu Gly Ser Ser Ala
 20 25 30

Leu Glu Gly Leu Gln Asp Pro Ser Gln Xaa
 35 40

10

(2) INFORMATION FOR SEQ ID NO: 209:

15 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 42 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 209:

20 Met Ser Ser Pro Ala Thr Trp Arg Leu Thr Leu Pro Ser Leu Leu Val
 1 5 10 15

Phe Leu Thr Gly Glu Ala Met Pro Trp Pro Ala His Ser Thr Ser Cys
 20 25 30

25

Thr His Val Leu Ser Thr Val Ser Thr Xaa
 35 40

30

(2) INFORMATION FOR SEQ ID NO: 210:

35 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 46 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 210:

40 Met Gln Ala Pro Leu Gln Asp Cys Gly Arg Ser Val Ser Leu Arg Leu
 1 5 10 15

Ala Cys Val Leu Ala Pro Leu Thr Thr Ser Ser Arg Gly Cys His Leu
 20 25 30

45 Gln Leu Pro Gln Asp Lys Gly Lys Ala Arg Xaa Asp Ser Xaa
 35 40 45

50

(2) INFORMATION FOR SEQ ID NO: 211:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 266 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 211:

Met Asn Gly Ser His Lys Asp Pro Leu Leu Pro Phe Pro Ala Ser Ala
 1 5 10 15

60

318

Arg Thr Pro Ser Leu Pro Pro Ala Pro Pro Ala Gln Ala Pro Leu Pro
 20 25 30
 5 Trp Lys Pro Ser Gly Phe Ala Arg Ile Ser Pro Pro Pro Pro Leu Ala
 35 40 45
 Ile Leu Gln Tyr Arg Gly Lys Ala Asp His Gly Glu Ser Gly Gln Gln
 50 55 60
 10 Leu Ala Ala Ala Pro Gly Asp Gly Arg Leu Pro Leu Leu Glu Ala Val
 65 70 75 80
 Arg Arg Leu Arg Gly Gln Asp Cys Gly Pro Leu Ser Ala Leu Cys His
 85 90 95
 15 Gly Gln Leu Leu Ala Gln Pro Val Pro Gln Val Leu Leu Leu Pro Gly
 100 105 110
 Ala Xaa Gly Asp Ile Gly Thr Ser Cys Tyr Thr Lys Ser Gly Met Ile
 115 120 125
 20 Leu Cys Arg Asn Asp Tyr Ile Arg Leu Phe Gly Asn Ser Gly Ala Cys
 130 135 140
 25 Ser Ala Cys Gly Gln Ser Ile Pro Ala Ser Glu Leu Val Met Arg Ala
 145 150 155 160
 Gln Gly Asn Val Tyr His Leu Lys Cys Phe Thr Cys Ser Thr Cys Arg
 165 170 175
 30 Asn Arg Leu Val Pro Gly Asp Arg Phe His Tyr Ile Asn Gly Ser Leu
 180 185 190
 35 Phe Cys Glu His Asp Arg Pro Thr Ala Leu Ile Asn Gly His Leu Asn
 195 200 205
 Ser Leu Gln Ser Asn Pro Leu Leu Pro Asp Gln Lys Val Cys Lys Val
 210 215 220
 40 Arg Val Met Gln Asn Ala Cys Leu His Leu Arg Phe Val His His Arg
 225 230 235 240
 Trp Ile Pro Cys Xaa Phe Ser Arg Gln Val Thr Phe Val Ala Ser Thr
 245 250 255
 45 Ser Ala Ser Ser Met Pro Leu His Leu Leu
 260 265

50

(2) INFORMATION FOR SEQ ID NO: 212:

(i) SEQUENCE CHARACTERISTICS:

55 (A) LENGTH: 94 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 212:

60 Met Ala Arg Thr Arg Thr Pro Ser Ser Pro Phe Leu Leu Leu Arg Glu
 1 5 10 15

Leu Pro Pro Ser Leu Gln Leu Arg Gln Pro Arg Arg Pro Phe Pro Gly
 20 25 30
 5 Ser Arg Ala Ala Ser Leu Ala Phe His Arg Arg Arg Leu Ser Gln Tyr
 35 40 45
 Cys Asn Ile Gly Glu Lys Gln Thr Met Val Asn Pro Gly Ser Ser Ser
 50 55 60
 10 Gln Pro Pro Pro Val Thr Ala Gly Ser Leu Ser Trp Lys Arg Cys Ala
 65 70 75 80
 15 Gly Cys Gly Gly Lys Ile Ala Asp Arg Phe Leu Leu Tyr Ala
 85 90

20 (2) INFORMATION FOR SEQ ID NO: 213:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 213:

Leu Phe Gly Asn Ser Gly Ala Cys Ser Ala Cys Gly Gln Ser Ile Pro
 1 5 10 15
 30 Ala Ser Glu Leu Val Met Arg Ala
 20

35 (2) INFORMATION FOR SEQ ID NO: 214:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 19 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 214:

His Asp Arg Pro Thr Ala Leu Ile Asn Gly His Leu Asn Ser Leu Gln
 1 5 10 15
 45 Ser Asn Pro

50 (2) INFORMATION FOR SEQ ID NO: 215:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 215:

Leu Val Pro Gly Asp Arg Phe His Tyr Ile Asn Gly
 1 5 10

60

(2) INFORMATION FOR SEQ ID NO: 216:

5

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 81 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 216:

Met Lys Tyr Met Gly Gly Cys Ala Lys Val Met Cys Lys Tyr Tyr Val
1 5 10 15

15

Ile Leu Tyr Gln Gly Leu Glu Tyr Pro Leu Leu Xaa Ser Gly Asp Pro
20 25 30

Glu Thr Ser Pro Pro Trp Ile Leu Arg Ala Asp Cys Ile Val Leu Ser
35 40 45

20

Ser Arg Asn Phe His Ser Asn Xaa Gly Arg Leu Thr Ile Asn Lys Ile
50 55 60

25

Tyr Val Ile Gly Gly Gly Lys Tyr Arg Gly Glu Val Thr Asn Gly Ala
65 70 75 80

Lys

30

(2) INFORMATION FOR SEQ ID NO: 217:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 41 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 217:

40

Met Gly Gln Ser Glu Leu Tyr Ser Ser Ile Leu Arg Asn Leu Gly Val
1 5 10 15

Leu Phe Leu Val Tyr Thr Arg Gly Gly Phe Leu Leu Ser Pro Leu Leu
20 25 30

45

His Gly Thr Leu Thr Cys Ala His Ser
35 40

50

(2) INFORMATION FOR SEQ ID NO: 218:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 35 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 218:

60

Met Val Leu Leu Leu Leu Thr Val Ala Ser Tyr Thr Val Phe Trp Met
1 5 10 15

321

Ile Gly Asp Val Leu Asp Ile Leu Phe Leu Trp Asn Phe Glu Tyr Thr
 20 25 30

5 Thr Leu Tyr
 35

10 (2) INFORMATION FOR SEQ ID NO: 219:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 38 amino acids

(B) TYPE: amino acid

15 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 219:

Met Glu Leu Tyr Asn Ser Leu Cys Pro Ile Cys Tyr Phe Ser Thr Val
 1 5 10 15

20 Leu Thr Thr Thr Tyr Tyr Ile Tyr Phe Val Tyr Ser Gln Ser Ser Xaa
 20 25 30

25 Ile Arg Met Lys Val Pro
 35

30 (2) INFORMATION FOR SEQ ID NO: 220:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 45 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 220:

Met Gln Ile Val Ile Val Leu Tyr Cys Val Arg Asn Lys Asp Lys Lys
 1 5 10 15

40 Lys Val Cys Thr Cys Ser Val Gln Thr Gln Phe Phe Phe Pro Ile Phe
 20 25 30

45 Pro Ile Leu Gly Cys Leu Asn Gly Cys Arg Thr Gln Glu
 35 40 45

(2) INFORMATION FOR SEQ ID NO: 221:

50 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 221:

Met Lys Tyr Met Gly Gly Cys Ala Lys Val Met Cys Lys Tyr Tyr Val
 1 5 10 15

60 Ile Leu Tyr Gln Gly Leu Glu Tyr Pro Leu Leu Xaa
 20 25

(2) INFORMATION FOR SEQ ID NO: 222:

5

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 35 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 222:

Leu Glu Tyr Pro Leu Leu Xaa Ser Gly Asp Pro Glu Thr Ser Pro Pro
 1 5 10 15

15

Trp Ile Leu Arg Ala Asp Cys Ile Val Leu Ser Ser Arg Asn Phe His
 20 25 30

Ser Asn Xaa
 35

20

(2) INFORMATION FOR SEQ ID NO: 223:

25

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 32 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 223:

Arg Asn Phe His Ser Asn Xaa Gly Arg Leu Thr Ile Asn Lys Ile Tyr
 1 5 10 15

35

Val Ile Gly Gly Gly Lys Tyr Arg Gly Glu Val Thr Asn Gly Ala Lys
 20 25 30

40

(2) INFORMATION FOR SEQ ID NO: 224:

45

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 145 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 224:

50

Val Thr Asn Glu Met Ser Gln Gly Arg Gly Lys Tyr Asp Phe Tyr Ile
 1 5 10 15

Gly Leu Gly Leu Ala Met Ser Ser Ser Ile Phe Ile Gly Gly Ser Phe
 20 25 30

55

Ile Leu Lys Lys Lys Gly Leu Leu Arg Leu Ala Arg Lys Gly Ser Met
 35 40 45

60

Arg Ala Gly Gln Gly Gly His Ala Tyr Leu Lys Glu Trp Leu Trp Trp
 50 55 60

Ala Gly Leu Leu Ser Met Gly Ala Gly Glu Val Ala Asn Phe Ala Ala
 65 70 75 80

5 Tyr Ala Phe Ala Pro Ala Thr Leu Val Thr Pro Leu Gly Ala Leu Ser
 85 90 95

Val Leu Val Ser Ala Ile Leu Ser Ser Tyr Phe Leu Asn Glu Arg Leu
 100 105 110

10 Asn Leu His Gly Lys Ile Gly Cys Leu Leu Ser Ile Leu Gly Ser Thr
 115 120 125

Val Met Val Ile His Ala Pro Lys Glu Glu Glu Ile Glu Thr Leu Asn
 15 130 135 140

Glu
 145

20

(2) INFORMATION FOR SEQ ID NO: 225:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 78 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 225:

30 Val Thr Asn Glu Met Ser Gln Gly Arg Gly Lys Tyr Asp Phe Tyr Ile
 1 5 10 15

Gly Leu Gly Leu Ala Met Ser Ser Ser Ile Phe Ile Gly Gly Ser Phe
 20 25 30

35 Ile Leu Lys Lys Lys Gly Leu Leu Arg Leu Ala Arg Lys Gly Ser Met
 35 40 45

40 Arg Ala Gly Gln Gly Gly His Ala Tyr Leu Lys Glu Trp Leu Trp Trp
 50 55 60

Ala Gly Leu Leu Ser Met Gly Ala Gly Glu Val Ala Asn Phe
 65 70 75

45

(2) INFORMATION FOR SEQ ID NO: 226:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 30 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 226:

50 Asn Phe Ala Ala Tyr Ala Phe Ala Pro Ala Thr Leu Val Thr Pro Leu
 1 5 10 15

Gly Ala Leu Ser Val Leu Val Ser Ala Ile Leu Ser Ser Tyr
 20 25 30

60

(2) INFORMATION FOR SEQ ID NO: 227:

5

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 36 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 227:

Glu Arg Leu Asn Leu His Gly Lys Ile Gly Cys Leu Leu Ser Ile Leu
 1 5 10 15

15

Gly Ser Thr Val Met Val Ile His Ala Pro Lys Glu Glu Glu Ile Glu
 20 25 30

Thr Leu Asn Glu
 35

20

(2) INFORMATION FOR SEQ ID NO: 228:

25

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 31 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 228:

30

Arg Phe Lys Thr Leu Met Thr Asn Lys Ser Glu Gln Asp Gly Asp Ser
 1 5 10 15

Ser Lys Thr Ile Glu Ile Ser Asp Met Lys Tyr His Ile Phe Gln
 20 25 30

35

(2) INFORMATION FOR SEQ ID NO: 229:

40

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 229:

45

Leu Val Glu Gly Lys Leu Phe Tyr Ala His Lys Val Leu Leu Val Thr
 1 5 10 15

50

Xaa Ser Asn Arg
 20

55

(2) INFORMATION FOR SEQ ID NO: 230:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 87 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

60

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 230:

5 CCTTAAAAGC TGACATTTTA TAATTGTGTT GTATAGCAGC AACTATATCC TTCCAAAAAT 60
CAAATGTTTT TTGACCATG TTCAGTT 87

10

(2) INFORMATION FOR SEQ ID NO: 231:

(i) SEQUENCE CHARACTERISTICS:

15

- (A) LENGTH: 38 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 231:

20

CCTTAAAAGC TGACATTTTA TAATTGTGTT GTATAGCA 38

25

(2) INFORMATION FOR SEQ ID NO: 232:

(i) SEQUENCE CHARACTERISTICS:

30

- (A) LENGTH: 38 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 232:

35

CTTCCAAAAA TCAAATGTTT TTGACCATG GTTCAGTT 38

40

(2) INFORMATION FOR SEQ ID NO: 233:

(i) SEQUENCE CHARACTERISTICS:

45

- (A) LENGTH: 455 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 233:

50

Met Ala Gln His Phe Ser Leu Ala Ala Cys Asp Val Val Gly Phe Asp
1 5 10 15

Leu Asp His Thr Leu Cys Arg Tyr Asn Leu Pro Glu Ser Ala Pro Leu
20 25 30

55

Ile Tyr Asn Ser Phe Ala Gln Phe Leu Val Lys Glu Lys Gly Tyr Asp
35 40 45

Lys Glu Leu Leu Asn Val Thr Pro Glu Asp Trp Asp Phe Cys Cys Lys
50 55 60

60

326

Gly Leu Ala Leu Asp Leu Glu Asp Gly Asn Phe Leu Lys Leu Ala Asn
 65 70 75 80
 5 Asn Gly Thr Val Leu Arg Ala Ser His Gly Thr Lys Met Met Thr Pro
 85 90 95
 Glu Val Leu Ala Glu Ala Tyr Gly Lys Lys Glu Trp Lys His Phe Leu
 100 105 110
 10 Ser Asp Thr Gly Met Ala Cys Arg Ser Gly Lys Tyr Tyr Phe Tyr Asp
 115 120 125
 Asn Tyr Phe Asp Leu Pro Gly Ala Leu Leu Cys Ala Arg Val Val Asp
 130 135 140
 15 Tyr Leu Thr Lys Leu Asn Asn Gly Gln Lys Thr Phe Asp Phe Trp Lys
 145 150 155 160
 Asp Ile Val Ala Ala Ile Gln His Asn Tyr Lys Met Ser Ala Phe Lys
 165 170 175
 20 Glu Asn Cys Gly Ile Tyr Phe Pro Glu Ile Lys Arg Asp Pro Gly Arg
 180 185 190
 25 Tyr Leu His Ser Cys Pro Glu Ser Val Lys Lys Trp Leu Arg Gln Leu
 195 200 205
 Lys Asn Ala Gly Lys Ile Leu Leu Leu Ile Thr Ser Ser His Ser Asp
 210 215 220
 30 Tyr Cys Arg Leu Leu Cys Glu Tyr Ile Leu Gly Asn Asp Phe Thr Asp
 225 230 235 240
 Leu Phe Asp Ile Val Ile Thr Asn Ala Leu Lys Pro Gly Phe Phe Ser
 245 250 255
 35 His Leu Pro Ser Gln Arg Pro Phe Arg Thr Leu Glu Asn Asp Glu Glu
 260 265 270
 40 Gln Glu Ala Leu Pro Ser Leu Asp Lys Pro Gly Trp Tyr Ser Gln Gly
 275 280 285
 Asn Ala Val His Leu Tyr Glu Leu Leu Lys Lys Met Thr Gly Lys Pro
 290 295 300
 45 Glu Pro Lys Val Val Tyr Phe Gly Asp Ser Met His Ser Asp Ile Phe
 305 310 315 320
 Pro Ala Arg His Tyr Ser Asn Trp Glu Thr Val Leu Ile Leu Glu Glu
 325 330 335
 50 Leu Arg Gly Asp Glu Gly Thr Arg Ser Gln Arg Pro Glu Glu Ser Glu
 340 345 350
 55 Pro Leu Glu Lys Lys Gly Lys Tyr Glu Gly Pro Lys Ala Lys Pro Leu
 355 360 365
 Asn Thr Ser Ser Lys Lys Trp Gly Ser Phe Phe Ile Asp Ser Val Leu
 370 375 380
 60

327

Gly Leu Glu Asn Thr Glu Asp Ser Leu Val Tyr Thr Trp Ser Cys Lys
 385 390 395 400
 5 Arg Ile Ser Thr Tyr Ser Thr Ile Ala Ile Pro Ser Ile Glu Ala Ile
 405 410 415
 Ala Glu Leu Pro Leu Asp Tyr Lys Phe Thr Arg Phe Ser Ser Ser Asn
 420 425 430
 10 Ser Lys Thr Ala Gly Tyr Tyr Pro Asn Pro Pro Leu Val Leu Ser Ser
 435 440 445
 Asp Glu Thr Leu Ile Ser Lys
 450 455
 15

(2) INFORMATION FOR SEQ ID NO: 234:

20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 27 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 234:
 25 Thr Ser Ser His Ser Asp Tyr Cys Arg Leu Leu Cys Glu Tyr Ile Leu
 1 5 10 15
 Gly Asn Asp Phe Thr Asp Leu Phe Asp Ile Val
 30 20 25

(2) INFORMATION FOR SEQ ID NO: 235:

35 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 327 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 235:
 Met Lys Thr Lys Asn Ile Pro Glu Ala His Gln Asp Ala Phe Lys Thr
 1 5 10 15
 45 Gly Phe Ala Glu Gly Phe Leu Lys Ala Gln Ala Leu Thr Gln Lys Thr
 20 25 30
 Asn Asp Ser Leu Arg Arg Thr Arg Leu Ile Leu Phe Val Leu Leu Leu
 35 40 45
 50 Phe Gly Ile Tyr Gly Leu Leu Lys Asn Pro Phe Leu Ser Val Arg Phe
 50 55 60
 Arg Thr Thr Thr Gly Leu Asp Ser Ala Val Asp Pro Val Gln Met Lys
 55 65 70 75 80
 Asn Val Thr Phe Glu His Val Lys Gly Val Glu Glu Ala Lys Gln Glu
 85 90 95
 60 Leu Gln Glu Val Val Glu Phe Leu Lys Asn Pro Gln Lys Phe Thr Ile

328

100 105 110
 Leu Gly Gly Lys Leu Pro Lys Gly Ile Leu Leu Val Gly Pro Pro Gly
 115 120 125
 5 Thr Gly Lys Thr Leu Leu Ala Arg Ala Val Ala Gly Glu Ala Asp Val
 130 135 140
 10 Pro Phe Tyr Tyr Ala Ser Gly Ser Glu Phe Asp Glu Met Phe Val Gly
 145 150 155 160
 Val Gly Ala Ser Arg Ile Arg Asn Leu Phe Arg Glu Ala Lys Ala Asn
 165 170 175
 15 Ala Pro Cys Val Ile Phe Ile Asp Glu Leu Asp Ser Val Gly Gly Lys
 180 185 190
 Arg Ile Glu Ser Pro Met His Pro Tyr Ser Arg Gln Thr Ile Asn Gln
 195 200 205
 20 Leu Leu Ala Glu Met Asp Gly Phe Lys Pro Asn Glu Gly Val Ile Ile
 210 215 220
 25 Ile Gly Ala Thr Asn Phe Pro Glu Ala Leu Asp Asn Ala Leu Ile Arg
 225 230 235 240
 Pro Gly Arg Phe Asp Met Gln Val Thr Val Pro Arg Pro Asp Val Lys
 245 250 255
 30 Gly Arg Thr Glu Ile Leu Lys Trp Tyr Leu Asn Lys Ile Lys Phe Asp
 260 265 270
 Xaa Ser Val Asp Pro Glu Ile Ile Ala Arg Gly Thr Val Gly Phe Ser
 275 280 285
 35 Gly Ala Glu Leu Glu Asn Leu Val Asn Gln Ala Ala Leu Lys Ala Ala
 290 295 300
 Val Asp Gly Lys Glu Met Val Thr Met Lys Glu Leu Gly Val Phe Gln
 40 305 310 315 320
 Arg Gln Asn Ser Asn Gly Ala
 325

45

(2) INFORMATION FOR SEQ ID NO: 236:

- 50 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 236:

55 Met Lys Thr Lys Asn Ile Pro Glu Ala His Gln Asp Ala Phe Lys Thr
 1 5 10 15

Gly Phe Ala Glu Gly
 20

60

(2) INFORMATION FOR SEQ ID NO: 237:

5

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 237:

10

Pro Val Gln Met Lys Asn Val Thr Phe Glu His Val Lys Gly Val Glu
1 5 10 15

15

Glu Ala Lys Gln Glu Leu Gln
20

(2) INFORMATION FOR SEQ ID NO: 238:

20

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 238:

Ser Arg Gln Thr Ile Asn Gln Leu Leu Ala Glu Met Asp Gly Phe Lys
1 5 10 15

30

Pro Asn Glu Gly Val Ile Ile
20

35

(2) INFORMATION FOR SEQ ID NO: 239:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 239:

Phe Ser Gly Ala Glu Leu Glu Asn Leu Val Asn Gln Ala Ala Leu Lys
1 5 10 15

45

Ala Ala Val Asp Gly Lys Glu Met
20

50

(2) INFORMATION FOR SEQ ID NO: 240:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 192 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 240:

60

Leu Pro Met Trp Gln Val Thr Ala Phe Leu Asp His Asn Ile Val Thr
1 5 10 15

330

Ala Gln Thr Thr Trp Lys Gly Leu Trp Met Ser Cys Val Val Gln Ser
 20 25 30

5 Thr Gly His Met Gln Cys Lys Val Tyr Asp Ser Val Leu Ala Leu Ser
 35 40 45

Thr Glu Val Gln Ala Ala Arg Ala Leu Thr Val Ser Ala Val Leu Leu
 50 55 60

10 Ala Phe Val Ala Leu Phe Val Thr Leu Ala Gly Ala Gln Cys Thr Thr
 65 70 75 80

15 Cys Val Ala Pro Gly Pro Ala Lys Ala Arg Val Ala Leu Thr Gly Gly
 85 90 95

Val Leu Tyr Leu Phe Cys Gly Leu Leu Ala Leu Val Pro Leu Cys Trp
 100 105 110

20 Phe Ala Asn Ile Val Val Arg Glu Phe Tyr Asp Pro Ser Val Pro Val
 115 120 125

Ser Gln Lys Tyr Glu Leu Gly Ala Xaa Leu Tyr Ile Gly Trp Ala Ala
 130 135 140

25 Thr Ala Leu Leu Met Val Gly Gly Cys Leu Leu Cys Cys Gly Ala Trp
 145 150 155 160

30 Val Cys Thr Gly Arg Pro Asp Leu Ser Phe Pro Val Lys Tyr Ser Ala
 165 170 175

Pro Arg Arg Pro Thr Ala Thr Gly Asp Tyr Asp Lys Lys Asn Tyr Val
 180 185 190

35

40 (2) INFORMATION FOR SEQ ID NO: 241:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 24 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 241:

Leu His Tyr Phe Ala Leu Ser Phe Val Leu Ile Leu Thr Glu Ile Cys
 1 5 10 15

50

Leu Val Ser Ser Gly Met Gly Phe
 20

55

(2) INFORMATION FOR SEQ ID NO: 242:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 31 amino acids

(B) TYPE: amino acid

60

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 242:

5 Gln Leu Arg Asn Gly Ile Pro Pro Gly Arg Lys Ala Leu Phe Cys Ser
 1 5 10 15
 Gly Lys Pro Arg Leu Phe Thr Leu Gly Gln Gly Arg Thr Cys Ala
 20 25 30

10

(2) INFORMATION FOR SEQ ID NO: 243:

(i) SEQUENCE CHARACTERISTICS:

15

(A) LENGTH: 39 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 243:

20 Trp Ser Gly Leu Trp Val Thr Thr Trp Asn Gly Ser Ser Gly Glu Arg
 1 5 10 15
 Thr Pro Ser Pro Trp Arg Arg Lys Arg Ala Ser Gln Ser Ala Gly Arg
 20 25 30
 Ile Ala Ser Trp Met Ser Phe
 35

30

(2) INFORMATION FOR SEQ ID NO: 244:

(i) SEQUENCE CHARACTERISTICS:

35

(A) LENGTH: 14 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 244:

40 Glu Tyr Asn Lys Glu Ser Glu Asp Lys Tyr Val Phe Leu Val
 1 5 10

45 (2) INFORMATION FOR SEQ ID NO: 245:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 14 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 245:

Ile Asp Val Glu Ile Ala Arg Ser Asp Cys Arg Lys Pro Leu
 1 5 10

55

(2) INFORMATION FOR SEQ ID NO: 246:

(i) SEQUENCE CHARACTERISTICS:

60

(A) LENGTH: 142 amino acids

332

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 246:

5 Met Pro Arg Cys Arg Trp Leu Ser Leu Ile Leu Leu Thr Ile Pro Leu
 1 5 10 15

Ala Leu Val Ala Arg Lys Asp Pro Lys Lys Asn Glu Thr Gly Val Leu
 20 25 30

10 Arg Lys Leu Lys Pro Val Asn Ala Ser Asn Ala Asn Val Lys Gln Cys
 35 40 45

Leu Trp Phe Ala Met Gln Glu Tyr Asn Lys Glu Ser Glu Asp Lys Tyr
 15 50 55 60

Val Phe Leu Val Val Lys Thr Leu Gln Ala Gln Leu Gln Val Thr Asn
 65 70 75 80

20 Leu Leu Glu Tyr Leu Ile Asp Val Glu Ile Ala Arg Ser Asp Cys Arg
 85 90 95

Lys Pro Leu Ser Thr Asn Glu Ile Cys Ala Ile Gln Glu Asn Ser Lys
 100 105 110

25 Leu Lys Arg Lys Leu Ser Cys Ser Phe Leu Val Gly Ala Leu Pro Trp
 115 120 125

Asn Gly Glu Phe Thr Val Met Glu Lys Lys Cys Glu Asp Ala
 30 130 135 140

(2) INFORMATION FOR SEQ ID NO: 247:

35

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 92 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 247:

Cys Leu Trp Phe Ala Met Gln Glu Tyr Asn Lys Glu Ser Glu Asp Lys
 1 5 10 15

45 Tyr Val Phe Leu Val Val Lys Thr Leu Gln Ala Gln Leu Gln Val Thr
 20 25 30

Asn Leu Leu Glu Tyr Leu Ile Asp Val Glu Ile Ala Arg Ser Asp Cys
 35 40 45

50 Arg Lys Pro Leu Ser Thr Asn Glu Ile Cys Ala Ile Gln Glu Asn Ser
 50 55 60

Lys Leu Lys Arg Lys Leu Ser Cys Ser Phe Leu Val Gly Ala Leu Pro
 55 65 70 75 80

Trp Asn Gly Glu Phe Thr Val Met Glu Lys Lys Cys
 85 90

60

(2) INFORMATION FOR SEQ ID NO: 248:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 123 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 248:

5
 10 Ala Arg Lys Asp Pro Lys Lys Asn Glu Thr Gly Val Leu Arg Lys Leu
 1 5 10 15
 Lys Pro Val Asn Ala Ser Asn Ala Asn Val Lys Gln Cys Leu Trp Phe
 20 25 30
 15 Ala Met Gln Glu Tyr Asn Lys Glu Ser Glu Asp Lys Tyr Val Phe Leu
 35 40 45
 Val Val Lys Thr Leu Gln Ala Gln Leu Gln Val Thr Asn Leu Leu Glu
 20 50 55 60
 Tyr Leu Ile Asp Val Glu Ile Ala Arg Ser Asp Cys Arg Lys Pro Leu
 65 70 75 80
 25 Ser Thr Asn Glu Ile Cys Ala Ile Gln Glu Asn Ser Lys Leu Lys Arg
 85 90 95
 Lys Leu Ser Cys Ser Phe Leu Val Gly Ala Leu Pro Trp Asn Gly Glu
 100 105 110
 30 Phe Thr Val Met Glu Lys Lys Cys Glu Asp Ala
 115 120

35

(2) INFORMATION FOR SEQ ID NO: 249:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 44 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 249:

40
 45 Asp Ser Pro Asp Thr Glu Pro Gly Ser Ser Ala Gly Pro Thr Gln Arg
 1 5 10 15
 Pro Ser Asp Asn Ser His Asn Glu His Ala Pro Ala Ser Gln Gly Leu
 20 25 30
 50 Lys Ala Glu His Leu Tyr Ile Leu Ile Gly Val Ser
 35 40

55

(2) INFORMATION FOR SEQ ID NO: 250:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 101 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 250:

5 His Arg Gln Asn Gln Ile Lys Gln Gly Pro Pro Arg Ser Lys Asp Glu
 1 5 10 15
 Glu Gln Lys Pro Gln Gln Arg Pro Asp Leu Ala Val Asp Val Leu Glu
 20 25 30
 10 Arg Thr Ala Asp Lys Ala Thr Val Asn Gly Leu Pro Glu Lys Asp Arg
 35 40 45
 Glu Thr Asp Thr Ser Ala Leu Ala Ala Gly Ser Ser Gln Glu Val Thr
 50 55 60
 15 Tyr Ala Gln Leu Asp His Trp Ala Leu Thr Gln Arg Thr Ala Arg Ala
 65 70 75 80
 Val Ser Pro Gln Ser Thr Lys Pro Met Ala Glu Ser Ile Thr Tyr Ala
 85 90 95
 20 Ala Val Ala Arg His
 100

25

(2) INFORMATION FOR SEQ ID NO: 251:

(i) SEQUENCE CHARACTERISTICS:

30 (A) LENGTH: 115 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 251:

35 Met Ser Pro His Pro Thr Ala Leu Leu Gly Leu Val Leu Cys Leu Ala
 1 5 10 15
 Gln Thr Ile His Thr Gln Glu Glu Asp Leu Pro Arg Pro Ser Ile Ser
 20 25 30
 40 Ala Glu Pro Gly Thr Val Ile Pro Leu Gly Ser His Val Thr Phe Val
 35 40 45
 Cys Arg Gly Pro Val Gly Val Gln Thr Phe Arg Leu Glu Arg Glu Ser
 50 55 60
 45 Arg Ser Thr Tyr Asn Asp Thr Glu Asp Val Ser Gln Ala Ser Pro Ser
 65 70 75 80
 Glu Ser Glu Ala Arg Phe Arg Ile Asp Ser Val Ser Glu Gly Asn Ala
 85 90 95
 50 Gly Pro Tyr Arg Cys Ile Tyr Tyr Lys Pro Pro Lys Trp Ser Glu Gln
 100 105 110
 55 Ser Asp Tyr
 115

60

(2) INFORMATION FOR SEQ ID NO: 252:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 18 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 252:

10 Thr Ala Leu Leu Gly Leu Val Leu Cys Leu Ala Gln Thr Ile His Thr
1 5 10 15
Gln Glu

- 15 (2) INFORMATION FOR SEQ ID NO: 253:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 14 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 253:

25 Leu Pro Arg Pro Ser Ile Ser Ala Glu Pro Gly Thr Val Ile
1 5 10

- 30 (2) INFORMATION FOR SEQ ID NO: 254:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 15 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 254:

35 Cys Arg Gly Pro Val Gly Val Gln Thr Phe Arg Leu Glu Arg Glu
1 5 10 15

- 40 (2) INFORMATION FOR SEQ ID NO: 255:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 31 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 255:

50 Val Leu Glu Arg Thr Ala Asp Lys Ala Thr Val Asn Gly Leu Pro Glu
1 5 10 15

55 Lys Asp Arg Glu Thr Asp Thr Ser Ala Leu Ala Ala Gly Ser Ser
20 25 30

- (2) INFORMATION FOR SEQ ID NO: 256:
60 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 438 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 256:

5 Met Asn Thr Pro Asn Gly Asn Ser Leu Ser Ala Ala Glu Leu Thr Cys
1 5 10 15

10 Gly Met Ile Met Cys Leu Ala Arg Gln Ile Pro Gln Ala Thr Ala Ser
20 25 30

Met Lys Asp Gly Lys Trp Glu Arg Lys Lys Phe Met Gly Thr Glu Leu
35 40 45

15 Asn Gly Lys Thr Leu Gly Ile Leu Gly Leu Gly Arg Ile Gly Arg Glu
50 55 60

Val Ala Thr Arg Met Gln Ser Phe Gly Met Lys Thr Ile Gly Tyr Asp
65 70 75 80

20 Pro Ile Ile Ser Pro Glu Val Ser Ala Ser Phe Gly Val Gln Gln Leu
85 90 95

25 Pro Leu Glu Glu Ile Trp Pro Leu Cys Asp Phe Ile Thr Val His Thr
100 105 110

Pro Leu Leu Pro Ser Thr Thr Gly Leu Leu Asn Asp Asn Thr Phe Ala
115 120 125

30 Gln Cys Lys Lys Gly Val Arg Val Val Asn Cys Ala Arg Gly Gly Ile
130 135 140

Val Asp Glu Gly Ala Leu Leu Arg Ala Leu Gln Ser Gly Gln Cys Ala
145 150 155 160

35 Gly Ala Ala Leu Asp Val Phe Thr Glu Glu Pro Pro Arg Asp Arg Ala
165 170 175

40 Leu Val Asp His Glu Asn Val Ile Ser Cys Pro His Leu Gly Ala Ser
180 185 190

Thr Lys Glu Ala Gln Ser Arg Cys Gly Glu Glu Ile Ala Val Gln Phe
195 200 205

45 Val Asp Met Val Lys Gly Lys Ser Leu Thr Gly Val Val Asn Ala Gln
210 215 220

Ala Leu Thr Ser Ala Phe Ser Pro His Thr Lys Pro Trp Ile Gly Leu
225 230 235 240

50 Ala Glu Ala Leu Gly Thr Leu Met Arg Ala Trp Ala Gly Ser Pro Lys
245 250 255

55 Gly Thr Ile Gln Val Ile Thr Gln Gly Thr Ser Leu Lys Asn Ala Gly
260 265 270

Asn Cys Leu Ser Pro Ala Val Ile Val Gly Leu Leu Lys Glu Ala Ser
275 280 285

60 Lys Gln Ala Asp Val Asn Leu Val Asn Ala Lys Leu Leu Val Lys Glu

337

290 295 300

Ala Gly Leu Asn Val Thr Thr Ser His Ser Pro Ala Ala Pro Gly Glu
305 310 315 320

5 Gln Gly Phe Gly Glu Cys Leu Leu Ala Val Ala Leu Ala Gly Ala Pro
 325 330 335

10 Tyr Gln Ala Val Gly Leu Val Gln Gly Thr Thr Pro Val Leu Gln Gly
 340 345 350

Leu Asn Gly Ala Val Phe Arg Pro Glu Val Pro Leu Arg Arg Asp Leu
355 360 365

15 Pro Leu Leu Leu Phe Arg Thr Gln Thr Ser Asp Pro Ala Met Leu Pro
370 375 380

Thr Met Ile Gly Leu Leu Ala Glu Ala Gly Val Arg Leu Leu Ser Tyr
385 390 395 400

20 Gln Thr Ser Leu Val Ser Asp Gly Glu Thr Trp His Val Met Gly Ile
 405 410 415

Ser Ser Leu Leu Pro Ser Leu Glu Ala Trp Lys Gln His Val Thr Glu
25 420 425 430

Ala Phe Gln Phe His Phe
435

30

(2) INFORMATION FOR SEQ ID NO: 257:

- 35 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 24 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 257:

40 Met Ala Phe Ala Asn Leu Arg Lys Val Leu Ile Ser Asp Ser Leu Asp
 1 5 10 15

Pro Cys Cys Arg Lys Ile Leu Gln
 20

45

(2) INFORMATION FOR SEQ ID NO: 258:

- 50 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 18 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 258:

55 Gly Gly Leu Gln Val Val Glu Lys Gln Asn Leu Ser Lys Glu Glu Leu
 1 5 10 15

Ile Ala
 60

5 (2) INFORMATION FOR SEQ ID NO: 259:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 29 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 259:

Met Cys Leu Ala Arg Gln Ile Pro Gln Ala Thr Ala Ser Met Lys Asp
1 5 10 15

15 Gly Lys Trp Glu Arg Lys Lys Phe Met Gly Thr Glu Leu
20 25

20 (2) INFORMATION FOR SEQ ID NO: 260:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 29 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 260:

Ala Leu Thr Ser Ala Phe Ser Pro His Thr Lys Pro Trp Ile Gly Leu
1 5 10 15

30 Ala Glu Ala Leu Gly Thr Leu Met Arg Ala Trp Ala Gly
20 25

35 (2) INFORMATION FOR SEQ ID NO: 261:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 36 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 261:

45 Glu Val Pro Leu Arg Arg Asp Leu Pro Leu Leu Leu Phe Arg Thr Gln
1 5 10 15

Thr Ser Asp Pro Ala Met Leu Pro Thr Met Ile Gly Leu Leu Ala Glu
20 25 30

50 Ala Gly Val Arg
35

55 (2) INFORMATION FOR SEQ ID NO: 262:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 109 amino acids

(B) TYPE: amino acid

60 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 262:

5 Phe Gly Thr Arg Phe Leu Ala Asn Leu Leu Leu Glu Glu Asp Asn Lys
 1 5 10 15
 Phe Cys Ala Asp Cys Gln Ser Lys Gly Pro Arg Trp Ala Ser Trp Asn
 20 25 30
 10 Ile Gly Val Phe Ile Cys Ile Arg Cys Ala Xaa Ile His Arg Asn Leu
 35 40 45
 Gly Val His Ile Ser Arg Val Lys Ser Val Asn Leu Asp Gln Trp Thr
 50 55 60
 15 Gln Val Gln Ile Gln Cys Met Gln Xaa Met Gly Asn Gly Lys Ala Asn
 65 70 75 80
 Arg Leu Tyr Glu Ala Tyr Leu Pro Glu Thr Phe Arg Arg Pro Gln Ile
 85 90 95
 20 Asp Pro Ala Val Glu Gly Phe Ile Arg Asp Xaa Tyr Glu
 100 105

25

(2) INFORMATION FOR SEQ ID NO: 263:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 amino acids
 30 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 263:

35 Glu Glu Asp Asn Lys Phe Cys Ala Asp Cys Gln Ser Lys Gly Pro Arg
 1 5 10 15
 Trp Ala Ser Trp Asn
 20

40

(2) INFORMATION FOR SEQ ID NO: 264:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 amino acids
 45 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 264:

50 Gly Val Phe Ile Cys Ile Arg Cys Ala Xaa Ile His Arg Asn Leu Gly
 1 5 10 15
 Val His Ile Ser
 20

55

(2) INFORMATION FOR SEQ ID NO: 265:

60 (i) SEQUENCE CHARACTERISTICS:

340

(A) LENGTH: 23 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 265:

5

Ser Val Asn Leu Asp Gln Trp Thr Gln Val Gln Ile Gln Cys Met Gln
 1 5 10 15

10

Xaa Met Gly Asn Gly Lys Ala
 20

15

(2) INFORMATION FOR SEQ ID NO: 266:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 245 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 266:

Met Asp Leu Leu Gly Leu Asp Ala Pro Val Ala Cys Ser Ile Ala Asn
 1 5 10 15

25

Ser Lys Thr Ser Asn Thr Leu Glu Lys Asp Leu Asp Leu Leu Ala Ser
 20 25 30

30

Val Pro Ser Pro Ser Ser Ser Gly Ser Arg Lys Val Val Gly Ser Met
 35 40 45

Pro Thr Ala Gly Ser Ala Gly Ser Val Pro Glu Asn Leu Asn Leu Phe
 50 55 60

35

Pro Glu Pro Gly Ser Lys Ser Glu Glu Ile Gly Lys Lys Gln Leu Ser
 65 70 75 80

Lys Asp Ser Ile Leu Ser Leu Tyr Gly Ser Gln Thr Xaa Gln Met Pro
 85 90 95

40

Thr Gln Ala Met Phe Met Ala Pro Ala Gln Met Ala Tyr Pro Thr Ala
 100 105 110

Tyr Pro Ser Phe Pro Gly Val Thr Pro Pro Asn Ser Ile Met Gly Ser
 115 120 125

45

Met Met Pro Pro Pro Val Gly Met Val Ala Gln Pro Gly Ala Ser Gly
 130 135 140

50

Met Val Ala Pro Met Ala Met Pro Ala Gly Tyr Met Gly Gly Met Gln
 145 150 155 160

Ala Ser Met Met Gly Val Pro Asn Gly Met Met Thr Thr Gln Gln Ala
 165 170 175

55

Gly Tyr Met Ala Gly Met Ala Ala Met Pro Gln Thr Val Tyr Gly Val
 180 185 190

Gln Pro Ala Gln Gln Leu Gln Trp Asn Leu Thr Gln Met Thr Gln Gln
 195 200 205

60

341

Met Ala Gly Met Asn Phe Tyr Gly Ala Asn Gly Met Met Asn Tyr Gly
 210 215 220

5 Gln Ser Met Ser Gly Gly Asn Gly Gln Ala Ala Asn Gln Thr Leu Ser
 225 230 235 240

Pro Gln Met Trp Lys
 245

10

(2) INFORMATION FOR SEQ ID NO: 267:

(i) SEQUENCE CHARACTERISTICS:
 15 (A) LENGTH: 315 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 267:

20 Met Asp Leu Leu Gly Leu Asp Ala Pro Val Ala Cys Ser Ile Ala Asn
 1 5 10 15

Ser Lys Thr Ser Asn Thr Leu Glu Lys Asp Leu Asp Leu Leu Ala Ser
 20 25 30

25 Val Pro Ser Pro Ser Ser Ser Gly Ser Arg Lys Val Val Gly Ser Met
 35 40 45

30 Pro Thr Ala Gly Ser Ala Gly Ser Val Pro Glu Asn Leu Asn Leu Phe
 50 55 60

Pro Glu Pro Gly Ser Lys Ser Glu Glu Ile Gly Lys Lys Gln Leu Ser
 65 70 75 80

35 Lys Asp Ser Ile Leu Ser Leu Tyr Gly Ser Gln Thr Xaa Gln Met Pro
 85 90 95

Thr Gln Ala Met Phe Met Ala Pro Ala Gln Met Ala Tyr Pro Thr Ala
 100 105 110

40 Tyr Pro Ser Phe Pro Gly Val Thr Pro Pro Asn Ser Ile Met Gly Ser
 115 120 125

Met Met Pro Pro Pro Val Gly Met Val Ala Gln Pro Gly Ala Ser Gly
 45 130 135 140

Met Val Ala Pro Met Ala Met Pro Ala Gly Tyr Met Gly Gly Met Gln
 145 150 155 160

50 Ala Ser Met Met Gly Val Pro Asn Gly Met Met Thr Thr Gln Gln Ala
 165 170 175

Gly Tyr Met Ala Gly Met Ala Ala Met Pro Gln Thr Val Tyr Gly Val
 180 185 190

55 Gln Pro Ala Gln Gln Leu Gln Trp Asn Leu Thr Gln Met Thr Gln Gln
 195 200 205

Met Ala Gly Met Asn Phe Tyr Gly Ala Asn Gly Met Met Asn Tyr Gly
 60 210 215 220

342

Gln Ser Met Ser Gly Gly Asn Gly Gln Ala Ala Asn Gln Thr Leu Ser
 225 230 235 240

5 Pro Gln Met Trp Lys Phe Gly Thr Arg Phe Leu Ala Asn Leu Leu Leu
 245 250 255

Glu Glu Asp Asn Lys Phe Cys Ala Asp Cys Gln Ser Lys Gly Pro Arg
 260 265 270

10 Trp Ala Ser Trp Asn Ile Gly Val Phe Ile Cys Ile Arg Cys Ala Xaa
 275 280 285

Ile His Arg Asn Leu Gly Val His Ile Ser Arg Val Lys Ser Val Asn
 15 290 295 300

Leu Asp Gln Trp Thr Gln Val Gln Ile Gln Cys
 305 310 315

20

(2) INFORMATION FOR SEQ ID NO: 268:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 39 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 268:

30 Met Gln Xaa Met Gly Asn Gly Lys Ala Asn Arg Leu Tyr Glu Ala Tyr
 1 5 10 15

Leu Pro Glu Thr Phe Arg Arg Pro Gln Ile Asp Pro Ala Val Glu Gly
 20 25 30

35 Phe Ile Arg Asp Xaa Tyr Glu
 35

40

(2) INFORMATION FOR SEQ ID NO: 269:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 67 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 269:

50 Lys Tyr Gly Lys Val Gly Lys Cys Val Ile Phe Glu Ile Pro Gly Ala
 1 5 10 15

Pro Asp Asp Glu Ala Val Arg Ile Phe Leu Glu Phe Glu Arg Val Glu
 20 25 30

55 Ser Ala Ile Lys Ala Val Val Asp Leu Asn Gly Arg Tyr Phe Gly Gly
 35 40 45

Arg Val Val Lys Ala Cys Phe Tyr Asn Leu Asp Lys Phe Arg Val Leu
 50 55 60

60

Asp Leu Ala
65

5

(2) INFORMATION FOR SEQ ID NO: 270:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 270:

Lys Ala Val Asp Leu Gly Arg Tyr Phe Gly Gly Arg
15 1 5 10

20

(2) INFORMATION FOR SEQ ID NO: 271:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 271:

Glu Ala Val Arg Ile Phe Phe Arg Glu
1 5

30

(2) INFORMATION FOR SEQ ID NO: 272:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 306 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 272:

40 Arg Met Gly Arg Phe His Arg Ile Leu Glu Pro Gly Leu Asn Ile Leu
1 5 10 15

Ile Pro Val Leu Asp Arg Ile Arg Tyr Val Gln Ser Leu Lys Glu Ile
20 25 30

45

Val Ile Asn Val Pro Glu Gln Ser Ala Val Thr Leu Asp Asn Val Thr
35 40 45

50

Leu Gln Ile Asp Gly Val Leu Tyr Leu Arg Ile Met Asp Pro Tyr Lys
50 55 60

Ala Ser Tyr Gly Val Glu Asp Pro Glu Tyr Ala Val Thr Gln Leu Ala
65 70 75 80

55

Gln Thr Thr Met Arg Ser Glu Leu Gly Lys Leu Ser Leu Asp Lys Val
85 90 95

60

Phe Arg Glu Arg Glu Ser Leu Asn Ala Ser Ile Val Asp Ala Ile Asn
100 105 110

344

Gln Ala Ala Asp Cys Trp Gly Ile Arg Cys Leu Arg Tyr Glu Ile Lys
 115 120 125
 5 Asp Ile His Val Pro Pro Arg Val Lys Glu Ser Met Gln Met Gln Val
 130 135 140
 Glu Ala Glu Arg Arg Lys Arg Ala Thr Val Leu Glu Ser Glu Gly Thr
 145 150 155 160
 10 Arg Glu Ser Ala Ile Asn Val Ala Glu Gly Lys Lys Gln Ala Gln Ile
 165 170 175
 Leu Ala Ser Glu Ala Glu Lys Ala Glu Gln Ile Asn Gln Ala Ala Gly
 180 185 190
 15 Glu Ala Ser Ala Val Leu Ala Lys Ala Lys Ala Lys Ala Glu Ala Ile
 195 200 205
 Arg Ile Leu Ala Ala Ala Leu Thr Gln His Asn Gly Asp Ala Ala Ala
 210 215 220
 Ser Leu Thr Val Ala Glu Gln Tyr Val Ser Ala Phe Ser Lys Leu Ala
 225 230 235 240
 25 Lys Asp Ser Asn Thr Ile Leu Leu Pro Ser Asn Pro Gly Asp Val Thr
 245 250 255
 Ser Met Val Ala Gln Ala Met Gly Val Tyr Gly Ala Leu Thr Lys Ala
 260 265 270
 30 Pro Val Pro Gly Thr Pro Asp Ser Leu Ser Ser Gly Ser Ser Arg Asp
 275 280 285
 Val Gln Gly Thr Asp Ala Ser Leu Asp Glu Glu Leu Asp Arg Val Lys
 290 295 300
 Met Ser
 305

40

(2) INFORMATION FOR SEQ ID NO: 273:

45

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 273:

50

Ala Ser Tyr Gly Val Glu Asp Pro Glu Tyr Ala Val Thr Gln Leu Ala
 1 5 10 15

Gln Thr Thr Met Arg Ser Glu Leu Gly Lys
 20 25

55

(2) INFORMATION FOR SEQ ID NO: 274:

60

(i) SEQUENCE CHARACTERISTICS:

345

(A) LENGTH: 27 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 274:

5

Met Gln Met Gln Val Glu Ala Glu Arg Arg Lys Arg Ala Thr Val Leu
 1 5 10 15

10

Glu Ser Glu Gly Thr Arg Glu Ser Ala Ile Asn
 20 25

15

(2) INFORMATION FOR SEQ ID NO: 275:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 275:

Leu Thr Val Ala Glu Gln Tyr Val Ser Ala Phe Ser Lys Leu Ala Lys
 1 5 10 15

25

Asp Ser Asn Thr Ile Leu Leu Pro Ser Asn
 20 25

30

(2) INFORMATION FOR SEQ ID NO: 276:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 70 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 276:

Leu Leu Gly Ala Thr Ala Pro Leu Val Ser Leu Val Pro Glu Val Ala
 1 5 10 15

40

Ala Ala Val Gly Asn Ala Gly Ala Arg Gly Ala Xaa His Trp Gly Pro
 20 25 30

45

Phe Ala Glu Gly Leu Ser Thr Gly Phe Trp Pro Arg Ser Ala Arg Ala
 35 40 45

Ser Ser Gly Leu Pro Arg Asn Thr Val Val Leu Phe Val Pro Gln Gln
 50 55 60

50

Glu Ala Trp Val Val Glu
 65 70

55

(2) INFORMATION FOR SEQ ID NO: 277:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 46 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 277:

5 Arg Met Trp Arg Asn Gly Thr His Phe Trp Glu Cys Lys Ile Val Gln
1 5 10 15
Pro Leu Trp Lys Thr Val Trp Trp Phe Pro Arg Lys Leu Ser Ile Glu
20 25 30
10 Leu Pro Glu Asn Leu Ala Ile Leu Ile Gly Thr Tyr Phe Lys
35 40 45

(2) INFORMATION FOR SEQ ID NO: 278:

15

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 33 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 278:

Leu Lys Arg His Phe Pro Lys Glu Ala Asn Lys His Val Lys Arg Cys
1 5 10 15
25 Ser Thr Ser Leu Asp Ile Arg Glu Ile Gln Ile Lys Ile Lys Met Arg
20 25 30
Tyr

30

(2) INFORMATION FOR SEQ ID NO: 279:

35

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 328 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 279:

Gly Thr Arg Pro Gly Glu Ser His Ala Asn Asp Leu Glu Cys Ser Gly
1 5 10 15
45 Lys Gly Lys Cys Thr Thr Lys Pro Ser Glu Ala Thr Phe Ser Cys Thr
20 25 30
Cys Glu Glu Gln Tyr Val Gly Thr Phe Cys Glu Glu Tyr Asp Ala Cys
35 40 45
50 Gln Arg Lys Pro Cys Gln Asn Asn Ala Ser Cys Ile Asp Ala Asn Glu
50 55 60
Lys Gln Asp Gly Ser Asn Phe Thr Cys Val Cys Leu Pro Gly Tyr Thr
65 70 75 80
55 Gly Glu Leu Cys Gln Ser Lys Ile Asp Tyr Cys Ile Leu Asp Pro Cys
85 90 95
60 Arg Asn Gly Ala Thr Cys Ile Ser Ser Leu Ser Gly Phe Thr Cys Gln
100 105 110

347

Cys Pro Glu Gly Tyr Phe Gly Ser Ala Cys Glu Glu Lys Val Asp Pro
 115 120 125
 5 Cys Ala Ser Ser Pro Cys Gln Asn Asn Gly Thr Cys Tyr Val Asp Gly
 130 135 140
 Val His Phe Thr Cys Asn Cys Ser Pro Gly Phe Thr Gly Pro Thr Cys
 145 150 155 160
 10 Ala Gln Leu Ile Asp Phe Cys Ala Leu Ser Pro Cys Ala His Gly Thr
 165 170 175
 Cys Arg Ser Val Gly Thr Ser Tyr Lys Cys Leu Cys Asp Pro Gly Tyr
 180 185 190
 15 His Gly Leu Tyr Cys Glu Glu Glu Tyr Asn Glu Cys Leu Ser Ala Pro
 195 200 205
 20 Cys Leu Asn Ala Ala Thr Cys Arg Asp Leu Val Asn Gly Tyr Glu Cys
 210 215 220
 Val Cys Leu Ala Glu Tyr Lys Gly Thr His Cys Glu Leu Tyr Lys Asp
 225 230 235 240
 25 Pro Cys Ala Asn Val Ser Cys Leu Asn Gly Ala Thr Cys Asp Ser Asp
 245 250 255
 Gly Leu Asn Gly Thr Cys Ile Cys Ala Pro Gly Phe Thr Gly Glu Glu
 260 265 270
 30 Cys Asp Ile Asp Ile Asn Glu Cys Asp Ser Asn Pro Cys His His Gly
 275 280 285
 35 Gly Ser Cys Leu Asp Gln Pro Asn Gly Tyr Asn Cys His Cys Pro His
 290 295 300
 Gly Trp Val Gly Ala Asn Cys Glu Ile His Leu Gln Trp Lys Ser Gly
 305 310 315 320
 40 His Met Ala Glu Ser Leu Thr Asn
 325

45

(2) INFORMATION FOR SEQ ID NO: 280:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 280:

50 Gly Lys Cys Thr Thr Lys Pro Ser Glu Ala Thr Phe Ser Cys Thr Cys
 1 5 10 15
 Glu Glu Gln Tyr Val Gly Thr Phe Cys
 20 25
 60

(2) INFORMATION FOR SEQ ID NO: 281:

5 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 281:

10 Cys Ala His Gly Thr Cys Arg Ser Val Gly Thr Ser Tyr Lys Cys Leu
 1 5 10 15

Cys Asp Pro Gly Tyr His
 20

15

(2) INFORMATION FOR SEQ ID NO: 282:

20 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 33 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 282:

25 Cys Ala Asn Val Ser Cys Leu Asn Gly Ala Thr Cys Asp Ser Asp Gly
 1 5 10 15

30 Leu Asn Gly Thr Cys Ile Cys Ala Pro Gly Phe Thr Gly Glu Glu Cys
 20 25 30

Asp

35

(2) INFORMATION FOR SEQ ID NO: 283:

40 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 299 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 283:

45 Met Ala Gln Asn Leu Lys Asp Leu Ala Gly Arg Leu Pro Ala Gly Pro
 1 5 10 15

Arg Gly Met Gly Thr Ala Leu Lys Leu Leu Gly Ala Gly Ala Val
 20 25 30

50

Ala Tyr Gly Val Arg Glu Ser Val Phe Thr Val Glu Gly Gly His Arg
 35 40 45

55 Ala Ile Phe Phe Asn Arg Ile Gly Gly Val Gln Gln Asp Thr Ile Leu
 50 55 60

Ala Glu Gly Leu His Phe Arg Ile Pro Trp Phe Gln Tyr Pro Ile Ile
 65 70 75 80

60 Tyr Asp Ile Arg Ala Arg Pro Arg Lys Ile Ser Ser Pro Thr Gly Ser

349

85 90 95
 Lys Asp Leu Gln Met Val Asn Ile Ser Leu Arg Val Leu Ser Arg Pro
 100 105 110
 5 Asn Ala Gln Glu Leu Pro Ser Met Tyr Gln Arg Leu Gly Leu Asp Tyr
 115 120 125
 10 Glu Glu Arg Val Leu Pro Ser Ile Val Asn Glu Val Leu Lys Ser Val
 130 135 140
 Val Ala Lys Phe Asn Ala Ser Gln Leu Ile Thr Gln Arg Ala Gln Val
 145 150 155 160
 15 Ser Leu Leu Ile Arg Arg Glu Leu Thr Glu Arg Ala Lys Asp Phe Ser
 165 170 175
 Leu Ile Leu Asp Asp Val Ala Ile Thr Glu Leu Ser Phe Ser Arg Glu
 180 185 190
 20 Tyr Thr Ala Ala Val Glu Ala Lys Gln Val Ala Gln Gln Glu Ala Gln
 195 200 205
 Arg Ala Gln Phe Leu Val Glu Lys Ala Lys Gln Glu Gln Arg Gln Lys
 25 210 215 220
 Ile Val Gln Ala Glu Gly Glu Ala Glu Ala Ala Lys Met Leu Gly Glu
 225 230 235 240
 30 Ala Leu Ser Lys Asn Pro Gly Tyr Ile Lys Leu Arg Lys Ile Arg Ala
 245 250 255
 Ala Gln Asn Ile Ser Lys Thr Ile Ala Thr Ser Gln Asn Arg Ile Tyr
 260 265 270
 35 Leu Thr Ala Asp Asn Leu Val Leu Asn Leu Gln Asp Glu Ser Phe Thr
 275 280 285
 Arg Gly Ser Asp Ser Leu Ile Lys Gly Lys Lys
 40 290 295

45 (2) INFORMATION FOR SEQ ID NO: 284:

 (i) SEQUENCE CHARACTERISTICS:

 (A) LENGTH: 18 amino acids

 (B) TYPE: amino acid

 (D) TOPOLOGY: linear

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 284:

Lys Ala Leu Ala Leu Ser Phe His Gly Trp Ser Gly Thr Gly Lys Asn
 1 5 10 15

55 Phe Val

60 (2) INFORMATION FOR SEQ ID NO: 285:

- 5 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 285:

10 Asn Leu Ile Asp Tyr Phe Ile Pro Phe Leu Pro Leu Glu Tyr Arg His
1 5 10 15
Val Arg Leu Cys Ala Arg
20

- 15 (2) INFORMATION FOR SEQ ID NO: 286:

- 20 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 286:

25 Asn Leu Ile Asp Tyr Phe Ile Pro Phe Leu Pro Leu Glu Tyr Arg His
1 5 10 15
Val Arg Leu Cys
20

- 30 (2) INFORMATION FOR SEQ ID NO: 287:

- 35 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 26 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 287:

40 Cys His Gln Thr Leu Phe Ile Phe Asp Glu Ala Glu Lys Leu His Pro
1 5 10 15
Gly Leu Leu Glu Val Leu Gly Pro His Leu
20 25

- 45 (2) INFORMATION FOR SEQ ID NO: 288:

- 50 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 288:

55 Pro Glu Lys Ala Leu Ala Leu Ser Phe His Gly Trp Ser Gly Thr Gly
1 5 10 15
Lys Asn Phe Val Ala
60 20

5 (2) INFORMATION FOR SEQ ID NO: 289:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 289:

Asn Leu Lys Glu Lys Ile Phe Ile Ser Phe Ala Trp Leu Pro Lys Ala
1 5 10 15

15 Thr Val Gln Ala Ala Ile Gly
20

20 (2) INFORMATION FOR SEQ ID NO: 290:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 290:

Trp Leu Pro Lys Ala Thr Val Gln Ala Ala Ile Gly Ser Val Ala Leu
1 5 10 15

30 Asp

35 (2) INFORMATION FOR SEQ ID NO: 291:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 18 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 291:

His Asp Arg Thr Met Gln Asp Ile Val Tyr Lys Leu Val Pro Gly Leu
45 1 5 10 15

Gln Glu

50 (2) INFORMATION FOR SEQ ID NO: 292:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 23 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 292:

60 Phe Ala Ser His Asp Arg Thr Met Gln Asp Ile Val Tyr Lys Leu Val

352

1 5 10 15
Pro Gly Leu Gln Glu Gly Glu
20

5

(2) INFORMATION FOR SEQ ID NO: 293:

10

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 293:

Leu Val Leu Ser Leu Gly Ala Trp Gly Trp Pro Ser Thr Cys Leu Trp
1 5 10 15

20

Trp

25

(2) INFORMATION FOR SEQ ID NO: 294:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 294:

Gln Gly Lys Leu Gln Met Trp Val Asp Val Phe Pro Lys Ser Leu
1 5 10 15

35

(2) INFORMATION FOR SEQ ID NO: 295:

40

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 295:

45

Pro Pro Phe Asn Ile Thr Pro Arg Lys Ala Lys Lys Tyr Tyr Leu Arg
1 5 10 15

50

(2) INFORMATION FOR SEQ ID NO: 296:

55

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 19 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 296:

Lys Thr Asp Val His Tyr Arg Ser Leu Asp Gly Glu Gly Asn Phe Asn
1 5 10 15

Trp Arg Phe

5

(2) INFORMATION FOR SEQ ID NO: 297:

10

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 297:

Pro Arg Leu Ile Ile Gln Ile Trp Asp Asn Asp Lys Phe Ser Leu Asp
1 5 10 15

20

Asp Tyr Leu Gly Phe Leu Glu Leu Asp Leu
20 25

25

(2) INFORMATION FOR SEQ ID NO: 298:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 298:

Ala Val Met Ile Gly Asp Asp Cys Arg Asp Asp Val Gly Gly Ala
1 5 10 15

35

(2) INFORMATION FOR SEQ ID NO: 299:

40

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 299:

45

Ile Leu Val Lys Thr Gly Lys Tyr Arg Ala Ser Asp Glu Glu Lys Ile
1 5 10 15

50

Asn

(2) INFORMATION FOR SEQ ID NO: 300:

55

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 277 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 300:

354

Met Asp Ser Met Pro Glu Pro Ala Ser Arg Cys Leu Leu Leu Leu Pro
 1 5 10 15
 5 Leu Leu Leu Leu Leu Leu Leu Leu Leu Pro Ala Pro Glu Leu Gly Pro
 20 25 30
 Ser Gln Ala Gly Ala Glu Glu Asn Asp Trp Val Arg Leu Pro Ser Lys
 35 40 45
 10 Cys Glu Val Cys Lys Tyr Val Ala Val Glu Leu Lys Lys Pro Leu Arg
 50 55 60
 Lys Arg Gln Asp Thr Glu Val Ile Gly Thr Val Tyr Gly Ile Leu Asp
 15 65 70 75 80
 Gln Lys Ala Ser Gly Val Lys Tyr Thr Lys Ser Asp Leu Arg Leu Ile
 85 90 95
 20 Glu Val Thr Glu Thr Ile Cys Lys Arg Leu Leu Asp Tyr Ser Leu His
 100 105 110
 Lys Glu Arg Thr Gly Ser Xaa Arg Phe Ala Lys Gly Met Ser Glu Thr
 115 120 125
 25 Phe Glu Thr Leu His Xaa Leu Val His Lys Gly Val Lys Val Val Met
 130 135 140
 Asp Ile Pro Tyr Glu Leu Trp Asn Glu Thr Ser Ala Glu Val Ala Asp
 145 150 155 160
 Leu Lys Lys Gln Cys Asp Val Leu Val Glu Glu Phe Glu Glu Val Ile
 165 170 175
 35 Glu Asp Trp Tyr Arg Asn His Gln Glu Glu Asp Leu Thr Glu Phe Leu
 180 185 190
 Cys Ala Asn His Val Leu Lys Gly Lys Asp Thr Ser Cys Leu Ala Glu
 195 200 205
 40 Gln Trp Ser Gly Lys Lys Gly Asp Thr Ala Ala Leu Gly Gly Lys Lys
 210 215 220
 Ser Lys Lys Lys Ser Ile Arg Ala Lys Ala Ala Gly Gly Arg Ser Ser
 45 225 230 235 240
 Ser Ser Lys Gln Arg Lys Glu Leu Gly Gly Leu Glu Gly Asp Pro Ser
 245 250 255
 50 Pro Glu Glu Asp Glu Gly Ile Gln Lys Ala Ser Pro Leu Thr His Ser
 260 265 270
 Pro Pro Asp Glu Leu
 275
 55

(2) INFORMATION FOR SEQ ID NO: 301:

60 (i) SEQUENCE CHARACTERISTICS:

355

(A) LENGTH: 199 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 301:

5 Met Asp Gly Gln Lys Lys Asn Trp Lys Asp Lys Val Val Asp Leu Leu
1 5 10 15

10 Tyr Trp Arg Asp Ile Lys Lys Thr Gly Val Val Phe Gly Ala Ser Leu
20 25 30

Phe Leu Leu Leu Ser Leu Thr Val Phe Ser Ile Val Ser Val Thr Ala
35 40 45

15 Tyr Ile Ala Leu Ala Leu Leu Ser Val Thr Ile Ser Phe Arg Ile Tyr
50 55 60

Lys Gly Val Ile Gln Ala Ile Gln Lys Ser Asp Glu Gly His Pro Phe
65 70 75 80

20 Arg Ala Tyr Leu Glu Ser Glu Val Ala Ile Ser Glu Glu Leu Val Gln
85 90 95

Lys Tyr Ser Asn Ser Ala Leu Gly His Val Asn Cys Thr Ile Lys Glu
100 105 110

Leu Arg Arg Leu Phe Leu Val Asp Asp Leu Val Asp Ser Leu Lys Phe
115 120 125

30 Ala Val Leu Met Trp Val Phe Thr Tyr Val Gly Ala Leu Phe Asn Gly
130 135 140

Leu Thr Leu Leu Ile Leu Ala Leu Ile Ser Leu Phe Ser Val Pro Val
145 150 155 160

35 Ile Tyr Glu Arg His Gln Ala Gln Ile Asp His Tyr Leu Gly Leu Ala
165 170 175

Asn Lys Asn Val Lys Asp Ala Met Ala Lys Ile Gln Ala Lys Ile Pro
180 185 190

Gly Leu Lys Arg Lys Ala Glu
195

45

(2) INFORMATION FOR SEQ ID NO: 302:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 302:

55 Met Ala Val Thr Leu Ser Leu Leu Leu Gly Gly Arg Val Cys Ala
1 5 10 15

60

(2) INFORMATION FOR SEQ ID NO: 303:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 41 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 303:

Pro Ser Leu Ala Val Gly Ser Arg Pro Gly Gly Trp Arg Ala Gln Ala
 1 5 10 15
 Leu Leu Ala Gly Ser Arg Thr Pro Ile Pro Thr Gly Ser Arg Arg Asn
 20 25 30
 Gly Ser Cys Arg Arg Trp Arg Ala Pro
 35 40

(2) INFORMATION FOR SEQ ID NO: 304:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 56 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 304:

Met Ala Val Thr Leu Ser Leu Leu Leu Gly Gly Arg Val Cys Ala Pro
 1 5 10 15
 Ser Leu Ala Val Gly Ser Arg Pro Gly Gly Trp Arg Ala Gln Ala Leu
 20 25 30
 Leu Ala Gly Ser Arg Thr Pro Ile Pro Thr Gly Ser Arg Arg Asn Gly
 35 40 45
 Ser Cys Arg Arg Trp Arg Ala Pro
 50 55

(2) INFORMATION FOR SEQ ID NO: 305:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 481 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 305:

GATGTTACAC AGCTCTTTAA TAATAGTGGC CATAGCTGTA ATAACAATGA CAACAGTAGG 60
 TAACGGTAGT CATAACCAACA GTAGGGCAGT GCATTTTATA TTACAACCTGG TTTCTTGCTC 120
 TAGTAGGCTT GGGGATGGGT GAAGACGGAC AGGGCTGGCG CAGACCCTTT CCTTCTCCTC 180
 TCCAGCCAC AGTGATCTGG GCTTTTACAA GACAGCCTGC TTCCATTGAG TAGTGTGGGA 240
 AAGTTCCTTC TTGGCTTAGC AATACCCCTG AGACCTTGTT CAGTGGGCTG TGTCTCTCCC 300

357

TGGGATGCTG GGAGCACCAA GTGTGGCCGA GCTAGGGCTG CTGACTTCCT CTGGGCGCCT 360
CTGGGCTGCG AGGGTCTCTT ATAGGAATTG AGGCCCTTTG CTGCTCCAAG AAATGCTGAG 420
5 GCTGTGGGCA RAGGGKTGTA CCAAGGGGA CTCITGCTCT GTGTCTGACT TTGGGGRATC 480
C 481

10

(2) INFORMATION FOR SEQ ID NO: 306:

(i) SEQUENCE CHARACTERISTICS:
15 (A) LENGTH: 58 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 306:

CACAGCTCTT TAATAATAGT GGCCATAGCT GTAATAACAA TGACAACAGT AGGTAACG 58

25

(2) INFORMATION FOR SEQ ID NO: 307:

(i) SEQUENCE CHARACTERISTICS:
30 (A) LENGTH: 59 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 307:

TGTGTCTCTC CCTGGGATGC TGGGAGCACC AAGTGTGGCC GAGCTAGGGC TGCTGACTT 59

40

(2) INFORMATION FOR SEQ ID NO: 308:

(i) SEQUENCE CHARACTERISTICS:
45 (A) LENGTH: 85 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double
(D) TOPOLOGY: linear

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 308:

GCGAGGGTCT CTTATAGGAA TTGAGGCCCT TTGCTGCTCC AAGAAATGCT GAGGCTGTGG 60

55

GCARAGGGKT GTACCCAAGG GGACT 85

60

(2) INFORMATION FOR SEQ ID NO: 309:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 34 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 309:

Met Val Gly Pro Val Thr Leu His Lys Lys Ile His Thr Thr Thr Val
 1 5 10 15

10 Leu Phe Ile Val Gln Ile His Ile Leu Leu Ile Gln Ala Ile Thr Gln
 20 25 30

Ala Lys

15

(2) INFORMATION FOR SEQ ID NO: 310:

20 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 67 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 310:

Leu Gln Met His Leu Met Ile Leu Gln Met Thr Gly Leu Ser Ile Leu
 1 5 10 15

30 Ala Leu Leu Gly Lys Ser Thr Thr Thr Ile Val Glu Gln Lys Phe His
 20 25 30

Asn Gly Lys Asn Gln Lys Ser Gly Leu Lys Glu Asn Arg Asp Lys Lys
 35 40 45

35 Lys Gln Thr Arg Trp Gln Ser Thr Ala Ser Gln Lys Ile Gly Ile Thr
 50 55 60

Glu Glu Arg
 65

40

(2) INFORMATION FOR SEQ ID NO: 311:

45 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 101 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

50 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 311:

Met Val Gly Pro Val Thr Leu His Lys Lys Ile His Thr Thr Thr Val
 1 5 10 15

55 Leu Phe Ile Val Gln Ile His Ile Leu Leu Ile Gln Ala Ile Thr Gln
 20 25 30

Ala Lys Leu Gln Met His Leu Met Ile Leu Gln Met Thr Gly Leu Ser
 35 40 45

60 Ile Leu Ala Leu Leu Gly Lys Ser Thr Thr Thr Ile Val Glu Gln Lys

359

50 55 60

Phe His Asn Gly Lys Asn Gln Lys Ser Gly Leu Lys Glu Asn Arg Asp
 65 70 75 80

5 Lys Lys Lys Gln Thr Arg Trp Gln Ser Thr Ala Ser Gln Lys Ile Gly
 85 90 95

10 Ile Thr Glu Glu Arg
 100

15 (2) INFORMATION FOR SEQ ID NO: 312:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 74 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 312:

Met Gln Thr Cys Pro Leu Val Gly Thr Leu Leu Thr Arg Asn Met Asp
 1 5 10 15

25 Gly Tyr Thr Cys Ala Val Val Thr Ser Thr Ser Phe Trp Ile Ile Ser
 20 25 30

Ala Trp Xaa Leu Trp Lys Gly Ser Pro Ser Thr Ser Met Pro Thr Met
 35 40 45

30 Pro Glu Thr Pro Leu Arg Thr Leu Cys Cys Thr Lys Met Pro Ser Ile
 50 55 60

Phe Ser Ser Leu Met Thr Asp Gly Arg Ala
 35 65 70

40 (2) INFORMATION FOR SEQ ID NO: 313:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 78 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 313:

Met Thr Leu Ile Gln Asn Cys Trp Tyr Ser Trp Leu Phe Phe Gly Phe
 1 5 10 15

50 Phe Phe His Phe Leu Arg Lys Ser Ile Ser Ile Phe Ser Ile Phe Leu
 20 25 30

Val Cys Phe Arg Ile Leu Ala Leu Gly Pro Thr Cys Phe Leu Val Trp
 35 40 45

55 Phe Trp Lys Ala Phe Phe Arg His Ile Leu Ile Phe Ile Cys Leu Ser
 50 55 60

Arg Glu Val Phe Arg Pro Arg Cys Phe Leu Val Tyr Phe Arg
 60 65 70 75

(2) INFORMATION FOR SEQ ID NO: 314:

5

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 71 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 314:

Met Gly Thr Arg Ala Gln Val Thr Pro Gly Arg Leu Pro Ile Pro Pro
 1 5 10 15

15

Pro Ala Pro Gly Leu Pro Phe Ser Ala Xaa Glu Pro Leu Gln Gly Gln
 20 25 30

Leu Arg Arg Val Ser Ser Ser Arg Gly Gly Phe Pro Gly Leu Ala Leu
 35 40 45

20

Gln Leu Leu Arg Ser Glu Thr Val Lys Ala Tyr Val Asn Asn Glu Ile
 50 55 60

25

Asn Ile Leu Ala Ser Phe Phe
 65 70

(2) INFORMATION FOR SEQ ID NO: 315:

30

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 40 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 315:

Met Leu Val Arg Thr Arg Pro Ser Gln Pro Leu Pro Leu Pro Gly Val
 1 5 10 15

40

Gly Leu Gly Gly Pro Arg Ser Gly Asp Pro Pro Glu Ser Thr Glu Leu
 20 25 30

Arg Lys Gly Pro Gly Phe Leu Ala
 35 40

45

(2) INFORMATION FOR SEQ ID NO: 316:

50

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 262 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 316:

Met Cys Pro Val Cys Gly Arg Ala Leu Ser Ser Pro Gly Ser Leu Gly
 1 5 10 15

60

Arg His Leu Leu Ile His Ser Glu Asp Gln Arg Ser Asn Cys Ala Val
 20 25 30

Cys Gly Ala Arg Phe Thr Ser His Ala Thr Phe Asn Ser Glu Lys Leu
 35 40 45
 5 Pro Glu Val Leu Asn Met Glu Ser Leu Pro Thr Val His Asn Glu Gly
 50 55 60
 Pro Ser Ser Ala Glu Gly Lys Asp Ile Ala Phe Ser Pro Pro Val Tyr
 65 70 75 80
 10 Pro Ala Gly Ile Leu Leu Val Cys Asn Asn Cys Ala Ala Tyr Arg Lys
 85 90 95
 Xaa Leu Glu Ala Gln Thr Pro Ser Val Xaa Lys Trp Ala Leu Arg Arg
 100 105 110
 15 Gln Asn Glu Pro Leu Glu Val Arg Leu Gln Arg Leu Glu Arg Glu Arg
 115 120 125
 Thr Ala Lys Lys Ser Arg Arg Asp Asn Glu Thr Pro Glu Glu Arg Glu
 130 135 140
 Val Arg Arg Met Arg Asp Arg Glu Ala Lys Arg Leu Gln Arg Met Gln
 145 150 155 160
 25 Glu Thr Asp Glu Gln Arg Ala Arg Arg Leu Gln Arg Asp Arg Glu Ala
 165 170 175
 Met Arg Leu Lys Arg Ala Asn Glu Thr Pro Glu Lys Arg Gln Ala Arg
 180 185 190
 30 Leu Ile Arg Glu Arg Glu Ala Lys Arg Leu Lys Arg Arg Leu Glu Lys
 195 200 205
 Met Asp Met Met Leu Arg Ala Gln Phe Gly Gln Asp Pro Ser Ala Met
 210 215 220
 Ala Ala Leu Ala Ala Glu Met Asn Phe Phe Gln Leu Pro Val Ser Gly
 225 230 235 240
 40 Val Glu Leu Asp Xaa Gln Leu Leu Gly Lys Met Ala Phe Glu Glu Gln
 245 250 255
 Asn Ser Ser Xaa Leu His
 260
 45

(2) INFORMATION FOR SEQ ID NO: 317:

50

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 190 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

55

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 317:

Met Asp His Ser His His Met Gly Met Ser Tyr Met Asp Ser Asn Ser
 1 5 10 15
 60 Thr Met Gln Pro Ser His His His Pro Thr Thr Ser Ala Ser His Ser

35 (2) INFORMATION FOR SEQ ID NO: 318:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 123 amino acids

(B) TYPE: amino acid

40 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 318:

	Met	Val	Gln	Pro	Cys	Gly	Ala	Cys	Ala	Lys	Thr	Xaa	Trp	Lys	Ala	Cys
45	1				5					10					15	
	Ser	Ser	Cys	Cys	Ser	Ser	Pro	Cys	Cys	Leu	Gln	Glu	Arg	Trp	Pro	Xaa
				20					25					30		
50	Pro	Xaa	Ala	Xaa	Cys	Pro	Glu	Xaa	Gly	Pro	Ser	Ser	His	Pro	Gly	Ile
			35						40					45		
	Gln	Ala	Leu	Cys	Ala	Val	Ala	Val	Val	Tyr	Leu	Ser	Pro	Ser	Ser	Arg
		50					55					60				
55	Leu	Asp	Trp	Ser	Leu	Ala	Pro	Leu	Phe	Val	Pro	Ser	Leu	Ala	Ala	Gly
	65					70					75					80
	Glu	Thr	Pro	Leu	Thr	Gln	Pro	Ala	Trp	Ala	Leu	Thr	Thr	Asn	Thr	Leu
60					85					90					95	

363

Gly His Gly Gln Pro Ala Gln Asp Arg Leu Pro Ala Leu Gly His Cys
 100 105 110

5 Ala Pro Ile Ser Val Leu Gly Leu Gly Ser Ser
 115 120

364

Applicant's or agent's file reference number	008PCT	International application ?	Unassigned
--	--------	-----------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page 75, line N/A	
B. IDENTIFICATION OF DEPOSIT	
Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit April 28, 1997	Accession Number 209012
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")	
For receiving Office use only	
<input checked="" type="checkbox"/> This sheet was received with the international application	For International Bureau use only
Authorized officer Lydel Meadows Paralegal Specialist IAPD-PCT Operations (703) 305-3745	<input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer

365

Applicant's or agent's file reference number	008PCT	International application number	Unassigned
--	--------	----------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page 75, line N/A	
B. IDENTIFICATION OF DEPOSIT	
Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit June 5, 1997	Accession Number 209089
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer Lydell Meadows Paralegal Specialist IAPD-PCT Operations (703) 305-3745	Authorized officer

366

Applicant's or agent's file reference number	2008PCT	International application	Unassigned
---	---------	---------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page 78, line N/A	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit June 5, 1997	Accession Number 209090
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")	

<p>For receiving Office use only</p> <p><input checked="" type="checkbox"/> This sheet was received with the international application</p> <p>Authorized officer Lydell Meadows Paralegal Specialist IAPD-PCT Operations (703) 305-3745</p>	<p>For International Bureau use only</p> <p><input type="checkbox"/> This sheet was received by the International Bureau on:</p> <p>Authorized officer</p>
---	--

Applicant's or agent's file reference number	008PCT	367	International application 1	Unassigned
---	--------	-----	-----------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page 80, line N/A	
B. IDENTIFICATION OF DEPOSIT	
Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit May 22, 1997	Accession Number 209076
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (If the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer Lydell Meadows Paralegal Specialist IAPD-PCT Operations (703) 305-3745	Authorized officer

368

Applicant's or agent's file reference number	008PCT	International application ?	Unassigned
--	--------	-----------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page 82, line N/A	
B. IDENTIFICATION OF DEPOSIT	
Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit May 29, 1997	Accession Number 209086
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input checked="checked" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer Lydell Meadows Paralegal Specialist IAPD-PCT Operations (703) 305-3745	Authorized officer

369

Applicant's or agent's file reference number	008PCT	International application ?	Unassigned
--	--------	-----------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page 83, line N/A	
B. IDENTIFICATION OF DEPOSIT	
Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit June 19, 1997	Accession Number 209126
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications, e.g., "Accession Number of Deposit")	
For receiving Office use only	For International Bureau use only
<input checked="" type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer Lydell Meadows Paralegal Specialist IAPD-PCT Operations (703) 305-3745	Authorized officer

What Is Claimed Is:

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:
 - (a) a polynucleotide fragment of SEQ ID NO:X or a polynucleotide fragment of the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - (b) a polynucleotide encoding a polypeptide fragment of SEQ ID NO:Y or a polypeptide fragment encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - (c) a polynucleotide encoding a polypeptide domain of SEQ ID NO:Y or a polypeptide domain encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - (d) a polynucleotide encoding a polypeptide epitope of SEQ ID NO:Y or a polypeptide epitope encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - (e) a polynucleotide encoding a polypeptide of SEQ ID NO:Y or the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X, having biological activity;
 - (f) a polynucleotide which is a variant of SEQ ID NO:X;
 - (g) a polynucleotide which is an allelic variant of SEQ ID NO:X;
 - (h) a polynucleotide which encodes a species homologue of the SEQ ID NO:Y;
 - (i) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(h), wherein said polynucleotide does not hybridize under stringent conditions to a nucleic acid molecule having a nucleotide sequence of only A residues or of only T residues.
2. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding a secreted protein.
3. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding the sequence identified as SEQ ID NO:Y or the polypeptide encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X.

4. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises the entire nucleotide sequence of SEQ ID NO:X or the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X.
5. The isolated nucleic acid molecule of claim 2, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.
6. The isolated nucleic acid molecule of claim 3, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.
7. A recombinant vector comprising the isolated nucleic acid molecule of claim 1.
8. A method of making a recombinant host cell comprising the isolated nucleic acid molecule of claim 1.
9. A recombinant host cell produced by the method of claim 8.
10. The recombinant host cell of claim 9 comprising vector sequences.
11. An isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence selected from the group consisting of:
- (a) a polypeptide fragment of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
 - (b) a polypeptide fragment of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z, having biological activity;
 - (c) a polypeptide domain of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
 - (d) a polypeptide epitope of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
 - (e) a secreted form of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
 - (f) a full length protein of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;

- (g) a variant of SEQ ID NO:Y;
- (h) an allelic variant of SEQ ID NO:Y; or
- (i) a species homologue of the SEQ ID NO:Y.

12. The isolated polypeptide of claim 11, wherein the secreted form or the
5 full length protein comprises sequential amino acid deletions from either the C-terminus
or the N-terminus.

13. An isolated antibody that binds specifically to the isolated polypeptide of
claim 11.
10

14. A recombinant host cell that expresses the isolated polypeptide of claim
11.

15. A method of making an isolated polypeptide comprising:
15 (a) culturing the recombinant host cell of claim 14 under conditions such that
said polypeptide is expressed; and
(b) recovering said polypeptide.

16. The polypeptide produced by claim 15.
20

17. A method for preventing, treating, or ameliorating a medical condition,
comprising administering to a mammalian subject a therapeutically effective amount of
the polypeptide of claim 11 or the polynucleotide of claim 1.

18. A method of diagnosing a pathological condition or a susceptibility to a
25 pathological condition in a subject comprising:
(a) determining the presence or absence of a mutation in the polynucleotide of
claim 1; and
(b) diagnosing a pathological condition or a susceptibility to a pathological
30 condition based on the presence or absence of said mutation.

19. A method of diagnosing a pathological condition or a susceptibility to a
pathological condition in a subject comprising:
(a) determining the presence or amount of expression of the polypeptide of
35 claim 11 in a biological sample; and
(b) diagnosing a pathological condition or a susceptibility to a pathological
condition based on the presence or amount of expression of the polypeptide.

20. A method for identifying a binding partner to the polypeptide of claim 11 comprising:

- 5 (a) contacting the polypeptide of claim 11 with a binding partner; and
(b) determining whether the binding partner effects an activity of the polypeptide.

21. The gene corresponding to the cDNA sequence of SEQ ID NO:Y.

10 22. A method of identifying an activity in a biological assay, wherein the method comprises:

- (a) expressing SEQ ID NO:X in a cell;
(b) isolating the supernatant;
(c) detecting an activity in a biological assay; and
15 (d) identifying the protein in the supernatant having the activity.

23. The product produced by the method of claim 22.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/12125

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : 435/69.1, 70.1, 71.1, 235.1, 243, 325, 410; 536/23.1, 23.5

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/69.1, 70.1, 71.1, 235.1, 243, 325, 410; 536/23.1, 23.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 679 016 A1 (MATSUBARA et al.) 11 February 1995, see entire document and sequence listing, especially SEQ ID NO. 12, position 585-605 versus reference sequence at position 42-62; SEQ ID NO. 13, position 1942-5189 versus reference sequence at position 1-248; SEQ ID NO. 15, position 569-817 versus reference sequence at position 1-249; SEQ ID NO. 16, position 233-586 versus reference sequence at position 1-354; and SEQ ID NO. 18, position 1309-1699 versus reference sequence at position 12-393.	1-10, 14, 15, and 21
Y	WO 96/40917 A1 (YALE UNIVERSITY.) 19 December 1996. See entire document and sequence listing, especially SEQ ID NO. 11, position 444-692 versus reference sequence at position 2-250.	1-10, 14, 15, and 21

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

08 SEPTEMBER 1998

Date of mailing of the international search report

01 OCT 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

BRIAN R. SPANTON

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/12125

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 95/27791 A1 (DAVIES et al.) 19 October 1995, See entire document and sequence listing, especially SEQ ID NO. 17, position 742-799 versus reference sequence at position 1334-1391.	1-10, 14, 15, and 21
Y	WO 95/14100 A1 (THE WELLCOME FOUNDATION LIMITED) 26 May 1995. See entire document and sequence listing, especially SEQ ID NO. 97, position 966-991 versus reference sequence at position 747-772.	1-10, 14, 15, 21
Y	WO 94/28133 A1 (AMGEN INC.) 08 December 1994, see entire document and sequence listing, especially SEQ ID NO. 14, position 758-808 versus reference sequence at position 1599-1649.	1-10, 14, 15, and 21
Y	WO 95/01437 A2 (REGENTS OF THE UNIVERSITY OF MINESOTA) 12 January 1995, see entire document and sequence listing, especially SEQ ID NO. 19, position 69-122 versus reference sequence at position 604-657.	1-10, 14, 15, and 21

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/12125

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-10, 14 15 and 21

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/12125

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

C07H 21/02, 04; C12N 5/00, 5/04, 5/06, 5/10, 5/16; 15/00, 15/09, 15/10, 15/11, 15/12; C12P 21/04, 21/06

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

Databases: Genbank, embase, biosis, medline

Search Terms/Strategy: Sequence search of Sequences 11-19 and 97; est; secret?; moore?/au; shi?/au; rosen?/au; ruben?/au; lafleuer?/au; olsen?/au; ebner?/au; brewer?/au; young?/au; greene?/au; ferrie?/au; yu ?/au; ni ?/au; feng ?/au

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I:

Claims 1-10, 14, 15, and 21 drawn to a polynucleotide(s), vector(s) containing the polynucleotide, host cells containing the vector(s) which are SEQ ID NO: X or a polynucleotide encoding the polypeptide Y or a cDNA in the material deposited with American Type Culture Collection with accession number Z wherein the cDNA in Z hybridizes to X. Additionally Group I contains the first method making the cells (claim 14) containing the vector(s) containing the polynucleotide(s) and the first method of use of the cells (claim 15) to make a product. There appear to be a total of 46 polynucleotide sequences of which the first ten (10) are selected for examination and therefore, there are nine (9) remaining additional groups of four (4) polynucleotide sequences.

Group II:

Claims 11, 12, 16, and 23 drawn to polypeptides and/or fragments thereof with the amino acid sequence defined by SEQ ID NO: Y as found in the material deposited with the American Type Culture Collection with accession number Z. There appear to be a total of 74 polypeptide sequences and therefore 73 additional species of proteins.

Group III:

Claim 13, drawn to an antibody that binds to a polypeptide with the amino acid sequence defined by SEQ ID NO: Y as found in the material deposited with the American Type Culture Collection with accession number Z. There appear to be a total of 74 antibodies that correspond to the SEQ ID NOs: for the "Y" and "Z" sequences and therefore 73 additional species of proteins.

Group IV:

Claim 17, drawn to a process of preventing, treating, or ameliorating a medical condition by administering a polypeptide or a polynucleotide which a second/alternative process of use of the second product and of an alternative process of use of the first claimed product in Group I.

In Group IV, and where additional fees are paid, the claims are searched only insofar as they are applicable to the selected polypeptide and its corresponding SEQ ID NO: as the first species is directed to a process practiced using a polypeptide. The second species is the practice of the process using a polynucleotide. In each instance, the same selected polypeptide as for the first species of Group II and for the first 10 polynucleotide sequences for Group I would be examined. Applicant may elect to pay additional fees for each additional one of the 73 different polypeptide species beyond the first one (1) polypeptide and/or the first 10 polynucleotides as set forth in the above paragraphs directed to Group I and II.

Group V:

Claim 18, drawn to a method of diagnosis of a pathological condition and another alternative process of use of the first claimed product in Group I. Additionally Group V contains indications that there are a total of 46 polynucleotide sequences and therefore, nine (9) additional groups of four (4) polynucleotide sequences beyond the first ten (10) sequences.

Group VI:

Claim 19, drawn to a method of diagnosis of a pathological condition and another alternative process of use of the polypeptide. There appear to be a total of 74 polypeptide sequences and therefore 73 additional species of proteins.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/12125

Group VII:

Claim 20, drawn to a method of identification of a binding partner for a polypeptide. There appear to be a total of 74 polypeptide sequences and therefore 73 additional species of proteins.

Group VIII:

Claim 22, drawn to a method of identification of function of a protein is another alternative process of use of the product in Group I. Additionally Group V contains indications that there are a total of 46 polynucleotide sequences and therefore, nine(9) additional groups of four (4) polynucleotide sequences beyond the first ten (10) sequences.

The inventions listed as Groups I through VIII do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons.

Claims of Group I are drawn to nucleotides, nucleotide constructs, and/or methods requiring the use of nucleotides or nucleotide constructs that contain more than ten individual, independent, and distinct nucleotide sequences in alternative form. Accordingly, these claims are subject to lack of unity as outlined in 1192 O.G. 68 (19 November 1996).

For Group I, the first ten (10) of the individual polynucleotide sequences designated as "X" by SEQ ID NO: as set forth in the application (see for example page 29+ and/or the SEQUENCE LISTING) are included for search. The corresponding SEQ ID NO: for "Y" and "Z" for each selected "X" should also be noted. The search of the no more than ten sequences may include the complements of the selected sequences and, where appropriate, may include subsequences within the selected sequences (e.g., oligomeric probes and/or primers).

In Group IV (as directed to the species which are polynucleotides) should applicant pay the additional fee for the second appearing species in Group IV which are polynucleotides, first ten (10) of the individual polynucleotide sequences designated as "X" by SEQ ID NO: as set forth in the application (see for example page 29+ and/or the SEQUENCE LISTING) are included for search of Group IV should the fees for Group IV be paid. This is also applied to Groups V and VIII. The corresponding SEQ ID NO: for "Y" and "Z" for each selected "X" should also be noted. The search of the no more than ten sequences may include the complements of the selected sequences and, where appropriate, may include subsequences within the selected sequences (e.g., oligomeric probes and/or primers).

Where Applicant may elect to pay additional fees for a search of sequences beyond the initial ten (10) polynucleotide sequences, and in accordance with 1192 O.G. 68 (19 November 1996), applicant may select additional groups of polynucleotides consisting of four (4) sequences beyond the initial ten (10) sequences for Group I which would then be searched with Group I upon payment of the requisite fees for the requisite Groups beyond Group I.

As to the polypeptides of Groups II, III, IV (as directed to a species which is a polypeptide), VI, and VII each is a distinct and different protein. Should additional fees for the above indicated Groups be paid, the first amino acid sequence identified from the SEQUENCE LISTING by applicant would be searched with the additional group for which the additional search fees were paid.

Applicant may select additional proteins and or antibodies to be searched by specifying the appropriate SEQ ID NOs and payment of the requisite additional fees for each single additional particular species that are selected beyond the one (1) protein identified by SEQ ID NO:.

The SEQ ID NOs in Group I define, absent evidence to the contrary, structurally distinct and different proteins. Note the present application written description (page 5+) refers to the protein encoded by gene 1 as likely to be involved in promotion of a variety of cancers whereas gene 2 (pages 6-7) is directed to apparently a variety but not correlated immune system disorder(s) whereas gene 3 (pages 7-8) is asserted at page 7 to be a mediator of ligand dependent AF-2. Each of which and absent factual evidence to the contrary, are directed to genes encoding distinct and different proteins and are therefore distinct and different genes and appear to map to different chromosomes.

As to the protein of Group II and the antibody of Group III, each is distinct and different for the reasons indicated in the preceding paragraph and because the proteins have distinct and different chemical, physical, and biological properties from that of DNA/polynucleotides/vectors and cells containing same.

Groups IV through VIII are directed to alternative processes of use of the Group I and II compositions where